



UNITED STATES DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
NATIONAL MARINE FISHERIES SERVICE
Alaska Fisheries Science Center
Resource Assessment and Conservation Engineering Division
7600 Sand Point Way Northeast
BIN C15700, Building 4
Seattle, Washington 98115-0070

April 9, 1997

NOAA SHIP MILLER FREEMAN
Cruise No. 97-02
Preliminary Cruise Results
Echo Integration-Trawl Survey of the Southeastern
Aleutian Basin near Bogoslof Island

Midwater Assessment and Conservation Engineering

Scientists from the Alaska Fisheries Science Center (AFSC) investigated the population distribution and characteristics of spawning walleye pollock (*Theragra chalcogramma*) in the southeastern Aleutian Basin near Bogoslof Island between February 28 and March 11, 1997, (12 sea days) using echo integration-trawl (EIT) survey techniques aboard the NOAA ship *Miller Freeman*. This research cruise was the ninth in a series that began in 1988 and has been continued annually with the exception of 1990 as part of long-term monitoring of Bering Sea walleye pollock. In 1997, the cruise began and ended in Dutch Harbor, Alaska, and was a cooperative effort involving scientists from the United States, China, Poland and South Korea.

The vessel's itinerary was as follows:

Feb 28	Embark scientists during touch and go in Dutch Harbor, standard sphere calibration in Captain's Bay, AK.
Mar 1-10	Echo integration-trawl survey of the Bogoslof Island region
Mar 11	Arrive Dutch Harbor; end of cruise

The primary cruise objectives were to collect echo integration data and midwater trawl data essential to determine the distribution, biomass, and biological composition of walleye pollock in the southeastern Aleutian Basin. The 38 kHz and 120 kHz scientific acoustic systems were calibrated using standard sphere techniques. Physical oceanographic data (temperature and



salinity profiles) were collected at selected sites. Sea surface parameters (e.g., temperature, salinity, light level) were monitored continuously, and an acoustic Doppler current profiler was operated continuously in the water profiling mode.

Secondary objectives involved a number of separate projects and investigators from the AFSC, University of Alaska, and the Alaska Department of Fish and Game (ADF&G).

METHODS

Sampling Equipment

Acoustic data were collected with a Simrad EK500¹ quantitative echo-sounding system on board the NOAA ship *Miller Freeman*, a 66-m (216 ft) stern trawler equipped for fisheries and oceanographic research. The Simrad 38 kHz and 120 kHz split-beam transducers were mounted on the bottom of the vessel's centerboard. With the centerboard fully extended, the transducers were 9 m below the water surface. System electronics were housed inside the vessel in a new permanent laboratory space dedicated to acoustics that replaced the portable laboratory used on previous cruises. Data from the Simrad EK500 echo sounder/receiver were processed using Simrad BI500 echo integration and target strength data analysis software on a SUN workstation. Results presented in this document are based on the 38 kHz data.

Midwater echo sign was sampled using an Aleutian Wing 30/26 trawl (AWT), a full mesh wing trawl constructed of nylon except for polyethylene towards the aft section of the body and the codend. Headrope and footrope lengths each measured 81.7 m (268 ft) and mesh sizes tapered from 325.1 cm (128 in) in the forward section of the net to 8.9 cm (3.5 in) in the codend. The net was fitted with a 3.2 cm (1.25 in) codend liner. It was fished with 82.4 m (270 ft) of 1.9 cm (0.75 in) diameter 8x19 non-rotational dandyines, 454.5 kg (1,000-lb) tom weights on each side, and 5 m² "Fishbuster" doors [1,250 kg (2,750 lb)]. For one haul, 227.3 kg (500-lb) tom weights were used. Vertical net opening and depth were monitored with a WesMar third wire netsounder system attached to the headrope of the trawl.

Trawl hauls were made to identify selected echo sign and provide biological data and pollock samples. Haul duration was kept to

¹ Reference to trade names or commercial firms does not constitute U.S. Government endorsement.

the minimum necessary to ensure an adequate sample. Biological data collected from each haul included species composition, sex composition, length frequencies, whole fish and ovary weights, maturities, otoliths, pollock ovaries and whole stomachs from selected species.

Temperature/depth data were collected with a micro-bathythermograph (MBT) attached to the headrope of all trawls. Conductivity-temperature-depth (CTD) data were collected with a Seabird CTD system at calibration sites and selected locations. While transecting, we collected temperature-depth profile data at other locations using expendable bathythermographs (XBT's). Sea surface oceanographic data and environmental data were collected using the *Miller Freeman's* Scientific Collection System (SCS). Ocean current profile data were obtained using the vessel's acoustic Doppler current profiler system whose transducer is mounted in the centerboard.

Survey Methods

Four standard sphere calibrations of the acoustic systems were made in conjunction with the survey (Table 1). No significant differences were observed in the 38 kHz system parameters among the four calibrations. Results from calibration of the 120 kHz system are not presented here as that system was not used in the acoustic data analysis.

Between March 1 and 10, two EIT survey passes were run through the Bogoslof spawning area. Pass 1 (March 1-8) covered about 1400 nautical miles (nmi) of transects (Fig. 1). The trackline consisted of 27 north-south parallel transects beginning at about long. 165°44'W and ending near long. 170°20'W. Transect spacing at the eastern end was 10 nmi and decreased to 5 nmi between transects 7.0 and 17.0. Pass 2 (March 9-10) covered 525 nmi, and was run from west to east between longitudes 170°W and 167°30'W at 5 nmi spacing (Fig. 2). For both passes, southern transect endpoints were at approximately 100 m bottom depth on the Aleutian shelf but varied depending on bottom depth and fish echo sign. The northern extent of the 10 nmi-spaced transects was similar to that during previous winter surveys in the Bogoslof region, between latitudes 54°30' and 54°40'N east of long. 168°W and between latitudes 54°00' and 54°30'N west of long. 168°W.

Echo integration and trawl data were collected 24 hours a day. Vessel speeds averaged 11.5 kts during acoustic data collection and ranged between 7 and 12 kts, depending upon weather conditions. We collected echo integration data from 14 m below

the surface to within 0.5 m of the bottom, or to 1000 m depending on bottom depth. These data were thoroughly scrutinized by one or more scientists and stored in an INGRES database. When properly scaled they were used to provide estimates of pollock density.

Midwater trawl hauls were made at selected locations to identify echo sign and provide biological samples. The average trawling speed was about 3 kts. The AWT's vertical opening averaged about 32 m. Standard catch sorting and enumeration procedures were used to process all catches. Each trawl catch was completely sorted unless it exceeded about 1000 kg; representative splits of large catches were sorted. Total weights and numbers of individuals, by species, were determined for each catch.

Individual pollock were further sampled to determine sex, fork length (FL), body weight, age, maturity, and ovary weight. An electronic scale was used to determine all weights taken from individual pollock specimens. Fish FLs were determined to the nearest cm with a Polycorder measuring device (a combination of a bar code reader and a hand held computer). Since winter 1996, maturities have been determined by visual inspection of gonads based on an internationally accepted 8-stage scale. Expressed in terms of the older 5-stage scale, the stages were categorized as: immature, developing 1 & 2, pre-spawning 1 & 2, spawning, and post-spawning 1 & 2.

Several special projects were completed in addition to species collections associated with the estimation of pollock biomass. They included collecting and preserving stomachs from lanternfish (Myctophidae), and smoothtongues (Bathylagidae) in 10% formalin for food habits studies. Mature walleye pollock were spawned and the fertilized eggs cultured for laboratory experiments on eggs and larvae. Samples of pollock ovary tissue were collected from pre-spawning females for a study of interannual variation in fecundity. Fin clip, muscle and otolith samples were taken from walleye pollock for FOCI/ADF&G genetic research. Samples of forage fishes (myctophids) were frozen for a University of Alaska (Fairbanks) study on seabird food habits. Whole pollock were collected and frozen for the Observer Program at AFSC.

PRELIMINARY RESULTS

Biological data were collected and samples preserved from 16 midwater trawls (Figs. 1&2, Tables 2-4). Pollock dominated the catches in both weights and numbers (Table 5). Oceanographic data were collected from 19 MBT casts (Table 2), 3 CTD casts, and

5 XBT casts (Table 6). Temperature profiles showed a well-mixed water column with an average temperature of 3.9° C between the surface and 500 m. Near-surface temperatures averaged 3.2° C on the eastern edge of the survey area, and 3.7° C on the western (Aleutian Basin) side (Fig. 3).

During both passes, pollock distribution was similar to that in 1996. Pollock were moderately concentrated between longitudes 165°30'-166°30'W and in the southern portion of the survey area along the Aleutian Island chain between 167°30'-168°30'W (Fig 4). As in 1996, most pollock (60% of the biomass from pass 1) were concentrated in Samalga Pass, between Umnak Island and the Islands of Four Mountains (longitudes 169°-170° W, Figs. 4&5). They were distributed in several spawning aggregations 1-10 miles in horizontal extent and 200-400 m in vertical extent between 200-700 m in the water column. The average sizes of pollock increased from east to west. The eastern most haul (haul 1) caught pollock with lengths averaging 44.0 cm/41.9 cm for females/males, respectively (Fig. 6a). Between longitudes 166°W and 167° W, average lengths were 50.4 cm/46.8 cm for females/males (Fig. 6b). Lengths of pollock encountered west of long. 167°W averaged 55.4 cm/51.8 cm for females/males, respectively (Fig. 6c). Sampled lengths ranged from 33-66 cm for sexes combined. The sex ratio by haul spanned 0.29 to 0.96 female and averaged around 0.62 female. Maturities observed were mainly pre-spawning 1&2 (coded 4 and 5 in the current 8-point scale); 84% and 78% were pre-spawning among females/males, respectively (Fig. 7). The mean gonadosomatic index (gsi) for prespawning females was 0.20 (Fig. 8). Total pollock abundance in the Bogoslof area appeared to be lower than it was in 1996.

SCIENTIFIC PERSONNEL

<u>Name</u>	<u>Sex/ Nationality</u>	<u>Position</u>	<u>Organization</u>
Neal Williamson	M/USA	Chief Scientist	AFSC
Daniel Twohig	M/USA	Electronics Tech.	AFSC
Taina Honkalehto	F/USA	Fish. Biologist	AFSC
Steve de Blois	M/USA	Fish. Biologist	AFSC
Mike Guttormsen	M/USA	Fish. Biologist	AFSC
Kevin Landgraf	M/USA	Fish. Biologist	AFSC
Jerzy Janusz	M/Poland	Fish. Biologist	MIR
Seok-Gwan Choi	M/Korea	Fish. Acoustician	NFRDA
Xianyong Zhao	M/China	Fish. Acoustician	YSFRI

AFSC - Alaska Fisheries Science Center, Seattle, WA
MIR - Morski Intytut Rybacki, Gdynia, Poland
NFRDA - National Fisheries Research and Development Agency,
Pusan, Korea
YSFRI - Yellow Sea Fisheries Research Institute, Qingdao, China

For further information contact Dr. Gary Stauffer, Director,
Resource Assessment and Conservation Engineering Division,
Alaska Fisheries Science Center, National Marine Fisheries
Service, 7600 Sand Point Way Northeast, Building 4, BIN C15700,
Seattle, WA 98115-0070. Telephone (206) 526-4170.

Table 1. Summary of sphere calibrations conducted before, during and after the winter 1997 pollock echo integration-trawl survey of the Bogoslof Island region.

Date (1997)	Location	Freq (kHz)	Water Temp (deg. C)		Sphere Range from Transducer (m)	TS Gain (dB)	SV Gain (dB)	3dB Beam Width (deg.)	Angle Offset	
			at Transducer*	at Sphere					Along	Athwart
11 Feb	Port Susan, WA	38	7.5	8.2	31.8	27.2	27.1	6.81	-0.03	-0.06
28 Feb	Captains Bay, AK	38	3.5	3.8	31.5	27.1	27.1	6.80	0.00	-0.05
14 Mar	Nateekin Bay, AK	38	3.4	3.4	28.8	27.1	--	6.79	0.00	-0.03
26 Mar	Malina Bay, AK	38	3.8	3.8	26.6	27.1	27.1	6.71	-0.01	-0.06
Feb-Mar	System settings during surveys	38	--	--	--	27.1	27.1	6.70	-0.09	-0.02

* The transducer is located approximately 9 m below the water surface.

Note: Gain and beam pattern terms are defined in the "Operator Manual for Simrad EK500 Scientific Echo Sounder (1993)" available from Simrad Subsea A/S , Standpromenaden 50, P.O. Box 111 N-3191 Horten, Norway.

Table 2. Summary of midwater trawl stations and catch data from the winter 1997 pollock echo integration-trawl survey of the Bogoslof Island area.

Haul No.	Date (GMT)	Time (GMT)	Duration (minutes)	Start Position		Depth (m)		Temp. (deg. C)		MBT cast number	Pollock Catch		Other Catch	
				Latitude (N)	Longitude (W)	Gear	Bottom	Gear	Surface		kg	number	kg	number
1	1 Mar	15:47	5	54 24.81	165 43.67	144	267	4.0	2.6	4	1,404.6	2,416		
2	2 Mar	0:50	25	54 21.09	166 18.31	492	733	3.8	3.3	5	2,506.6	2,639	13.4	35
3	3 Mar	7:38	7	53 36.63	167 43.76	469	776	3.8	3.8	6	2,422.4	2,320	7.5	47
4	4 Mar	4:05	5	53 36.57	168 16.92	500	686	3.6	3.8	7	604.8	422	5.1	612
5	4 Mar	16:14	3	53 33.88	168 27.88	494	793	3.6	4.0	8	490.0	426	4.5	198
6	6 Mar	16:25	7	53 27.76	169 26.96	118	1190	3.9	3.9	9	5,000.0	3,423		
7	7 Mar	6:48	25	53 16.38	169 32.71	205	600	4.1	3.9	10	133.3	95	0.4	27
8	7 Mar	12:24	4	53 6.62	169 7.37	450	820	3.3	3.9	11	1,073.4	637	3.0	220
9	7 Mar	16:03	6	53 1.12	169 16.79	299	800	4.0	3.9	12	969.5	693	6.4	29
10	7 Mar	20:05	7	53 1.71	169 18.76	513	972	3.3	3.8	13	3,627.3	3,196	2.7	283
11	9 Mar	9:03	3	53 0.67	169 21.11	225	906	4.1	3.5	14	634.6	504	3.2	9
12	9 Mar	18:25	5	53 9.94	169 4.49	542	959	3.3	3.6	15	2,980.2	2,262	9.8	640
13	10 Mar	8:07	4	53 33.82	168 21.91	448	894	3.6	3.6	16	1,791.8	1,363	8.2	282
14	10 Mar	22:33	7	53 35.77	167 39.32	469	911	3.7	3.7	17	1,938.0	1,520	12.0	134
15	11 Mar	7:45	8	54 21.52	166 1.49	496	590	3.6	3.7	18	864.5	1,056	12.7	226
16	11 Mar	10:34	10	54 25.07	166 9.37	483	564	3.7	3.2	19	670.6	746	16.1	1,036

Table 3. Summary of pollock biological samples and measurements collected during the winter 1997 echo integration-trawl survey of the Bogoslof Island area.

Haul	Length ¹	Maturity	Otoliths	Fish Weight	Ovary Weight
1	334	78	78	78	10
2	388	67	67	67	26
3	296	72	72	72	36
4	265	63	63	63	8
5	336	59	59	59	36
6	319	60	60	38	34
7	95	95	95	95	77
8	277	62	62	62	49
9	320	77	77	61	41
10	394	58	58	58	16
11	387	69	69	69	25
12	351	63	63	63	0
13	331	100	50	50	0
14	375	100	75	75	24
15	286	98	98	98	49
16	312	100	100	100	49
Totals	5,066	1,221	1,146	1,108	480

¹ Length includes samples from random length frequency only

Table 4. Summary of biological samples collected for special projects during the winter 1997 pollock echo integration-trawl survey of the Bogoslof Island area.

Haul	Pollock Spawning (M/F)	Stomach Collection MYC ¹ /NST ²	Observ.Pro. Samples POL/RF ³	Frozen MYC ¹	Pollock Ovary Collection	Pollock #1 ⁴	Genetics #2 ⁵
1	-	-	Y	-	-	100	-
2	-	3/-	Y	-	26	-	-
3	-	-	Y	Y	-	-	20
4	-	14/5	Y	Y	8	100	-
5	-	12/-	Y	Y	1	-	20
6	-	-	Y	-	-	100	-
7	-	3/-	Y	-	-	-	-
8	-	10/-	-	Y	-	-	20
9	-	7/-	-	-	-	-	20
10	3/3	-	-	-	-	100	-
11	-	4/-	-	-	-	-	20
12	5/3	17/-	-	Y	-	-	-
13	-	10/1	-	Y	-	-	20
14	-	20/-	-	Y	-	-	-
15	-	15/-	-	-	1	-	-
16	-	20/-	-	Y	-	-	-
Total	8/6	120/6*	8 Boxes	~(15kg)	36	400	120

¹ Myctophidae

² Northern Smoothtongues

³ Pollock/Rockfish

⁴ #1-Fin clips only

⁵ #2-Muscle, fin clips, otoliths taken

*NST collection was limited due to poor condition after haul

Table 5. Summary of catch by species in 16 Aleutian Wing trawls during the 1997 echo integration-trawl survey of the Bogoslof Island area.

Common Name	Scientific Name	Weight (kg)	Weight (%)	Numbers
Walleye Pollock	<i>Theragra chalcogramma</i>	27,111.6	99.6	23,718
Lanternfish Unidentified	Myctophidae	39.5	0.1	2,810
Pacific Sleeper Shark	<i>Somniosus pacificus</i>	11.2	<0.1	1
Pacific Lamprey	<i>Lampetra tridentata</i>	10.6	<0.1	29
Squid Unidentified	Teuthoidea	9.9	<0.1	121
Smooth Lump sucker	<i>Aptocyclus ventricosus</i>	8.8	<0.1	5
Longnose Lancetfish	<i>Alepisaurus ferox</i>	6.5	<0.1	1
Northern Smoothtongue	<i>Leuroglossus schmidti</i>	3.4	<0.1	557
Chum Salmon	<i>Oncorhynchus keta</i>	2.5	<0.1	1
Magistrate Armhook Squid	<i>Berryteuthis magister</i>	2.4	<0.1	9
Arrowtooth Flounder	<i>Atheresthes stomias</i>	2.1	<0.1	2
Jellyfish Unidentified	Scyphozoa	2.1	<0.1	4
Chinook Salmon	<i>Oncorhynchus tshawytscha</i>	2.0	<0.1	1
Lump sucker Unidentified	Cyclopteridae	1.4	<0.1	1
Shortfin Eelpout	<i>Lycodes brevipes</i>	1.1	<0.1	4
Pacific Viperfish	<i>Chauliodus macouni</i>	0.4	<0.1	6
Shrimp Unidentified	Decapoda	0.4	<0.1	210
Eelpout Unidentified	Zoarcidae	0.2	<0.1	3
Pacific Herring	<i>Clupea pallasii</i>	0.2	<0.1	1
Slender Barracudina	<i>Lestidiops ringens</i>	0.1	<0.1	4
Barracudina Unidentified	Paralepididae	0.1	<0.1	4
Snailfish Unidentified	Cyclopteridae	<0.1	<0.1	1
Salps Unidentified	<i>Thaliacea unident.</i>	<0.1	<0.1	1
Ribbon Barracudina	<i>Notolepsis risso</i>	<0.1	<0.1	1
Invertebrate Unidentified		<0.1	<0.1	1
Totals		27,216.8		27,496

Table 6. Summary of conductivity-temperature-depth (CTD) and expendable bathythermograph (XBT) casts made during the winter 1997 echo integration-trawl survey of the Bogoslof Island area.

Cast type	Number	Date	Time (GMT)	Latitude (N)	Longitude (W)	Bottom depth (m)
CTD	1	11 Feb	21:25	48 8.36	122 25.11	109
CTD	2	1 Mar	2:49	53 50.88	166 35.21	61
CTD	3	1 Mar	11:03	54 19.18	166 0.15	514
XBT	1	3 Mar	5:05	53 47.84	167 43.77	>1,500
XBT	2	3 Mar	22:26	54 16.65	168 17.58	>1,500
XBT	3	4 Mar	11:09	53 48.50	168 26.16	>1,500
XBT	4	5 Mar	0:04	54 7.37	168 51.43	>1,500
XBT	5	5 Mar	3:21	53 41.71	169 25.18	>1,500




Figure 1. Pass 1 trackline and haul locations of the winter 1997 echo integration-trawl survey of the Bogoslof Island area.




Figure 2. Trackline and haul locations for pass 2 of the winter 1997 echo integration-trawl survey of the Bogoslof Island area.




Figure 3. Temperature profiles from micro-bathymographs (MBTs) attached to the headrope of trawls during the winter 1997 echo integration-trawl survey of pollock in the Bogoslof Island area. Representative profiles from (a) eastern and (b) western sections of the survey area.




Figure 4. Relative pollock density along trackline from pass 1 of the winter 1997 echo integration-trawl survey of the Bogoslof Island area. Transect numbers are indicated.




Figure 5. Relative pollock density along trackline from pass 2 of the winter 1997 echo integration-trawl survey of the Bogoslof Island area. Transect numbers are indicated.




Figure 6. Raw length frequencies from midwater trawls east of 166 deg. W (a), between 166 and 167 deg. W (b), and west of 167 deg. W (c) from the winter 1997 echo integration-trawl survey of the Bogoslof Island area.




Figure 7. Maturity stages of pollock observed during the winter 1997 echo integration-trawl survey of the Bogoslof Island area. Haul 1 (E of 166° W) is contrasted with hauls 2, 15, and 16 (166 - 167° W), and hauls 3-14 (W of 167° W).




Figure 8. Pollock gonado-somatic indices for mature females plotted as a function of length from the winter 1997 echo integration-trawl survey of the Bogoslof Island area.