

CHAPTER 10

OTHER FLATFISH

by

Paul D. Spencer, Gary E. Walters, and Thomas K. Wilderbuer

EXECUTIVE SUMMARY

The following changes have been made to this assessment relative to the November 2003 SAFE:

Changes in the input data

- 1) The 2003 catch (total and discarded) was updated, and catch through 2 October, 2004 were included in the assessment.
- 2) 2004 Eastern Bering Sea and Aleutian Islands trawl survey biomass estimates and standard errors of other flatfish species were included in the assessment. A linear regression between the EBS shelf biomass estimates and the AI survey estimates was used to predict the AI biomass in years where an AI survey did not occur.

Changes in assessment results

- 1) A summary of the 2004 assessment harvest recommendations relative to the 2003 assessment harvest recommendations is as follows:

	2004 Assessment recommendations	2003 Assessment recommendations
Exploitable biomass	142,610 t	97,327 t
ABC	21,391 t	13,549 t
Overfishing	28,522 t	18,065 t
F_{ABC}	0.15	0.15
$F_{\text{overfishing}}$	0.20	0.20

Introduction

The Bering Sea/Aleutian Islands “other flatfish” group have typically included those flatfish besides rock sole, yellowfin sole, arrowtooth flounder, and Greenland turbot. Flathead sole (*Hippoglossoides elassodon*) were part of the other flatfish complex until they were removed in 1995, and Alaska plaice was removed from the complex in 2002, as sufficient biological data exists for these species to construct age-structured population models. In contrast, survey biomass estimates are the principal data source used to assess the remaining other flatfish. Although over a dozen species (Table 10.1) of flatfish are found in the BSAI area, the other flatfish biomass consists primarily of starry flounder, rex sole, longhead dab, and butter sole.

Catch History

The miscellaneous species found in the other flatfish species category are listed in Table 10.1, and their catches from 1995-2004 are shown in Table 10.2. Catch from 1995-2002 were obtained from the NMFS Regional Office “blend” data, and the catch for some species are reported by species and in an aggregate flatfish group. The catch estimates for these years were produced by applying the proportional catch, by species, from fishery observer data to the estimated total catch for the aggregate other flatfish group, and adding this total to the catch that was reported by species. In the newer catch accounting system used in 2003 and 2004, catches of other flatfish are reported only in an aggregate group, and the catch estimates for these years were produced by applying the proportional catch, by species, from fishery observer data to the estimated total catch of the aggregate group. In recent years, starry flounder (*Platichthys stellatus*) and rex sole (*Glyptocephalus zachirus*) account for most of the harvest of other flatfish, and contributed 84% of the harvest of other flatfish in 2004.

Other flatfish are grouped with Alaska plaice, rock sole, and flathead sole and other flatfish fisheries in a single prohibited species class (PSC) classification, with seasonal and total annual allowances of prohibited bycatch applied to the classification. In recent years, this group of fisheries has been closed prior to attainment of the TAC due to the bycatch of halibut (Table 10.3). Additionally, beginning on August 14th, 2004, retention of other flatfish was prohibited due to the total allowable catch being attained. The total catch of other flatfish, as of October 2nd, 2004, was 4669 t, or approximately 168 % of the TAC level of 2775 t.

DATA

Absolute Abundance and Exploitation Rates

The biomass of the other flatfish complex on the eastern Bering Sea shelf has been relatively stable from 1983-1995, averaging 50,200 t, and has slightly increased from 1996 to 2004, averaging 81,900 t (Table 10.4). The 2004 biomass estimate of other flatfish on the EBS shelf is 127,630 t. Increases in biomass have also been seen in the Aleutian Islands trawl survey, and the 2004 estimates of 14,980 t is larger than any previous AI survey estimate. The 2004 BSAI estimate of total biomass is 142,610 t. An estimate of total BSAI biomass for the years in which an AI survey was not conducted was produced by regressing the AI survey biomass against the EBS survey biomass and adding the predicted AI biomass estimate to the observed EBS estimate. Individual species biomass estimates for the EBS and AI areas from 1997-2004 are shown in Table

10.5. Estimates of species biomass for starry flounder, rex sole, and butter sole in the Aleutian Islands were computed by fitting a linear trend to the observed survey data from 1991-2004, and using this trend to estimate biomass in years without an Aleutian Island survey. Estimate of total BSAI biomass (Table 10.6) were then used to compute species-specific exploitation rates.

Exploitation rates for starry flounder and rex sole have been low, not exceeding 0.10 from 1997 to 2003 (Table 10.6). The exploitation rates for butter sole have been slightly higher, exceeding 0.14 in 1997, 2000, 2001, and 2003, but the biomass estimates for butter sole have large sampling variances, with coefficients of variation ranging from 0.5 to 0.86 in recent EBS trawl surveys dating back to 1999.

The 2003 biomass estimate of butter sole of 429 t is less than one-fourth the 2002 estimate of 2382, and results in an estimated exploitation rate of nearly 70%. However, butter sole were only captured in four hauls in the 2003 EBS trawl survey, leading to the large coefficient of variation of 0.74 in the estimated biomass. In addition, the bulk of the 2003 fishery records come primarily from waters less than 50 m in January and February, a depth and time not covered by the trawl survey. Thus, it is likely that the population of butter sole is larger than that indicated from the survey, and the comparison of survey biomass to harvest should be interpreted accordingly. The 2004 biomass estimate of butter sole was 1076 t, more than twice the 2003 estimate, and has an associated CV of 0.86.

Several species of other flatfish are relatively rare on the EBS shelf, including Dover sole, Sakhalin sole, and English sole, and it is useful to identify whether the EBS represents the edge of the distribution for these species. The distribution of English sole has been identified as Baja California to Unimak Island, and the distribution of Dover sole has been identified as from Baja California to the Bering Sea (Hart 1973). Thus, the eastern Bering Sea can be considered the edge of the range for these species, and more significant populations of these species are found in the Gulf of Alaska. For example, the abundance of Dover sole in the 1984-2001 GOA surveys has fluctuated between 63,000 t and 96,000 t, the abundance of butter sole has fluctuated between 17,000 t and 30,000 t, and the abundance of English sole has fluctuated between 3,000 t and 14,000 t (Turnock et al. 2001). Dover sole and English sole were most common in the eastern portion of the GOA, consistent with their reported distribution along the west coast of North America.

PROJECTIONS AND HARVEST ALTERNATIVES

Reference Fishing Mortality Rates and Yields

Other flatfish are assessed under Tier 5 of Amendment 56 to the BSAI groundfish management plan, and thus require estimates of biomass and natural mortality. The natural mortality rates used in age-structured BSAI flatfish assessments can be used as guidance and are presented below:

Species	Natural mortality rate used for stock assessment
Yellowfin sole	0.12
Rock sole	0.18
Flathead sole	0.20
Alaska plaice	0.25

Given this range of values, an assumption of 0.20 appears reasonable. The estimates of F_{abc} and F_{ofl} under tier 5 are $0.75M$ and M , respectively, and the ABC and OFL levels are the product of the fishing mortality rate and the biomass estimate. Given the F_{abc} and F_{ofl} levels of 0.15 and 0.20, and the biomass estimate of 142,610 t, the resulting ABC and OFL levels are 21,391 and 28,522 t.

F level (value)	Projected yield for year 2005
Tier 5 F_{ABC} (0.15)	21,391 t
Tier 5 F_{OFL} (0.20)	28,522 t

Summary

In summary, several quantities pertinent to the management of the other flatfish are listed below.

Quantity	Value
M	0.20
Tier	5
Year 2004 Total Biomass	142,610 t
F_{OFL}	0.20
Maximum F_{ABC}	0.15
Recommended F_{ABC}	0.15
OFL	28,522 t
Maximum allowable ABC	21,391 t
Recommended ABC	21,391 t

REFERENCES

- Hart, J.L. 1973. Pacific fishes of Canada. Fisheries Research Board of Canada, Bulletin 180, Ottawa. 740 pp.
- Spencer, P.D., T.K. Wilderbuer, and C.I. Zhang. 2002. A mixed-species yield per recruit model for eastern Bering Sea flatfish fisheries. Can J. Fish. Aquat. Sci. 59:291-302.
- Turnock, B.J., T.K. Wilderbuer, and E.S. Brown. 2001. Gulf of Alaska flatfish. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Gulf of Alaska Region as Projected for 2002. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.

Table 10.1. Flatfish species of the Bering Sea/Aleutian Islands “other flatfish” management complex.

<u>Common Name</u>	<u>Scientific Name</u>
Arctic flounder	<i>Liopsetta glacialis</i>
butter sole	<i>Isopsetta isolepis</i>
curlfin sole	<i>Pleuronectes decurrens</i>
deepsea sole	<i>Embassichthys bathybus</i>
Dover sole	<i>Microstomus pacificus</i>
English sole	<i>Parophrys vetulus</i>
longhead dab	<i>Limanda proboscidea</i>
Pacific sanddab	<i>Citharichthys sordidus</i>
petrale sole	<i>Eopsetta jordani</i>
rex sole	<i>Glyptocephalus zachirus</i>
roughscale sole	<i>Clidodoerma asperrimum</i>
sand sole	<i>Psettichthys melanostictus</i>
slender sole	<i>Lyopsetta exilis</i>
starry flounder	<i>Platichthys stellatus</i>
Sakhalin sole	<i>Pleuronectes sakhalinensis</i>

Table 10.2. Harvest (t) of other flatfish from 1995-2004. 2004 data is through October 2, 2004.

Year	Starry Founder	Rex sole	Butter sole	Remaining Species	Total
1995	398	673	157	97	1324
1996	1171	1148	218	211	2748
1997	1043	687	448	312	2490
1998	402	998	229	136	1765
1999	725	998	230	164	2117
2000	1151	1069	458	349	3027
2001	755	869	244	160	2028
2002	1075	1192	222	142	2631
2003	887	1399	296	166	2749
2004	2062	1858	514	235	4669

Table 10.3. Restrictions on the “other flatfish” fishery from 1994 to 2004 in the Bering Sea – Aleutian Islands management area. Note that in 1994, the other flatfish category included flathead sole. Unless otherwise indicated, the closures were applied to the entire BSAI management area. Zone 1 consists of areas 508, 509, 512, and 516, whereas zone 2 consists of areas 513, 517, and 521.

Year	Dates	Bycatch Closure
1994	2/28 – 12/31	Red King crab cap (Zone 1 closed)
	5/7 – 12/31	Bairdi Tanner crab (Zone 2 closed)
	7/5 – 12/31	Annual halibut allowance
1995	2/21 – 3/30	First Seasonal halibut cap
	4/17 – 7/1	Second seasonal halibut cap
	8/1 – 12/31	Annual halibut allowance
1996	2/26 – 4/1	First Seasonal halibut cap
	4/13 – 7/1	Second seasonal halibut cap
	7/31 – 12/31	Annual halibut allowance
1997	2/20 – 4/1	First Seasonal halibut cap
	4/12 – 7/1	Second seasonal halibut cap
	7/25 – 12/31	Annual halibut allowance
1998	3/5 – 3/30	First Seasonal halibut cap
	4/21 – 7/1	Second seasonal halibut cap
	8/16 – 12/31	Annual halibut allowance
1999	2/26 – 3/30	First Seasonal halibut cap
	4/27 – 7/04	Second seasonal halibut cap
	8/31 – 12/31	Annual halibut allowance
2000	3/4 – 3/31	First Seasonal halibut cap
	4/30 – 7/03	Second seasonal halibut cap
	8/25 – 12/31	Annual halibut allowance
2001	3/20 – 3/31	First Seasonal halibut cap
	4/27 – 7/01	Second seasonal halibut cap
	8/24 – 12/31	Annual halibut allowance
2002	2/22 – 12/31	Red King crab cap (Zone 1 closed)
	3/1 – 3/31	First Seasonal halibut cap
	4/20 – 6/29	Second seasonal halibut cap
	7/29 – 12/31	Annual halibut allowance
2003	2/18 – 3/31	First Seasonal halibut cap
	4/1 – 6/21	Second seasonal halibut cap
	7/31 – 12/31	Annual halibut allowance
2004	2/24 – 3/31	First Seasonal halibut cap
	4/16 – 6/29	Second seasonal halibut cap
	6/4 – 8/13	Bycatch status
	8/14 – 12/31	Prohibited species status

Table 10.4. Estimated biomass (t) of other flatfish from the eastern Bering Sea and Aleutian Islands trawl survey. Species included are dover sole, longhead dab, rex sole, Sakhalin sole, starry flounder, and butter sole. A linear regression between EBS and AI survey abundance was used to predicted AI abundance in years in which an AI survey did not exist.

Year	Area		Total
	EBS	AI	
1982	117763		129518
1983	66131	2700	68831
1984	59647		64956
1985	34572		37101
1986	39517	6100	45617
1987	49764		53977
1988	43751		47298
1989	49592		53786
1990	46649		50517
1991	72399	2144	74543
1992	53817		58480
1993	44399		48017
1994	54045	5464	59509
1995	37786		40671
1996	60225		65599
1997	70225	7580	77805
1998	73936		80830
1999	67713		73917
2000	70538	8149	78687
2001	78844		86282
2002	98052	8801	106853
2003	90327		99039
2004	127630	14980	142610

Table 10.5 --Estimated biomass (t) and coefficient of variation (in parentheses) for the miscellaneous species of the "other flatfish" management complex in the Bering Sea trawl and Aleutian Islands surveys.

Eastern Bering Sea Shelf survey

Year	Species						
	Dover Sole	Rex Sole	longhead dab	Sakhalin sole	starry flounder	butter sole	English sole
1982	--	5994 (0.16)	103806 (0.16)	--	7781 (0.32)	182 (0.82)	--
1983	--	7272 (0.18)	51386 (0.38)	--	7436 (0.25)	37 (0.45)	--
1984	--	13058 (0.28)	35308 (0.16)	137 (0.43)	8913 (0.36)	2231 (0.64)	--
1985	10 (1.04)	10751 (0.20)	9107 (0.13)	102 (0.37)	12181 (0.24)	2421 (0.83)	--
1986	15 (1.00)	12886 (0.22)	10889 (0.14)	274 (0.48)	9112 (0.33)	6341 (0.58)	--
1987	81 (0.91)	12931 (0.19)	11897 (0.19)	110 (0.59)	22702 (0.63)	2043 (0.38)	--
1988	38 (0.59)	15445 (0.15)	16710 (0.19)	253 (0.63)	9222 (0.30)	2083 (0.47)	--
1989	--	12939 (0.15)	13086 (0.16)	58 (0.57)	22205 (0.35)	1304 (0.54)	--
1990	47 (0.58)	11857 (0.21)	18601 (0.15)	110 (0.51)	15048 (0.26)	986 (0.60)	--
1991	55 (0.70)	16014 (0.28)	18680 (0.14)	291 (0.79)	34303 (0.23)	3056 (0.50)	--
1992	137 (0.58)	14001 (0.24)	10827 (0.17)	75 (0.48)	27544 (0.22)	1233 (0.70)	--
1993	37 (0.75)	14567 (0.32)	11690 (0.21)	78 (0.34)	16510 (0.22)	1517 (0.75)	--
1994	73 (0.72)	15943 (0.38)	18533 (0.26)	183 (0.41)	18218 (0.22)	1095 (0.97)	--
1995	--	10420 (0.28)	8402 (0.15)	109 (0.32)	17652 (0.29)	1203 (0.54)	--
1996	--	10532 (0.40)	8567 (0.20)	34 (0.34)	40409 (0.45)	683 (0.53)	--
1997	--	8233 (0.27)	18003 (0.21)	87 (0.49)	41018 (0.21)	2884 (0.43)	--
1998	41 (0.44)	7588 (0.22)	14737 (0.19)	34 (0.49)	49605 (0.30)	1942 (0.38)	--
1999	16 (0.65)	8020 (0.28)	12087 (0.21)	63 (0.29)	43375 (0.25)	4152 (0.62)	--
2000	11 (1.02)	9348 (0.19)	13511 (0.30)	145 (0.88)	45810 (0.19)	1713 (0.56)	--
2001	16 (0.84)	21660 (0.23)	12764 (0.26)	31 (0.43)	43026 (0.25)	796 (0.50)	--
2002	7 (0.80)	26053 (0.20)	9740 (0.22)	7 (0.69)	59877 (0.23)	2254 (0.64)	--
2003	350 (0.66)	28023 (0.15)	8827(0.22)	55 (0.40)	52893 (0.17)	179 (0.61)	--
2004	31(0.51)	28762 (0.19)	11290 (0.23)	8 (0.64)	86698 (0.38)	841 (0.86)	--

Aleutian Islands Surveys

Year	Species						
	Dover Sole	Rex Sole	longhead dab	Sakhalin sole	starry flounder	butter sole	English sole
1991 AI	174 (0.45)	1694 (0.18)	--	--	142 (0.85)	86 (0.73)	47 (0.80)
1994 AI	438 (0.41)	4306 (0.15)	--	--	134 (0.69)	505 (0.98)	83 (0.81)
1997 AI	386 (0.34)	6378 (0.16)	--	--	459 (0.90)	346 (0.98)	12 (0.72)
2000 AI	630 (0.38)	6526 (0.18)	--	--	590 (0.71)	310 (0.99)	95 (0.97)
2002 AI	575 (0.28)	7381 (0.15)	--	--	671 (0.72)	127 (0.83)	47 (0.94)
2004 AI	870 (0.28)	13717 (0.18)	--	--	123 (0.72)	235 (0.93)	35 (1.00)

Table 10.6. Estimated exploitation rates of rex sole, starry flounder and butter sole from 1997 to 2004.

Year	Rex sole			Starry Flounder			Butter sole		
	Biomass (t)	Harvest (t)	Exp. Rate	Biomass (t)	Harvest (t)	Exp. Rate	Biomass (t)	Harvest (t)	Exp. Rate
1997	14611	401	0.03	41477	814	0.02	3230	336	0.10
1998	14250	569	0.04	49950	242	0.00	2210	157	0.07
1999	15415	516	0.03	43750	597	0.01	4416	167	0.04
2000	15874	569	0.04	46400	770	0.02	2023	266	0.13
2001	30524	507	0.02	43829	479	0.01	1059	147	0.14
2002	33411	1227	0.04	60633	1023	0.02	2382	187	0.08
2003	38349	1399	0.04	53353	887	0.02	429	296	0.69
2004	42479	1858	0.04	86821	2062	0.02	1076	514	0.48