

Assessment of Walleye Pollock in the Gulf of Alaska

by

Martin Dorn, Steven Barbeaux, Sarah Gaichas, Michael Guttormsen,
Bernard Megrey, Kally Spalinger, and Mark Wilkins

Summary of major changes

Relative to last year's assessment, the following changes have been made in the current assessment.

New Input data:

1. Fishery: 2003 total catch and catch at age.
2. Shelikof Strait EIT survey: 2004 biomass and age composition.
3. ADF&G crab/groundfish trawl survey: 2004 biomass and length composition.
4. A section on ecosystem considerations has been added.

Assessment model

The age-structured assessment model developed using ADModel Builder (a C++ software language extension and automatic differentiation library) and used for assessments in 1999-2003 is fundamentally unchanged. Model exploration focused on evaluating the contribution of each survey time series to the assessment.

Assessment results

The model estimate of spawning biomass in 2005 is 211,660 t, which is 37.0% of unfished spawning biomass and below $B_{40\%}$ (229,000 t), thereby placing Gulf of Alaska pollock in sub-tier "b" of Tier 3. Estimates of an improvement in stock status in 2004 are due to increases in biomass estimates from the 2004 Shelikof Strait EIT survey (8% increase) and the 2004 ADFG trawl survey (48% increase). Increases in spawning biomass are due to the maturation of the relatively strong 1999 and 2000 year classes. However, spawning biomass is projected to peak in 2005 and decline in following years due to lack of significant recruitment since the 2000 year class. The author's 2005 ABC recommendation for pollock in the Gulf of Alaska west of 140° W lon. (W/C/WYK) is 78,980 t. This recommendation is based on a more conservative alternative to the maximum permissible F_{ABC} introduced in the 2001 SAFE, and a temporary change to an F50% harvest rate in 2005 and 2006 to dampen yield variability and lessen short-term stock decline.

For pollock in southeast Alaska (East Yakutat and Southeastern areas), the ABC recommendation in Appendix A is 6,520 t and the OFL is 8,694 t.

Responses to Comments of the Scientific and Statistical Committee (SSC)

From the December, 2003 minutes: "*The SSC recommends that the Plan Team carefully examine the method of seasonal and area apportionments and provide a detailed explanation and rationale.*"

Appendix C of this document is a detailed explanation and rationale for the methods used to apportion the TAC among management areas on a seasonal basis.

Introduction

Walleye pollock (*Theragra chalcogramma*) is a semi-pelagic schooling fish widely distributed in the North Pacific Ocean. Pollock in the Gulf of Alaska are managed as a single stock independently of pollock in the Bering Sea and Aleutian Islands. The separation of pollock in Alaskan waters into eastern Bering Sea and Gulf of Alaska stocks is supported by analysis of larval drift patterns from spawning locations (Bailey et al. 1997), genetic studies of allozyme frequencies (Grant and Utter 1980), mtDNA variability (Mulligan et al. 1992), and microsatellite allele variability (Bailey et al. 1997).

The results of studies of stock structure in the Gulf of Alaska are equivocal. There is evidence from allozyme frequency and mtDNA that spawning populations in the northern part of the Gulf of Alaska (Prince William Sound and Middleton Island) may be genetically distinct from the Shelikof Strait spawning population (Olsen et al. 2002). However significant variation in allozyme frequency was found between Prince William Sound samples in 1997 and 1998, indicating a lack of stability in genetic structure for this spawning population. Olsen et al. (2002) suggest that interannual genetic variation may be due to variable reproductive success, adult philopatry, source-sink population structure, or utilization of the same spawning areas by genetically distinct stocks with different spawning timing. Peak spawning at the two major spawning areas in the Gulf of Alaska occurs at different times. In the Shumagin Island area, peak spawning occurs between February 15- March 1, while in Shelikof Strait peak spawning occurs between March 15 and April 1. It is unclear whether the difference in timing is genetic or caused by differing environmental conditions in the two areas.

Fishery

The commercial fishery for walleye pollock in the Gulf of Alaska started as a foreign fishery in the early 1970s (Megrey 1989). Catches increased rapidly during the late 1970s and early 1980s (Table 1.1). A large spawning aggregation was discovered in Shelikof Strait in 1981, and a fishery developed for which pollock roe was an important product. The domestic fishery for pollock developed rapidly in the Gulf of Alaska with only a short period of joint venture operations in the mid-1980s. The fishery was fully domestic by 1988.

The fishery for pollock in the Gulf of Alaska is entirely shore-based with approximately 90% of the catch taken with pelagic trawls. During winter, fishing effort targets on pre-spawning aggregations in Shelikof Strait and near the Shumagin Islands (Fig. 1.1). Fishing in summer is less predictable, but typically occurs on the east side of Kodiak Island and in nearshore waters along the Alaska Peninsula.

Incidental catch in the Gulf of Alaska pollock fishery is low. For tows classified as pollock targets in the Gulf of Alaska, more than 95% of the catch by weight consists of pollock (Fig. 1.2). Arrowtooth flounder and Pacific cod account for approximately 60% of the incidental catch, with the remainder consisting of a variety of species, including flatfish, jellyfish, smelts, sharks, and skates.

Kodiak is the major port for pollock in the Gulf of Alaska, with 61% of the 1999-2003 landings. Sand Point and Dutch Harbor are also important ports, sharing 28% of 1998-2002 landings. Secondary ports, including Cordova, Port Moller, King Cove, Akutan, and Kenai, account for the remaining 11% of the 1999-2003 landings.

Since 1992, the Gulf of Alaska pollock TAC has been apportioned spatially and temporally to reduce impacts on Steller sea lions. The details of the apportionment scheme have evolved over time, but the general objective is to allocate the TAC to management areas based on the distribution of surveyed biomass, and to establish three or four seasons between mid-January and autumn during which some

fraction of the TAC can be taken. The Steller Sea Lion Protection Measures implemented in 2001 establish four seasons in the Central and Western GOA beginning January 20, March 10, August 25, and October 1, with 25% of the total TAC allocated to each season. Allocations to management areas 610, 620 and 630 are based on the seasonal biomass distribution as estimated by groundfish surveys. In addition, a new harvest control rule was implemented that requires a cessation of fishing when spawning biomass declines below 20% of unfished stock biomass.

Data Used in the Assessment

The data used in the assessment model consist of estimates of annual catch in tons, fishery age composition, NMFS summer bottom trawl survey estimates of biomass and age composition, echo integration trawl (EIT) survey estimates of biomass and age composition in Shelikof Strait, egg production estimates of spawning biomass in Shelikof Strait, ADF&G bottom trawl survey estimates of biomass and length and age composition, and historical estimates of biomass and length and age composition from surveys conducted prior to 1984 using a 400-mesh eastern trawl. Binned length composition data are used in the model only when age composition estimates are unavailable, such as the fishery in the early part of the modeled time period.

Total Catch

Estimated catch was derived by the NMFS Regional Office from shoreside electronic logbooks and observer estimates of at-sea discard (Table 1.2). Catches include the state-managed pollock fishery in Prince William Sound. In 1996-2004, the pollock Guideline Harvest Level (GHL) for the PWS fishery was deducted from the Acceptable Biological Catch (ABC) by the NPFMC Gulf of Alaska Plan Team.

Fishery Age Composition

Estimates of fishery age composition were derived from at-sea and port sampling of the pollock catch for length and ageing structures (otoliths). Pollock otoliths collected during the 2003 fishery were aged using the revised criteria described in Hollowed et al. (1995). Catch age composition was estimated using methods described by Kimura and Chikuni (1989). Age samples were used to construct age-length keys by sex and stratum. These keys were applied to length frequency data to obtain stratum-specific age composition estimates, which were then weighted by the catch in numbers in each stratum to obtain an overall age composition. Age and length samples from the 2003 fishery were stratified by half year and statistical area as follows:

Time strata		Shumagin-610	Chirikof-620	Kodiak-630	W. Yakutat and PWS-640 and 649
1st half (A and B seasons)	No. ages	137	447	241	217
	No. lengths	926	6719	1432	1214
2nd half (C and D seasons)	No. ages	461	321	386	---
	No. lengths	2456	1726	2122	---

In the first half of 2003, a broad range of ages (ages 3-9) were relatively common in all areas, except in area 620, where the age-4 fish from the 1999 year class were clearly dominant. In the second half of 2003, mean age decreased substantially in all areas and catches were dominated by age-3 and age-4 fish (2000 and 1999 year classes, Fig. 1.3).

Fishery catch at age in 1976-2003 is presented in Table 1.3 (See also Fig. 1.4). Sample sizes for ages and lengths are given in Table 1.4.

Gulf of Alaska Bottom Trawl Survey

Trawl surveys have been conducted by Alaska Fisheries Science Center (AFSC) every three years (beginning in 1984) to assess the abundance of groundfish in the Gulf of Alaska (Table 1.5). Starting in 2001, the survey frequency was increased to every two years. The year 2004 is an “off-year” in the survey schedule, so no new data are available for the current assessment. The survey uses a stratified random design, with 49 strata based on depth, habitat, and management area (Martin 1997). Area-swept biomass estimates are obtained using mean CPUE (standardized for trawling distance and mean net width) and stratum area. The survey is conducted from chartered commercial bottom trawlers using standardized poly-Nor’eastern high opening bottom trawls rigged with roller gear. Surveying effort averages 800 tows, 70% of which contain pollock (Table 1.6).

The time series of pollock biomass used in the assessment model is based on the surveyed area in the Gulf of Alaska west of 140° W lon., obtained by adding the biomass estimates for the Shumagin, Chirikof, Kodiak INPFC areas, and the western portion of Yakutat INPFC area. Biomass estimates for 1990, 1993, 1996, 1999, and 2003 for the west Yakutat region were obtained by splitting strata and survey CPUE data at 140° W lon. (M. Martin, AFSC, Seattle, WA, pers. comm. 1998). For surveys in 1984 and 1987, the average percent in West Yakutat in the 1990-99 surveys was used (2.7% increase). The average was also used in 2001, when West Yakutat was not surveyed.

An adjustment was made to the survey time series to account for unsurveyed pollock in Prince William Sound. This adjustment was derived from an area-swept biomass estimate for PWS from a trawl survey conducted by ADF&G in 1999, using a standard ADF&G 400 mesh eastern trawl. The 1999 biomass estimate for PWS was 6,304 t ± 2,812 t (95% CI) (W. Bechtol, ADF&G, 1999, pers. comm.). The PWS biomass estimate should be considered a minimum estimate because ADF&G survey gear is less effective at catching pollock compared to the triennial survey gear (von Szalay and Brown 2001). For 1999, the biomass estimates for the NMFS bottom trawl survey and the PWS survey were simply added to obtain a total biomass estimate. The adjustment factor for the 1999 survey, (PWS + NMFS)/NMFS, was applied to other triennial surveys, and increased biomass by 1.05%. We consider this an interim approach to assessing PWS pollock, and anticipate improvements from increased surveying effort in PWS and additional comparative work.

Bottom Trawl Age Composition

Estimates of numbers at age from the bottom trawl survey were obtained from random otolith samples and length frequency samples (Table 1.7). Numbers at age were estimated for three strata: Western GOA (Shumagin INPFC area), Central GOA (Chirikof and Kodiak INPFC areas), Eastern GOA (Yakutat and Southeastern INPFC areas) using age-length keys and CPUE-weighted length frequency data. The combined Western and Central age composition was used in the assessment model.

Shelikof Strait Echo Integration Trawl Survey

Echo integration trawl surveys to assess the biomass of pollock in the Shelikof Strait area have been conducted annually since 1981 (except 1982 and 1999). Survey methods and results for 2004 are presented in an Appendix to the SAFE (Guttormsen et al. 2004). The 2004 biomass estimate for age 2+ pollock in Shelikof Strait is 291,200 t, an increase of 8% from the 2003 biomass (Table 1.5). Biomass ≥43 cm biomass (a proxy for spawning biomass) increased by approximately five times from the 2003 estimate primarily due to the maturation of the relatively strong 1999 year class. in Shelikof Strait (Fig. 1.5). The estimated abundance of age-5 fish (0.318 billion) (1999 year class) was the largest since the early 1980s in the Shelikof Strait EIT time series.

Scheduled ship maintenance in 2004 precluded more extensive survey effort in winter spawning areas, as in 2002 and 2003. Only the shelf break area near Chirikof Island was surveyed. Estimated biomass was 30,400 t, nearly identical to 2003 estimate, but considerably lower than the 2002 estimate of 82,100 t.

Since the assessment model only includes individuals age 2 and older, the biomass of age-1 fish in the 1995 and 2000 surveys was subtracted from the total biomass for those years (reducing the biomass by 15% and 14% respectively (Table 1.5). In all other years, the biomass of age-1 fish was less than 2% of the total EIT biomass estimate.

Echo Integrated Trawl Survey Length Frequency

Annual biomass distributions by length from the Shelikof Strait EIT survey show the progression of strong year classes through the population (Fig. 1.6). In the 2004 survey, the length frequency is dominated by the age-5 fish from the 1999 year class. Length frequency data are not used in the assessment model because estimates of age composition are available for all surveys.

Echo Integrated Trawl Survey Age Composition

Estimates of numbers at age from the Shelikof Strait EIT survey (1981 - 1991, 1994 -1998, 2000-2003 (Table 1.7) were obtained from random otolith samples and length frequency samples. Otoliths collected during the 1994 - 2003 EIT surveys were aged using the revised criteria described in Hollowed et al. (1995). Sample sizes for ages and lengths are given Table 1.6.

Egg Production Estimates of Spawning Biomass

Estimates of spawning biomass in Shelikof Strait based on egg production methods were included in the assessment model. A complete description of the estimation process is given in Picquelle and Megrey (1993). The estimates of spawning biomass in Shelikof Strait show a pattern similar to the acoustic survey (Table 1.5). The annual egg production spawning biomass estimate for 1981 is questionable because of sampling deficiencies during the egg surveys for that year (Kendall and Picquelle 1990). Coefficients of variation (CV) associated with these estimates were included in the assessment model. Egg production estimates were discontinued because the Shelikof Strait EIT survey provided similar information.

Alaska Department of Fish and Game Crab/Groundfish Trawl Survey

The Alaska Department of Fish and Game (ADF&G) has conducted bottom trawl surveys of nearshore areas of the Gulf of Alaska since 1987. Although these surveys are designed to monitor population trends of Tanner crab and red king crab, walleye pollock and other fish are also sampled. Standardized survey methods using a 400-mesh eastern trawl were employed from 1987 to the present. The survey is designed to sample a fixed number of stations from mostly nearshore areas from Kodiak Island to Unimak Pass, and does not cover the entire shelf area. The average number of tows completed during the survey is 360. Details of the ADF&G trawl gear and sampling procedures are in Blackburn and Pengilly (1994).

The 2004 biomass estimate for pollock for the ADF&G crab/groundfish survey was 99,400 t, an increase of 48% from the 2003 biomass estimate (Table 1.5).

ADF&G Survey Length Frequency

Pollock length-frequency for the ADF&G survey in 1989-2002 (excluding 1991 and 1995) typically show a primary mode at lengths greater than 45 cm (Fig. 1.7). The predominance of large fish in the ADF&G survey may result from the selectivity of the gear, or because of greater abundance of large pollock in the areas surveyed.

ADF&G Survey Age Composition

Ages were determined by age readers in the AFSC age and growth unit from samples of pollock otoliths collected during the 2000 and 2002 ADF&G surveys (N = 559 & 538). Comparison with fishery age composition shows that older fish (> age-8) are more common in the ADF&G crab/groundfish survey. This is consistent with the assessment model, which estimates a domed-shaped selectivity pattern for the fishery, but an asymptotic selectivity pattern for the ADF&G survey.

Pre-1984 bottom trawl surveys

Considerable survey work was carried out in the Gulf of Alaska prior to the start of the NMFS triennial bottom trawl surveys in 1984. Between 1961 and the mid-1980s, the most common bottom trawl used for surveying was the 400-mesh eastern trawl. This trawl (or minor variants thereof) was used by IPHC for juvenile halibut surveys in the 1960s, 1970s, and early 1980s, and by NMFS for groundfish surveys in the 1970s.

Comparative work using the ADF&G 400-mesh eastern trawl and the NMFS poly-Nor' eastern trawl produced estimates of relative catchability (von Szalay and Brown 2001), making it possible to evaluate trends in pollock abundance from these earlier surveys in the pollock assessment. Von Szalay and Brown (2001) estimated a fishing power correction (FPC) for the ADF&G 400-mesh eastern trawl of 3.84 (SE = 1.26), indicating that 400-mesh eastern trawl CPUE for pollock would need to be multiplied by this factor to be comparable to the NMFS poly-Nor' eastern trawl.

In most cases, earlier surveys in the Gulf of Alaska were not designed to be comprehensive, with the general strategy being to cover the Gulf of Alaska west of Cape Spencer over a period of years, or to survey a large area to obtain an index for group of groundfish, i.e., flatfish or rockfish. For example, Ronholt et al. (1978) combined surveys for several years to obtain gulfwide estimates of pollock biomass for 1973-6. There are several difficulties with such an approach, including the possibility of double-counting or missing a portion of the stock that happened to migrate between surveyed areas.

We obtained an annual gulfwide index of pollock abundance using generalized linear models (GLM). Based on examination of historical survey trawl locations, we identified four index sites (one per INPFC area) that were surveyed relatively consistently during the period 1961-1983, and during the triennial survey time series (1984-99). The index sites were designed to include a range of bottom depths from nearshore to the continental slope. We fit a generalized linear model (GLM) to pollock CPUE data with year, site, depth strata (0-100 m, 100-200 m, 200-300 m, >300 m), and a site-depth interaction as factors. Both the pre-1984 400-mesh eastern trawl data and post-1984 triennial trawl survey data were used. For the earlier period, analysis was limited to sites where at least 20 trawls were made during the summer (May 1-Sept 15).

Pollock CPUE data consist of observations with zero catch and positive values otherwise, so we used a GLM model with Poisson error and a logarithmic link (Hastie and Tibshirani 1990). This form of GLM has been used in other marine ecology applications to analyze trawl survey data (Smith 1990, Swartzman et al. 1992). The fitted model was used to predict mean CPUE by site and depth for each year with survey data. Predicted CPUEs (kg km^{-2}) were multiplied by the area within a depth strata by INPFC area (km^2) and summed to obtain proxy biomass estimates by INPFC area. Since each INPFC area contained only a single non-randomly selected index site, these proxy biomass estimates are potentially biased and would not incorporate the variability in relationship between the mean CPUE at an index site and the mean CPUE for the entire INPFC area. We used a comparison between these proxy biomass estimates by INPFC area and the actual NMFS triennial survey estimates by INPFC area for 1984-99 to obtain correction factors and variance estimates. Correction factors had the form of a ratio estimate (Cochran 1977), in which the sum of the NMFS survey biomass estimates for an INPFC area for 1984-99 is divided

by the sum of the proxy biomass estimates for the same period.

Variances were obtained by bootstrapping data within site-depth strata and repeating the biomass estimation algorithm. A parametric bootstrap assuming a lognormal distribution was used for the INPFC area correction factors. Variance estimates do not reflect the uncertainty in the FPC estimate. In the assessment model, we do not apply the FPC to the biomass estimates, but instead include the information about FPC estimate (mean and variance) as a likelihood component for relative survey catchability,

$$\log L = \frac{(q_1/q_2 - \hat{FPC})^2}{2\sigma_{FPC}^2},$$

where q_1 is the catchability of the NMFS bottom trawl survey, q_2 is the catchability of historical 400-mesh eastern trawl surveys, \hat{FPC} is the estimated fishing power correction (= 3.84), and σ_{FPC} is the standard error of the FPC estimate (= 1.26).

Estimates of pollock biomass were very low (<300,000 t) between 1961 and 1971, increased by at least a factor of ten in 1974 and 1975, and then declined to approximately 900,000 t in 1978 (Table 1.8). No trend in pollock abundance is noticeable since 1978, and biomass estimates during 1978-1982 are in the same range as the post-1984 triennial survey biomass estimates. The coefficients of variation (CV) for GLM-based biomass estimates range between 0.24 and 0.64, and, as should be anticipated, are larger than the triennial survey biomass estimates, which range between 0.12 and 0.38.

Results were generally consistent with the multi-year combined survey estimates published previously (Table 1.8), and indicate a large increase in pollock biomass in the Gulf of Alaska occurred between the early 1960s (~200,000 t) and the mid 1970s (>2,000,000 t). Increases in pollock biomass between the 1960s and 1970s were also noted by Alton et al. (1987). In the 1961 survey, pollock were a relatively minor component of the groundfish community with a mean CPUE of 16 kg/hr (Ronholt et al. 1978). Arrowtooth flounder was the most common groundfish with a mean CPUE of 91 kg/hr. In the 1973-76 surveys, the CPUE of arrowtooth flounder was similar to the 1961 survey (83 kg/hr), but pollock CPUE had increased 20-fold to 321 kg/hr, and was by far the dominant groundfish species in the Gulf of Alaska. Meuter and Norcross (2002) also found that pollock was low in the relative abundance in 1960s, became the dominant species in Gulf of Alaska groundfish community in the 1970s, and subsequently declined in relative abundance.

Questions concerning the comparability of pollock CPUE data from historical trawl surveys with later surveys probably can never be fully resolved. However, because of the large magnitude of the change in CPUE between the surveys in the 1960s and the early 1970s using similar trawling gear, the conclusion that there was a large increase pollock biomass seems robust. Model results suggest that population biomass in 1961, prior to large-scale commercial exploitation of the stock, may have been the lowest observed. Early speculation about the rise of pollock in the Gulf of Alaska in the early 1970s implicated the large biomass removals of Pacific Ocean perch, a potential competitor for euphausiid prey (Somerton et al. 1979, Alton et al. 1987). More recent work has focused on role of climate change (Anderson and Piatt 1999, Bailey 2000). The occurrence of large fluctuations in pollock abundance without large changes in direct fishing impacts suggests a need for conservative management. If pollock abundance is controlled primarily by the environment, or through indirect ecosystem effects, it may be difficult to reverse population declines, or to achieve rebuilding targets should the stock become depleted. Reliance on sustained pollock harvests in the Gulf of Alaska, whether by individual fishermen, processing companies, or fishing communities, may be difficult over the long-term.

Qualitative trends

To qualitatively assess recent trends in abundance, we standardized each survey time series by dividing the annual estimate by the average since 1986 so all could be plotted on the same scale. The Shelikof Strait EIT survey was split into separate time series corresponding to the two acoustic systems used for the survey. Although there is considerable variability in each survey time series, a fairly clear downward trend is evident (Fig. 1.8). A lowess scatterplot smoother (SPLUS 1993) fit to the relative abundance data in aggregate shows a similar, but more gradual, decline than the estimated biomass trend from the assessment model.

We also evaluated indices derived from fisheries catch data for trends in biological characteristics (Fig. 1.9). The percent of females in the catch shows no obvious trend that would indicate differential mortality on the more valuable roe-bearing females. The mean age shows interannual variability due to strong year classes passing through the population, but no downward trends that would suggest excessive mortality rates. The percent of old fish in the catch (nominally defined as age 8 and older) is also highly variable due to variability in year class strength, but has apparently increased over time, opposite to the typical pattern in an exploited population. This pattern may be partly attributable to changes in fishery selectivity or to changes in ageing criteria, but is consistent with assessment results showing low fishing mortality rates. We computed an index of catch at age diversity using the Shannon-Wiener information index,

$$- \sum p_a \ln p_a ,$$

where p_a is the proportion at age. Increases in fishing mortality would tend to reduce age diversity, but year class variability would also influence age diversity. The index of age diversity is relatively stable during 1976-2002 (Fig. 1.9).

McKelvey Index

McKelvey (1996) found a significant correlation between the abundance of age-1 pollock in the Shelikof Strait EIT survey and subsequent estimates of year-class strength. The McKelvey index is defined as the estimated abundance of 9-16 cm fish in the Shelikof Strait EIT survey, and is an index of recruitment at age 2 in the following year (Table 1.9). The relationship between the abundance of age-1 pollock in the Shelikof Strait EIT survey and year-class strength provides a recruitment forecast for the year following the most recent Shelikof Strait EIT survey. The 2004 Shelikof EIT survey age-1 estimate is 8.3 million (19th in abundance out of 21 surveys), a continuation of the low values for this index since 2002.

2004 FOCI Year Class Prediction

Data

This forecast is based on five data sources: three physical properties and two biological data sets. The sources are: 1) observed 2004 Kodiak monthly precipitation. Monthly precipitation totals (inches) are prepared by the Kodiak, Alaska, National Weather Service (NWS) Office from hourly observations. Data were obtained from the NOAA National Climate Data Center, Asheville, North Carolina, and the NWS office in Kodiak, Alaska. 2) wind mixing energy at [57N, 156W] estimated from 2004 sea-level pressure analyses; Monthly estimates of wind mixing energy ($W m^{-2}$) were computed for a location [57°N, 156°W] near the southwestern end of Shelikof Strait. To make the estimates, twice daily gradient winds are computed for that location using the METLIB utility (NOAA Tech. Memo. ERL PMEL-54). Gradient winds are converted to surface winds using an empirical formula based on the work of Macklin *et al.* (*J. Geophys. Res.*, 98, 16,555–16,569, 1993). Wind mixing energy are computed using constant air density ($1.293 kg m^{-3}$) and the drag coefficient formulation of Large and Pond (*J. Phys. Oceanogr.*, 2, 464-482, 1982). 3) advection of ocean water in the vicinity of Shelikof Strait inferred from drogued drifters

deployed during the spring of 2004, 4) rough counts of pollock larvae from a survey conducted in May 2004, and 5) estimates of age 2 pollock abundance from this years assessment.

Analysis

Kodiak Precipitation: FOCI believes that Kodiak precipitation is a valid proxy for fresh-water runoff that contributes to the density contrast between coastal and Alaska Coastal Current water in Shelikof Strait. The greater the contrast, the more likely that eddies and other instabilities will form. Such secondary circulations have attributes that make them beneficial to survival of larval pollock.

The season began with typical precipitation during January, and then it turned into one of the six wettest seasons since the beginning of the record in 1961. Three of the six months in the reporting period had significantly greater than normal precipitation. April 2004 was the second wettest April since 1961.

Kodiak precipitation for 2004

<i>Month</i>	<i>% 30-yr average</i>
Jan	94
Feb	183
Mar	76
Apr	220
May	103
June	161

Based on this information, the forecast element for Kodiak 2004 rainfall has a score of 2.48. This is "average to strong" on the continuum from 1 (weak) to 3 (strong).

Wind Mixing: Following the decadal trend established in the late 1990s, wind mixing at the southern end of Shelikof Strait was again below the long-term average for the winter and spring months of 2004.

Wind mixing at the exit of Shelikof Strait for 2004

<i>Month</i>	<i>% 30-yr average</i>
Jan	39
Feb	58
Mar	80
Apr	92
May	50
June	78

Strong mixing in winter helps transport nutrients into the upper ocean layer to provide a basis for the spring phytoplankton bloom. Weak spring mixing is thought to better enable first feeding pollock larvae to locate and capture food. Weak mixing in winter is not conducive to high survival rates, while weak mixing in spring favors recruitment. This year's scenario produces a wind mixing score of 1.99, which equates to "average".

Advection: From an examination of drifter trajectories and wind forcing, the transport in Shelikof Strait for spring of 2004 was average. We have hypothesized that very strong transport is bad for pollock survival, and that moderate transport is best and that very weak transport is, while not as disastrous as strong transport, still detrimental to larval survival. Advection was given a score of 2.0.

Relating Larval Index to Recruitment: As in last years analysis, a nonlinear neural network model with one input neuron (larval abundance), 3 hidden neurons, and one output neuron (recruitment) was used to


relate larval abundance (catch/m²) to age 2 recruitment abundance (billions). The model estimated 6 weighting parameters.

Data used in the neural network model

<i>Year Class</i>	<i>Average larval abundance (catch/m²)</i>	<i>Age-2 recruitment (billions)</i>
1982	66.44347	0.439853
1985	80.4266	0.475708
1987	324.9025	0.363344
1988	256.9029	1.6131
1989	537.2943	1.04753
1990	335.0086	0.420166
1991	54.2223	0.240026
1992	563.6741	0.141524
1993	45.80764	0.212711
1994	124.9386	0.831484
1995	600.9925	0.403814
1996	472.0225	0.172953
1997	561.1063	0.180909
1998	72.81539	0.26786
1999	102.3862	1.14857
2000	486.1835	0.769093
2001	174.624	0.109923
2002	276.6972	0.183527

The neural network model, which used the first 18 observation pairs were fit to the model and had a R² of 0.143. A plot of the observed recruitment (actual) and that predicted from larval abundance (predicted) are given below where row number corresponds to the rows of the data matrix given above.

Actual values vs Predicted values



Observed and predicted recruitment values from the larval index-recruitment neural network model.


The trained network was then used to predict the recruitment for 2003 and 2004.

The predictions are

<i>Year</i>	<i>Actual Recruitment</i>	<i>Predicted Recruitment</i>
2003	n/a	0.538
2004	n/a	0.108

These values, using the 33% (0.340938) and 66% (0.691343) cutoff points given below correspond to a weak 2003 year class and a weak 2004 year class.


Larval Index Counts: Plotting the data by year and binning the data into catch/10 m² categories (given below) provides another view of the data. The pattern for 2004 (based on rough counts) shows that all data fall into the three lowest binning categories indicating that the 2004 year class may one of the weakest in this time series.



A series of histograms for larval walleye pollock densities in late May from 1982 to 2004. Data were binned into catch/10 m² categories. The data from 2000-2004 are rough counts taken at sea, and the 2004 data are from the 5MF04 cruise that was completed on June 1.

The data for the above figure are taken from a reference area that is routinely sampled and that usually contains the majority of the larvae. This year's distribution of pollock appears to be centered in the typical reference area and the larval abundance figures in the middle of the reference area seem to be average.

Given these two pieces of information, the score for larval index is set to the high end of the weak range or 1.65.



Mean catch per 10 m² for late May cruises during 1982-2004.

Spawner/Recruit Time Series: The time series of recruitment from this year's assessment was analyzed in the context of a probabilistic transition. The data set consisted of estimates of age 2 abundance from 1961-2004, representing the 1959-2002 year classes. There were a total of 44 recruitment data points. The 33% (0.341 billion) and 66% (0.691 billion) percentile cutoff points were calculated from the full time series and used to define the three recruitment states of weak, average and strong. The lower third of the data points were called weak, the middle third average and the upper third strong. Using these definitions, nine transition probabilities were then calculated:

- Probability of a weak year class following a weak
- Probability of a weak year class following an average
- Probability of a weak year class following a strong
- Probability of an average year class following a weak
- Probability of an average year class following an average
- Probability of an average year class following a strong
- Probability of a strong year class following a weak
- Probability of a strong year class following an average
- Probability of a strong year class following a strong

The probabilities were calculated with a time lag of two years so that the 2004 year class could be predicted from the size of the 2002 year class. The 2002 year class was estimated to be 0.184 billion and

was classified as weak. The probabilities of other recruitment states following a weak year class for a lag of 2 years (n=44) are given below:

2004 Year Class		2002 Year Class		Probability	N
Weak	follows	Weak		0.095	4
Average	follows	Weak		0.070	3
Strong	follows	Weak		0.146	6

The probability of a strong year class following a weak year class had the highest probability. We classified this data element as a strong, giving it a score at the low end of strong 2.34. Each of the data elements was weighted equally.

Conclusion

Based on these five elements and the weights assigned in the table below, the FOCI forecast of the 2004 year class is average.

<i>Element</i>	<i>Weights</i>	<i>Score</i>	<i>Total</i>
Time Sequence of R	0.2	2.34	0.468
Rain	0.2	2.48	0.496
Wind Mixing	0.2	1.99	0.398
Advection	0.2	2.00	0.4
Larval Index-abundance	0.2	1.65	0.33
Total	1.0		2.092 = Average

Analytic Approach

Model description

Age-structured models for the period 1961 to 2004 (44 yrs) were used to assess Gulf of Alaska pollock. Population dynamics were modeled using standard formulations for mortality and fishery catch (e.g. Fournier and Archibald 1982, Deriso et al. 1985, Hilborn and Walters 1992). Year- and age-specific fishing mortality was modeled as a product of a year effect, representing the full-recruitment fishing mortality, and an age effect, representing the selectivity of that age group to the fishery. The age effect was modeled using a double-logistic function with time-varying parameters (Dorn and Methot 1990, Sullivan et al. 1997). The model was fit to time series of catch biomass, survey indices of abundance, and estimates of age and length composition from the fishery and surveys. Details of the population dynamics and estimation equations are presented in an appendix.

Model parameters were estimated by maximizing the log likelihood of the data, viewed as a function of the parameters. Lognormal likelihoods were used for survey biomass and total catch estimates, and multinomial likelihoods were used for age and length composition data.

<i>Likelihood component</i>	<i>Statistical model for error</i>	<i>Variance assumption</i>
Fishery total catch (1964-2004)	Log-normal	CV = 0.05
POP fishery length comp. (1964-71)	Multinomial	Sample size = 60
Fishery age comp. (1972-2003)	Multinomial	Year-specific sample size = 60-400
Shelikof EIT survey biomass (1981-2004)	Log-normal	Survey-specific CV = 0.10-0.35
Shelikof EIT survey age comp. (1981-2004)	Multinomial	Sample size = 60
NMFS bottom trawl survey biomass (1984-2003)	Log-normal	Survey-specific CV = 0.11-0.38
NMFS bottom trawl survey age comp. (1984-2003)	Multinomial	Survey-specific sample size = 38-74
Egg production biomass (1981-92)	Log-normal	Survey specific CV = 0.10-0.25
ADF&G trawl survey biomass (1989-2004)	Log-normal	CV = 0.25
ADF&G survey age comp. (2000,2002)	Multinomial	Sample size = 10
ADF&G survey length comp. (1989-2004)	Multinomial	Sample size = 10
Historical trawl survey biomass (1961-1982)	Log-normal	Survey-specific CV = 0.24-0.64
Historical trawl survey age comp. (1973)	Multinomial	Sample size = 60
Historical trawl survey length comp. (1961-1982)	Multinomial	Sample size = 10
Fishery selectivity random walk process error	Log-normal	Slope CV = 0.10 (0.001 for 1961-71)
	Normal	Inflection age SD = 0.40 (0.004 for 1961-71)
Recruit process error (1961-1968,2004)	Log-normal	CV = 1.0

Recruitment

In most years, year-class abundance at age 2 was estimated as a free parameter. Constraints were imposed on recruitment at the start of the modeled time period to improve parameter estimability. Instead of estimating the abundance of each age of the initial age composition independently, we parameterized the initial age composition with mean log recruitment plus a log deviation from an equilibrium age structure based on that mean initial recruitment. A penalty was added to the log likelihood so that the log deviations would have the same variability as recruitment during the assessment period. We also used the same penalty for log deviations in recruitment for 1961-68, and in 2004. Log deviations were estimated as free parameters in other years. These relatively weak constraints were sufficient to obtain fully converged parameter estimates.

Modeling fishery data

A four-parameter double logistic equation was used to model fisheries selectivity. Instead of grouping years with similar selectivity patterns as in previous assessments (Hollowed et al., 1994, 1995, 1998), we allowed the parameters of the double logistic function to vary according to a random walk process (Sullivan et al. 1997). This approach allows selectivity to vary from one year to the next, but restricts the amount of variation that can occur. The resulting selectivity patterns are similar to those obtained by grouping years, but transitions between selectivity patterns occur gradually rather than abruptly. Constraining the selectivity pattern for a group of years to be similar can be done simply by reducing the year-specific standard deviation of the process error term. Since limited data are available from the Pacific Ocean perch fishery years (1964-71) and in 2004, the process error standard deviation for those years was assumed to be very small, so that annual changes in selectivity are not allowed during that period.

Modeling survey data

Survey abundance was assumed to be proportional to total abundance as modified by the estimated survey selectivity pattern. Expected population numbers at age for the survey were based on the mid-date of the survey, assuming constant fishing and natural mortality throughout the year. Standard deviations in the log-normal likelihood were set equal to the sampling error CV (coefficient of variation) associated with each survey estimate of abundance (Kimura 1991).

Survey catchability coefficients can be fixed or freely estimated. In previous assessments, the NMFS bottom trawl survey catchability was fixed at one as a precautionary constraint on the total biomass estimated by the model. In the 2001 assessment (Dorn et al. 2001), a likelihood profile on trawl catchability showed that the maximum likelihood estimate of trawl catchability was approximately 0.7. This result is reasonable because pollock are known to form pelagic aggregations and occur in nearshore areas not intensively sampled by the NMFS bottom trawl survey. In this assessment we carry forward a model with estimated trawl catchability as an alternative for consideration. Catchability coefficients for other surveys were estimated as free parameters. Egg production estimates of spawning stock biomass were included in the model by setting the age-specific selectivity equal to the estimated percent mature at age (Hollowed et al. 1991).

The EK500 acoustic system has been used to estimate biomass since 1992. Earlier surveys (1981-91) were obtained with an older Biosonics acoustic system (Table 1.5). Biomass estimates similar to the Biosonics acoustic system can be obtained using the EK500 when a volume backscattering (S_v) threshold of -58.5 dB is used (Hollowed et al. 1992). Because of the newer system's lower noise level, abundance estimates since 1992 have been based on a S_v threshold of -69 dB. We split the Shelikof Strait EIT survey time series into two periods corresponding to the two acoustic systems, and estimated separate survey catchability coefficients for each period. For the 1992 and 1993 surveys, biomass estimates using both noise thresholds were used to provide information on relative catchability.

Ageing error

An ageing error transition matrix is used in the assessment model to convert population numbers at age to expected fishery and survey catch at age (Table 1.10). Dorn et al. (2003) estimated this matrix using an ageing error model fit to the observed percent agreement at ages 2 and 9. Mean percent agreement is close to 100% at age 1 and declines to 40% at age 10. Annual estimates of percent agreement are variable, but show no obvious trend, from which it was concluded that using a single transition matrix for all years in the assessment model was appropriate. The model is based on a linear increase in the standard deviation of ageing error and the assumption that ageing error is normally distributed. The model predicts percent agreement by taking into account the probability that both readers are correct, both readers are off by one year in the same direction, and both readers are off by two years in the same direction (Methot 2000). The probability that both agree and were off by more than two years was considered negligible. A cooperative project between AFSC and ADF&G is in progress to validate pollock ageing criteria using radiometric methods (D. Kimura, pers. comm.)

Length frequency data

The assessment model was fit to length frequency data from various sources by converting predicted age distributions (as modified by age-specific selectivity) to predicted length distributions using an age-length transition matrix. Because seasonal differences in pollock length at age are large, several transition matrices were used. For each matrix, unbiased length distributions at age were estimated for several years using age-length keys, then averaged across years. A transition matrix estimated by Hollowed et al. (1998) was used for length-frequency data from the early period of the fishery. A transition matrix was estimated using 1992-98 Shelikof Strait EIT survey data and used for winter survey length frequency data. The following length bins were used: 17 - 27, 28 - 35, 36 - 42, 43 - 50, 51 - 55, 56 - 70 (cm). Finally, a transition matrix was estimated using second and third trimester fishery age and length data

during the years (1989-98) and was used for the ADF&G survey length frequency data. The following length bins were used: 25 - 34, 35 - 41, 42 - 45, 46 - 50, 51 - 55, 56 - 70 (cm), so that the first three bins would capture most of the summer length distribution of the age-2, age-3 and age-4 fish, respectively. Bin definitions were different for the summer and the winter transition matrices to account for the seasonal growth of the younger fish (ages 2-4).

Parameter estimation

A large number of parameters are estimated when using this modeling approach. More than half of these parameters are year-specific deviations in fishery selectivity coefficients. Parameters were estimated using ADModel Builder, a C++ software language extension and automatic differentiation library. Parameters in nonlinear models are estimated in ADModel Builder using automatic differentiation software extended from Greiwank and Corliss (1991) and developed into C++ class libraries. The optimizer in ADModel builder is a quasi-Newton routine (Press et al. 1992). The model is determined to have converged when the maximum parameter gradient is less than a small constant (set to 1×10^{-4}). ADModel builder includes post-convergence routines to calculate standard errors (or likelihood profiles) for any quantity of interest.

A list of model parameters is shown below:

<i>Population process modeled</i>	<i>Number of parameters</i>	<i>Estimation details</i>
Initial age structure	Ages 3-10 = 8	Estimated as log deviances from the log mean; constrained by random deviation process error from an equilibrium unfished age structure
Recruitment	Years 1961-2004 = 44	Estimated as log deviances from the log mean; recruitment in 1961-68, and 2004 constrained by random deviation process error.
Natural mortality	Age- and year-invariant = 1	Not estimated in the model
Fishing mortality	Years 1961-2004 = 44	Estimated as log deviances from the log mean
Mean fishery selectivity	4	Slope parameters estimated on a log scale
Annual changes in fishery selectivity	$4 * (\text{No. years} - 1) = 172$	Estimated as deviations from mean selectivity and constrained by random walk process error
Survey catchability	No. of surveys + 1 = 7	AFSC bottom trawl survey catchability not estimated, other catchabilities estimated on a log scale. Two catchability periods were estimated for the EIT survey.
Survey selectivity	10 (EIT survey: 2, BT survey: 4, ADF&G survey: 2, Historical 400-mesh eastern trawls: 2)	Slope parameters estimated on a log scale. The egg production survey uses a fixed selectivity pattern equal to maturity at age.
Total	116 primary parameters + 172 process error parameters + 2 fixed parameters = 290	

Parameters Estimated Independently

Pollock life history characteristics, including natural mortality, growth, and maturity, were estimated independently. These parameters are used in the model to estimate spawning and population biomass, and obtain predictions of fishery and survey biomass. Pollock life history parameters include:

- Natural mortality (M)
- Proportion mature at age.
- Weight at age and year by fishery and by survey

Natural mortality

Hollowed and Megrey (1990) estimated natural mortality using a variety of methods including estimates based on: a) growth parameters (Alverson and Carney 1975, and Pauly 1980), b) GSI (Gunderson and Dygert, 1988), c) monitoring cohort abundance, and d) estimation in the assessment model. These methods produced estimates of natural mortality that ranged from 0.24 to 0.30. The maximum age observed was 22 years. For the assessment modeling, natural mortality was assumed to be 0.3 for all ages.

Hollowed et al. (2000) developed a model for Gulf of Alaska pollock that accounted for predation mortality. The model suggested that natural mortality declines from 0.8 at age 2 to 0.4 at age 5, and then remains relatively stable with increasing age. In addition, stock size was higher when predation mortality was included. A theoretical analysis of a simple age-structured model by Clark (1999) evaluated the effect of an erroneous M on both estimated abundance and target harvest rates. He found that “errors in estimated abundance and target harvest rate were always in the same direction, with the result that, in the short term, extremely high exploitation rates can be recommended (unintentionally) in cases where the natural mortality rate is overestimated and historical exploitation rates in the catch-at-age data are low.” He proposed that this error could be avoided by using a conservative (low) estimate of natural mortality. This suggests that the current approach of using a potentially low but still credible estimate of M for assessment modeling is consistent with the precautionary approach. However, it should be emphasized that the role of pollock as prey in the Gulf of Alaska ecosystem cannot be fully evaluated using a single species assessment model (Hollowed et al. 2000).

Maturity at age

In the 2002 assessment, maturity at age for Gulf of Alaska pollock was estimated using maturity stage data collected during winter EIT surveys in the Gulf of Alaska during 1983-2002. These new estimates replaced a maturity at age vector estimated by Hollowed et al. (1991) using maturity stage data collected during 1983-89. Maturity stages for female pollock describe a continuous process of ovarian development between immature and post-spawning. For the purposes of estimating a maturity vector (the proportion of an age group that has been or will be reproductively active during the year) for stock assessment, all fish greater than or equal to a particular maturity stage are assumed to be mature, while those less than that stage are assumed to be immature. We assumed that maturity stages in which ovarian development had progressed to the point where ova were distinctly visible were mature. Maturity stage data should not be considered the most reliable data to estimate maturity at age. The stages are qualitative rather than quantitative, so there is subjectivity in assigning stages, and a potential for different technicians to apply criteria differently. Because the link between pre-spawning maturity stages and eventual reproductive activity later in the season is not well established, the division between mature and immature stages is problematic. Changes in the timing of spawning could also affect maturity at age estimates. Merati (1993) compared visual maturity stages with ovary histology and a blood assay for vitellogenin and found general consistency between the different approaches. Merati (1993) noted that ovaries classified as late developing stage (i.e. immature) may contain yolked eggs, but it was unclear whether these fish would spawn later in the year. The average sample size of female pollock maturity stage data per year from winter EIT surveys in the Gulf of Alaska is 850 (Table 1.11).

Estimates of maturity at age in 2004 from winter EIT surveys were above the long-term average for most ages (Fig. 1.10). For example, the proportion of mature age-4 fish was 56% compared to 26% for the long-term average. Because there did not appear to be an objective basis for excluding data, we used the 1983-2004 average maturity at age in assessment.

Logistic regression (McCullagh and Nelder 1983) was also used to estimate the age and length at 50% mature at age for each year. Annual estimates of age at 50% maturity are highly variable and range from 3.7 years in 1984 to 6.1 years in 1991, with an average of 5.0 years. Length at 50% mature is less variable than the age at 50% mature, suggesting that at least some of the variability in the age at maturity can be attributed to changes in length at age (Fig 1.11). There is less evidence of trends in the length at 50% mature, with only the 1983 and 1984 estimates as unusually low values. In 2004, the high proportion of mature age-4 fish reduced the estimated length at 50% mature to 38 cm. The average length at 50% mature for all years is approximately 43 cm.

Weight at age

Year-specific weight-at-age estimates are used in the model to obtain expected catches in biomass. Where possible, year and survey-specific weight-at-age estimates are used to obtain expected survey biomass. For each data source, unbiased estimates of length at age were obtained using year-specific age-length keys. Bias-corrected parameters for the length-weight relationship, $W = a L^b$, were also estimated. Weights at age was estimated by multiplying length at age by the predicted weight based on the length-weight regressions.

Model selection and evaluation

Model Selection

A range of different model configurations were used to assess the sensitivity of the results to model assumptions and different data sources. As in last year's assessment, we compared models with estimated and fixed NMFS trawl survey catchability. Models with different strategies to projecting yields were compared, including a model which assumed the 1999 year class was average, as in last year's yield projections. We also evaluated the effect of the ADF&G survey and the Shelikof Strait EIT survey on estimated stock status by reducing the weights used in fitting data from these surveys.

Model 1: Estimated NMFS trawl survey catchability. In previous assessments, catchability has been fixed at one as a precautionary assumption. In the previous assessments, a likelihood profile on trawl catchability showed that this parameter could be estimated. In most assessment models in the North Pacific, survey catchability is estimated as a free parameter when possible to do so, e.g., assessments for eastern Bering Sea pollock, sablefish, and Gulf of Alaska Pacific Ocean perch. Since catchability is estimated for all other surveys in the pollock assessment, there is no *a priori* reason from a technical perspective for treating the NMFS trawl survey differently.

Model 2a: A model that conforms to last year's model assumptions: trawl catchability fixed at 1.0, and all other catchabilities freely estimated.

Model 2b: As in model 2a, except for a temporary change to F50% in 2005 and 2006.

Model 2c: As in model 2a, except that the 1999 year class was assumed to be average in abundance for yield projections. The author's ABC recommendation in the 2003 assessment was based on this assumption as a precautionary measure.

Model 3: As in model 2a, except that weights used to fit the model to ADF&G survey time series were reduced (higher assumed CVs for biomass index, and lower nominal samples for length and age composition).

Model 4: As in model 2a, except that weights used to fit the model to Shelikof Strait EIT survey time series were reduced (higher assumed CVs for biomass index, and lower nominal samples for length and age composition).

Comparison of Model 1 (estimated trawl catchability) with Model 2a (fixed trawl catchability) indicate that a despite consistent difference in stock biomass (14% decrease for Model 2a), the difference in total log likelihood is small (0.78) (Table 1.12). When a similar analysis was performed in previous assessments, the estimate of catchability was ranged from 0.70 to 0.85, rather than 0.80 in the current assessment, suggesting some tendency for the estimate to jump around. Although Model 1 would be preferred by maximum likelihood criterion, the difference in model fit probably is not significant. Until a more precise estimate of catchability is possible, we consider that the historical convention of fixing catchability to be warranted. It should be noted that this represents a “hidden” element of conservatism in the assessment, since estimates of stock biomass and yield are lower when catchability is fixed. Not surprisingly, the uncertainty in biomass estimates are higher (and more realistic) for Model 1, since the assumption of known catchability in Model 2a artificially reduces uncertainty in the assessment.

Comparison of models that down weight either the ADFG trawl survey or the Shelikof EIT survey (models 3 and 4) indicate the estimated biomass trends is are broadly consistent with the base model (Fig. 1.12). All show a similar pattern of increase and decline, suggesting that no survey has a dominant influence on the estimated trend in abundance. For the full time period, down weighting the Shelikof Strait time series results in a much low peak abundance in the mid-1980s. For the period since 1990, down weighting the Shelikof Strait EIT survey results in higher biomass, while down weighting the ADFG trawl survey results in lower biomass. This suggests some lack of consistency between spawning survey in Shelikof Strait and the ADFG trawl survey.

Model Evaluation

Residual plots for model 2a (provisionally identified as the base model) were prepared to examine the goodness of fit of the base-run model to the age composition data. The Pearson residuals for a multinomial distribution are

$$r_i = \frac{p_i - \hat{p}_i}{\sqrt{(\hat{p}_i(1 - \hat{p}_i)/m)}}$$

where p_i is the observed proportion at age, \hat{p}_i is the expected proportion at age, and m is the sample size (McCullagh and Nelder 1983). Figure 1.13 is a comparison of observed and predicted fisheries age composition, and Figures 1.14-1.16 show residuals for the fit to the fishery, the Shelikof Strait EIT survey and the NMFS trawl survey age compositions, and the ADFG trawl survey length composition. Although there are large residuals for some ages and years, no severe pattern of residuals is evident in the fishery age composition. Two moderate patterns were apparent in the fishery data. The first is a tendency for strong year classes to gain strength from adjacent weaker year classes as they become older, producing a pattern of negative residuals for the adjacent year classes. This pattern is most apparent for the strong 1984 year class beginning in 1990 at age 6. In addition, there is a tendency for strong year classes to shift a year as they become older. This pattern is most obvious for the 1988 year class, which began to change into a 1989 year class in 1995.

In the Shelikof Strait EIT survey age composition, the most extreme residuals tend to be for juvenile fish of ages two and three. Since the Shelikof Strait survey covers only a portion of winter habitat of juvenile fish, this pattern could be explained by differences in spatial distribution of different year classes. For example, the 1995 year class was uncommon in the Shelikof Strait EIT survey at age two and age three, but first appeared as large numbers in the fishery age composition data as three-year-old fish in the Shumagin area in 1998. In contrast, the 1994 year class was very abundant in the Shelikof Strait EIT survey as juveniles, but was not nearly as strong in later fishery age composition data. A similar pattern seems to be developing for the 1999 year class.

Model fits to survey biomass estimates are similar to previous assessments (Dorn et al. 2003) (Figs. 1.17-1.19). General trends in survey time series are fit reasonably well. For example, both the model and all surveys show a declining trend in the 1990s. But since each survey time series shows a different pattern of decline, the model is unable to fit all surveys simultaneously. The ADF&G survey matches the model trend better than any other survey, despite receiving less weight in model fitting. The discrepancy between the NMFS trawl survey and the Shelikof Strait EIT survey biomass estimates in the 1980s accounts for the poor model fit to both time series during in those years. More recently, the model fits extremely well the 2003 biomass estimates from the NMFS bottom trawl survey and the ADF&G trawl survey, but shows a poorer fit to the 2003 and 2004 Shelikof Strait EIT survey biomass estimates and the 2004 ADF&G trawl survey biomass estimate.

A likelihood profile for NMFS trawl survey catchability shows that the likelihood is higher for models with catchability equal to 0.80 (Fig. 1.20). The change in log likelihood is very small (less than one) between models with fixed and estimated catchability, indicating that despite the large change in biomass, there is little objective basis for choosing one model over the other.

Assessment Model Results

Parameter estimates and model output for Model 2a are presented in a series of tables and figures. Estimated selectivity for different periods in the fishery and for surveys is given in Table 1.13 (see also Figure 1.21). Table 1.14 gives the estimated population numbers at age for the years 1961-2004. Table 1.15 gives the estimated time series of age 3+ population biomass, age-2 recruitment, and harvest rate (catch/3+ biomass) for 1969-2004 (see also Fig. 1.22). Stock size peaked in the early 1980s at approximately twice unfished stock size. In 1998, the stock dropped below the $B_{40\%}$ for the first time since the 1970s, and in 2004 is estimated to be at 29% of unfished stock size.

Retrospective comparison of assessment results

A retrospective comparison of assessment results for the years 1993-2003 indicates the current estimated trend in spawning biomass for 1969-2003 is consistent with previous estimates (Fig. 1.23). All time series show a similar pattern of decreasing spawning biomass in the 1990s. Retrospective biases in the assessment are small, but based on the current assessment, there was some tendency to underestimate ending year abundance from 1993 to 1997, followed by several years of overestimating ending year abundance. Assessment results from 2002, 2003 and 2004 are very consistent. The estimated 2004 age composition from the current assessment is very similar to the estimated age composition in the 2003 assessment (Fig. 1.23). The estimated number of age-2 recruits from the 2002 year class is much lower than the assumption of average recruitment in last year's assessment.

Stock and recruitment

Recruitment of Gulf of Alaska pollock is more variable ($CV = 1.06$) than Eastern Bering Sea pollock (0.61). Among North Pacific groundfish stocks with age-structured assessments, GOA pollock ranks

third in recruitment variability after sablefish and Pacific Ocean perch (<http://www.afsc.noaa.gov/refm/stocks/estimates.htm>). However, unlike sablefish and Pacific Ocean perch, pollock have a short generation time (5 yrs), so that large year classes do not persist in the population long enough to have a buffering effect on population variability. Because of these intrinsic population characteristics, the typical pattern of biomass variability for Gulf of Alaska pollock will be sharp increases due to strong recruitment, followed by periods of gradual decline until the next strong year class recruits to the population. Gulf of Alaska pollock is more likely to show this pattern than any other groundfish stock in the North Pacific due to the combination of a short generation time and high recruitment variability.

Since 1980, strong year classes have occurred every four to six years (Fig. 1.22). Because of high recruitment variability, the mean relationship between stock size and recruitment abundance is not apparent despite good contrast in stock abundance. Strong and weak year classes have been produced both at high spawning biomass and low spawning biomass. The 1972 year class (one of the largest on record) was produced by an estimated spawning biomass close to current levels, suggesting that the stock has the potential to produce strong year classes. Spawner productivity is higher at low spawning biomass compared to high spawning biomass, indicating that survival of eggs to recruitment is density-dependent (Fig. 1.24). However, this pattern of density-dependent survival emerges from strong decadal trends in spawner productivity. These decadal trends in spawner productivity have produced the pattern of increase and decline in the GOA pollock population. The last two decades have been a period of relatively low spawner productivity.

We summarize information on recent year classes in the table below. Subsequent to the 2000 year class, which appears to be moderate in abundance, information is sketchy. The 2001, 2002, and the 2003 year classes have not been common in the Shelikof Strait EIT surveys or fishery sampling, and apparently are weak in comparison to the 1999 and 2000 year classes. If the pattern of relatively strong pollock recruitment every 4-6 years continues, then the next episode of strong recruitment would be expected occur in 2005-07.

Year of recruitment	2004	2005	2006
Year class	2002	2003	2004
FOCI prediction	<i>Average</i>	<i>Average</i>	<i>Average</i>
Survey information	2003 NMFS bottom trawl age-1 estimate is 75.5 million (6th in abundance out of 9 surveys)	2004 Shelikof EIT survey age-1 estimate is 8.3 million (19th in abundance out of 21 surveys)	Anecdotal reports of exceptional abundance of age-0 pollock from 2004 east side Kodiak EIT depletion survey

Projections and Harvest Alternatives

Reference fishing mortality rates and spawning biomass levels

Since 1997, Gulf pollock have been managed under Tier 3 of NPFMC harvest guidelines. In Tier 3, reference mortality rates are based on the spawning biomass per recruit (SPR), while biomass reference levels are estimated by multiplying the SPR by average recruitment. Estimates of the FSPR harvest rates were obtained using the life history characteristics of Gulf of Alaska pollock (Table 1.16). Spawning biomass reference levels were based on mean 1979-2003 recruitment (786 million), which is 4% lower than the post-1979 mean in the 2003 assessment due to the inclusion of the weak 2001 year class in the

average. The average did not include the recruitment in 2004 (2002 year class) due to uncertainty in the estimates of year class strength. Spawning was assumed to occur on March 15th, and female spawning biomass was calculated using mean weight at age for the Shelikof Strait EIT surveys in 2000-2004 to estimate current reproductive potential. The SPR at $F=0$ was estimated as 0.729 kg/recruit, which is 4% lower than the estimate in last year's assessment primarily due to lower mean weight at age at spawning. This F_{SPR} rates depend the selectivity pattern of the fishery. Selectivity in the Gulf of Alaska pollock fishery changed as the fishery evolved from a foreign fishery occurring along the shelf break to a domestic fishery on spawning aggregations and in nearshore waters (Fig. 1.1). Since 1992, Gulf of Alaska pollock have been managed with time and area restrictions, and selectivity has been fairly stable (Fig. 1.21). For SPR calculations, we used a selectivity pattern based on an average for 1992-2003.

Gulf of Alaska pollock F_{SPR} harvest rates are given below:

F_{SPR} rate	Fishing mortality	Equilibrium under average 1979-2002 recruitment				
		Avg. Recr. (Million)	Total 3+ biom. (1000 t)	Female spawning biom. (1000 t)	Catch (1000 t)	Harvest rate
100.0%	0.000	786	1867	573	0	0.0%
50.0%	0.212	786	1261	286	139	11.1%
45.0%	0.251	786	1196	258	153	12.8%
40.0%	0.296	786	1130	229	166	14.7%
35.0%	0.351	786	1062	200	178	16.8%

The $B_{40\%}$ estimate of 229,000 t is about 8% lower than the $B_{40\%}$ estimate of 248,000 t in the 2003 assessment due to both lower post-1977 mean recruitment and lower SPR at $F=0$. The model estimate of spawning biomass in 2005 is 211,660 t, which is 37.0% of unfished spawning biomass and below $B_{40\%}$ (229,000 t), thereby placing Gulf of Alaska pollock in sub-tier "b" of Tier 3. In sub-tier "b" the OFL and maximum permissible ABC fishing mortality rates are adjusted downwards as described by the harvest guidelines (see SAFE Summary Chapter).

2004 acceptable biological catch

The definitions of OFL and maximum permissible F_{ABC} under Amendment 56 provide a buffer between the overfishing level and the intended harvest rate, as required by NMFS national standard guidelines. Since estimates of stock biomass from assessment models are uncertain, the buffer between OFL and ABC provides a margin of safety so that assessment error will not result in the OFL being inadvertently exceeded. For Gulf of Alaska pollock, the maximum permissible F_{ABC} harvest rate is 84.3% of the OFL harvest rate. In the 2001 assessment, based on an analysis that showed that the buffer between the maximum permissible F_{ABC} and OFL decreased when the stock is below approximately $B_{50\%}$, we developed a more conservative alternative that maintains a constant buffer between ABC and F_{ABC} at all stock levels (Table 1.17). While there is always some probability of exceeding F_{OFL} due to imprecise stock assessments, it did not seem reasonable to reduce safety margin as the stock declines.

This alternative is given by the following

$$\text{Define } B^* = B_{40\%} \frac{F_{35\%}}{F_{40\%}}$$

Stock status: $B / B^* > 1$, then $F = F_{40\%}$

Stock status: $0.05 < B / B^* \leq 1$, then $F = F_{40\%} \times (B / B^* - 0.05) / (1 - 0.05)$

Stock status: $B / B^* \leq 0.05$, then $F = 0$

This alternative has the same functional form as the maximum permissible F_{ABC} ; the only difference is that it declines linearly from B^* (= $B_{47\%}$) to $0.05B^*$ (Fig. 1.25).

Projections for 2005 for F_{OFL} , the maximum permissible F_{ABC} , and an adjusted $F_{40\%}$ harvest rate with a constant buffer between F_{ABC} and F_{OFL} are given for Models 1-4 in Table 1.12.

ABC recommendation

There are two major sources of new information about abundance trends in 2004. The 2004 Shelikof Strait EIT survey indicated a 8% increase in total biomass since 2003, but a strong increase (5-fold) increase in adult biomass (≥ 43 cm) due to maturation of the 1999 and 2000 year classes. The 2004 ADF&G crab/groundfish survey biomass increased by 48% from the 2003 estimate, but is only 3% higher than the 2002 estimate, suggesting that these differences are mostly sampling variability. The estimate of the 1999 year class is 14% lower than last year's estimate, but is counterbalanced by a 9% increase in 2000 year class. The overall picture both from surveys and assessment results suggests continued increase in pollock abundance since last year and a reasonable consistency with model projections in previous assessments.

The primary concerns about Gulf of Alaska pollock for the short-term are 1) the lack of recruitment to the population after the 2000 year class, 2) lower than expected spawning biomass estimates for Shelikof Strait. Since the early 1980s, there has been a pattern of relatively strong pollock recruitment every 4-6 years. If this pattern continues, the next episode of strong recruitment would be expected occur in 2005-07. As yet, there is no strong evidence that this has occurred, notwithstanding the anecdotal reports of large numbers of young-of-the-year pollock in surveys this year. The decline in spawning activity in Shelikof Strait is somewhat mitigated by the additional winter surveying efforts in 2002 and 2003, which found significant aggregations of spawning pollock elsewhere in the Gulf of Alaska. Nevertheless, the cause of these apparent changes in utilization of spawning habitat is unknown, and there is concern that changes in spawning behavior alone could impact pollock abundance in the future.

We consider Models 2a or 2b as the strongest candidates on which to base yield recommendations. The change in the estimates of NMFS trawl catchability with an additional year of data suggests that basing an assessment on an estimated trawl catchability could increase interannual variability in ABC recommendations. The model comparison suggests that the assumption that NMFS trawl catchability equals one is a reasonable precautionary assumption. Models which down weight an entire survey time series are useful for sensitivity analyses, but we are reluctant to de-emphasize a survey unless there is good evidence to think it is biased. No survey covers the entire spatial distribution of pollock (or distance above bottom). If the different components of the population sampled by each survey show different trends than the population as a whole, it may be advisable to use each survey time series as is, despite some lack of model fit, to obtain the most robust estimates of overall population trends.

The final issue is which model to use for yield projections. In previous assessments, the 1999 year class was set to average recruitment for yield projections due to uncertainty in the estimate of its magnitude. Although the estimate of the 1999 year class is 14% lower than last year's estimate, the percent change in the estimate is smaller than in previous assessments, suggesting it is reasonable to expect some consistency in subsequent assessments (Fig. 1.26). For the author's recommended ABC, we used the model estimate of the 1999 year class, but have included a projection with an assumed average 1999 year class (Model 2c) for consideration by the Plan Team or the SSC. A more important issue may be the appropriate response to the apparent lack of strong recruitment since the 2000 year class, and the projected increase in the ABC in 2005 followed by a relatively sharp decrease. Evaluation of several alternatives suggested that a temporary change to an adjusted $F_{50\%}$ harvest rate in 2005 and 2006 would stabilize yields over the short-term, reduce the rate of biomass decline, and at the same time address any residual concerns about the strength on the 1999 year class. Based on these considerations, we used Model 2b with an adjusted $F_{50\%}$ harvest rate in 2005 and 2006 for the author's recommended ABC of 78,980 t. The elements of risk-aversion in this recommendation relative to using the point estimate of the model and the maximum permissible F_{ABC} are the following: 1) fixing trawl catchability at 1.0; 2) applying a more conservative harvest rate than the maximum permissible F_{ABC} , and 3) a temporary change to an adjusted $F_{50\%}$ harvest rate in 2005 and 2006. Collectively these risk-averse elements reduce the recommended ABC to approximately 55% of the model point estimate.

To evaluate the probability that the stock is below the $B_{20\%}$ threshold, we modified the assessment model to include 2005, and assumed 2005 catches will be equal to the ABC recommendation. For 2006 and subsequent years, catch is derived from the estimate of spawning biomass in that year and the author's recommended fishing mortality schedule. We then sampled from the joint marginal likelihood of spawning biomass and fishing mortality in 2005 using Markov chain Monte Carlo (MCMC) (Fig. 1.27). A chain of 1,000,000 samples was thinned by selecting every 200th sample. Analysis of the thinned MCMC chain indicates that probability of the stock being below $B_{20\%}$ will be less than 1% in all years.

Projections and Status Determination

A standard set of projections is required for stocks managed under Tier 3 of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Protection Act, and the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA). For each scenario, the projections begin with the 2005 numbers at age as projected by the assessment model. In each year, the fishing mortality rate is determined by the spawning biomass in that year and the respective harvest scenario. Recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments during 1979-2003 as estimated by the assessment model. Spawning biomass is computed in each year based on the time of peak spawning (March 15) using the maturity and weight schedules in Table 1.16. This projection scheme is run 1000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios are used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2004, are as follows (" $max F_{ABC}$ " refers to the maximum permissible value of F_{ABC} under Amendment 56):

Scenario 1: In all future years, F is set equal to $max F_{ABC}$. (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs.)

Scenario 2: In all future years, F is set equal to the F_{ABC} recommended in the assessment.

Scenario 3: In all future years, F is set equal to 50% of $\max F_{ABC}$. (Rationale: This scenario provides a likely lower bound on F_{ABC} that still allows future harvest rates to be adjusted downward when stocks fall below reference levels.)

Scenario 4: In all future years, F is set equal to the 2000-2004 average F . (Rationale: For some stocks, TAC can be well below ABC, and recent average F may provide a better indicator of F_{TAC} than F_{ABC} .)

Scenario 5: In all future years, F is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

Two other scenarios are needed to satisfy the MSFCMA's requirement to determine whether a stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follow (for Tier 3 stocks, the MSY level is defined as $B_{35\%}$):

Scenario 6: In all future years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is overfished.)

Scenario 7: In 2005 and 2006, F is set equal to $\max F_{ABC}$, and in all subsequent years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is approaching an overfished condition.)

Results from scenarios 1-5 are presented in Table 1.18. Under all harvest policies except the $F=0$ policy, spawning biomass is projected to decrease from 2005 to 2008 due to the lack of recent recruitment, then increase gradually (Fig. 1.28).

Scenarios 6 and 7 are used to make the MSFCMA's required status determination as follows:

Spawning biomass is projected to be 210,059 t in 2005 for an FOFL harvest rate, which is less than $B_{35\%}$ (200,000 t), but greater than $\frac{1}{2}$ of $B_{35\%}$. Under scenario 6, the projected mean spawning biomass in 2015 is 217,856 t, 109% of $B_{35\%}$. Therefore, Gulf of Alaska pollock are not currently overfished.

Under scenario 7, projected mean spawning biomass in 2007 is 157,595 t, which is less than $B_{35\%}$, but greater than $\frac{1}{2}$ of $B_{35\%}$. Projected mean spawning biomass in 2017 is 219,549 t, 110% of $B_{35\%}$. Therefore, Gulf of Alaska pollock is not approaching an overfished condition.

Ecosystem considerations

Physical forcing

Pollock spawn in places and times that are conducive to egg and larval survival, and transport to suitable nursery areas (Bailey 1997). Pollock eggs and larvae are pelagic and drift in ocean currents. For pollock spawning in Shelikof Strait, formation of eddies in Alaska Coastal Current may keep larvae from being advected off the continental shelf, and may help concentrate prey. FOCI has developed pollock survival index based on Kodiak rainfall as a proxy for freshwater discharge that promotes formation of eddies. The index has high interannual variability, but generally increased from the 1960s to the mid-1980s, and has remained stable to 2004 (Fig. 23, Ecosystem Considerations Appendix). This index apparently is not associated with large-scale climate indices such as the Pacific decadal oscillation or the Arctic oscillation. A second pollock survival index developed by FOCI is based on wind mixing near the southern end of Shelikof Strait. Stronger than average mixing prior to spawning helps transport nutrients into the upper ocean layer, while weak mixing after spawning is thought to better enable first feeding pollock larvae to

locate and capture food. The wind-mixing index shows wide interannual variability and slightly increasing trend from the 1960s to present (Fig. 24, Ecosystem Considerations Appendix).

Hollowed et al. (2001) found similarities in the recruitment patterns of the dominant gadids in the North Pacific. In general, gadids exhibited low autocorrelation in recruitment, with most of the variability occurring at the interannual level. Pollock and Pacific cod in the Gulf of Alaska and Pacific hake off the US West Coast showed a higher incidence of strong year classes in those El Niño years when anomalous conditions propagated to northern latitudes (“Niño North” conditions).

Prey of pollock

An ECOPATH model was assembled to characterize food web structure in Gulf of Alaska using diet data and population estimates during 1990-94. We use ECOPATH here simply as a tool to integrate diet data and stock abundance estimates in a consistent way to evaluate ecosystem interactions. We focus primarily on first-order trophic interactions: prey of pollock and the predators of pollock.

Pollock consumption is primarily in the pelagic pathway in the food web, leading from phytoplankton through various categories of zooplankton to planktivorous fish species such as capelin and sandlance (Fig. 1.29). A potential exception are shrimp, which are more associated with the benthic pathway, and make up approximately 18% of pollock diet. Although pollock consume zooplankton during all life history stages, there is an ontogenetic shift in diet from copepods to large zooplankton (primarily euphausiids) (Fig. 1.30). Cannibalism is not as prevalent in the Gulf of Alaska as in the Eastern Bering Sea, and fish consumption is low even for large pollock (Yang and Nelson 2000). There are no extended time series of zooplankton abundance for the shelf waters of the Gulf of the Alaska. Brodeur and Ware (1995) provide evidence that biomass of zooplankton in the center of the Alaska Gyre was twice as high in the 1980s than in the 1950s and 1960s, consistent with a shift to positive values of the PDO since 1977.

Predators of pollock

Based on ECOPATH results, the top five predators on pollock >20 cm by relative importance are arrowtooth flounder, Pacific halibut, Pacific cod, Steller sea lion, and the directed pollock fishery (Fig 1.31). All major predators show some diet specialization, and none depend on pollock for more than 50% of their total consumption (Fig 1.32). Pacific halibut is most dependent on pollock (48%), followed by SSL (39%), then arrowtooth flounder (24% for juvenile and adult pollock combined), and lastly Pacific cod (18%). The diet of Pacific cod and Pacific halibut are similar in that the majority of their diet besides pollock is from the benthic pathway of the food web. Alternate prey for Steller sea lions and arrowtooth flounder are similar, and come primarily from the pelagic pathway.

Steller sea lions and the fishing fleet are central place foragers, making foraging trips from a specific locations (ports in the case of the fishing fleet, and rookeries or haulouts for Steller sea lions). Foraging bouts (or trawl sets) begin at the surface, and foragers attack their prey from the top down. In contrast, predation by groundfish is not as constrained geographically, and captures are likely occur when the predator swims upwards from the bottom. Changes in the vertical distribution of pollock may tend to favor one mode of foraging over another. For example, if pollock move deeper in the water column due to surface warming, foraging groundfish might obtain an advantage over surface foragers. Alternatively, pollock may respond adaptively to predation risks from groundfish or surface foragers by changing its position in the water column.

Aside from long-recognized decline in Steller sea lion abundance, the major predators of pollock in the Gulf of Alaska are stable to increasing, in some cases notably so (Fig. 1.33). During the short time in which reliable estimates are available (1990-2001), per unit pollock consumption has been increasing for arrowtooth flounder, stable for Pacific halibut, and decreasing for Pacific cod (Fig. 1.33). There has been

a marked decline in Pacific halibut weight at age since the 1970s that Clark et al. (1999) attributed to the 1977 regime shift without being able to determine the specific biological mechanisms that produced the change. Possibilities suggested by Clark et al. (1999) include the physiological effect of an increase in temperature, intra- and interspecific competition for prey, or a change in prey quality. The two species most dependent on pollock in the early 1990s (Pacific halibut and Steller sea lion) have both shown an exceptional biological response during the post-1977 period consistent with a reduction in carrying capacity (growth for Pacific halibut, survival for Steller sea lions). In contrast, the dominant predator on pollock in the Gulf of Alaska (arrowtooth flounder) has increased steadily in abundance over the same period and shows no evidence of decline in size at age (Fig. 1.34).

Pair-wise correlations in predator trends were examined for consistent patterns (Fig. 1.35). For each pair-wise comparison, we used the maximum number of years available. Time series for Steller sea lions and Pacific cod begin in mid 1970s, while other time series extend back to the early 1960s. We make no attempt to evaluate statistical significance (biomass trends are highly autocorrelated), and emphasize that correlation does not imply causation. If two populations are strongly correlated in time, there are many possible explanations: both populations are responding to similar forcing, one or other is causative agent, etc.

Pollock abundance, fishery catches, and Steller sea lions are positively correlated (Fig. 1.36). Since the harvest policy for pollock is modified fixed harvest rate strategy, a positive correlation between catch and abundance would be expected. The Steller sea lion trend is more strongly correlated with pollock abundance than pollock catches, but this correlation is based on data since 1976, and does not include earlier years of low pollock abundance. The only strong inverse correlation is between arrowtooth flounder and Steller sea lions. A strong positive correlation exists between Pacific cod and Pacific halibut, and, from the 1960s to the present, between Pacific halibut and arrowtooth flounder.

Several patterns are apparent in abundance trends and the diet data. First, the two predators with alternate prey in the benthic pathway, Pacific cod and Pacific halibut, covary and have been relatively stable in the post-1977 period. Second, the long term increases in both Pacific halibut and arrowtooth flounder (with quite different diets apart from pollock) may be linked to similarities in their reproductive behavior. Both spawn offshore in late winter, and conditions that enhance onshore advection, such as El Niños, may play an important role in recruitment to nursery areas for these species (Bailey and Picquelle 2002).

Finally, it is apparent that the potential for competition between Steller sea lions and arrowtooth flounder is underappreciated, perhaps because arrowtooth flounder seem poorly designed to compete as forager in the pelagic zone. However, arrowtooth flounder consume both the primary prey of Steller sea lions (pollock), and alternate pelagic prey also utilized by Steller sea lions (capelin, herring, sandlance, salmon). Arrowtooth predation on pollock occurs at a smaller size than pollock targeted by Steller sea lions. The arrowtooth flounder population is nearly unexploited, is increasing in abundance, may be increasing its per unit consumption of pollock, and shows no evidence of density-dependent growth. And lastly, since 1976 there has been a strong inverse correlation between arrowtooth flounder and Steller sea lion abundance that is at least consistent with competition between these species.

Literature Cited

- Alton, M., S. Hughes, and G. Hirschhorn. 1977. Gulf of Alaska pollock—its fisheries and resource potential. Unpubl. Manusc., 25 p. Alaska Fisheries Science Center. National Marine Fisheries Service National Oceanic and Atmospheric Administration. 7600 Sand Point Way, NE Seattle, Washington 98115-6349. Submitted to the International North Pacific Fisheries Commission in 1977 as INPFC doc. 2019.
- Alton, M. S., M. O. Nelson, and B. A. Megrey. 1987. Changes in the abundance and distribution of walleye pollock (*Theragra chalcogramma*) in the western Gulf of Alaska. *Fish. Res.* 5: 185-197.
- Alverson, D. L. And M. J. Carney. 1975. A graphic review of the growth and decay of population cohorts. *Cons. int. Explor. Mer.* 133-143.
- Anderson, P. J. and J. F. Piatt 1999. Community reorganization in the Gulf of Alaska following ocean climate regime shift. *Mar. Ecol. Prog. Ser.* 189:117-123.
- Bailey, K.M., P.J. Stabeno, and D.A. Powers. 1997. The role of larval retention and transport features in mortality and potential gene flow of walleye pollock. *J. Fish. Biol.* 51(Suppl. A):135-154.
- Bailey, K.M., T.J. Quinn II, P. Bentzen, and W.S. Grant. 1999. Population structure and dynamics of walleye pollock, *Theragra chalcogramma*. *Advances in Mar. Biol.* 37: 179-255.
- Bailey, K.M. 2000. Shifting control of recruitment of walleye pollock *Theragra chalcogramma* after a major climatic and ecosystem change. *Mar. Ecol. Prog. Ser.* 198:215-224.
- Bailey, K. M and S. J. Picquelle. 2002. Larval distribution of offshore spawning flatfish in the Gulf of Alaska: potential transport pathways and enhanced onshore transport during ENSO events. *Mar. Ecol. Prog. Ser.* 236:205-217.
- Baranov, F.I. 1918. On the question of the biological basis of fisheries. *Nauchn. Issed. Ikhtiologicheskii Inst. Izv.* 1:81-128.
- Blackburn, J. and D. Pengilly. 1994. A summary of estimated population trends of seven most abundant groundfish species in trawl surveys conducted by Alaska Department of Fish and Game in the Kodiak and Alaska Peninsula areas, 1988 through 1993. Alaska Department of Fish and Game, Regional Information Report No. 4K94-31. 19p.
- Brodeur, R. D. and Ware, D.M. 1995. Interdecadal variability in distribution and catch rates of epipelagic nekton in the Northeast Pacific Ocean. pp. 329-356 in R. J. Beamish [Ed.] *Climate change and northern fish populations*. Canadian Special Publication of Fisheries and Aquatic Sciences 121. National Research Council of Canada, Ottawa.
- Clark, W.G. 1999. Effects of an erroneous natural mortality rate on a simple age-structured model. *Can. J. Fish. Aquat. Sci.* 56:1721-1731.
- Clark, W. G., S. R. Hare, A. M. Parma, P. J. Sullivan, and R. J. Trumble. 1999. Decadal changes in growth and recruitment of Pacific halibut (*Hippoglossus stenolepis*). *Can. J. Fish. Aquat. Sci.* 56(2): 242-252.
- Cochran, W. G. 1977. *Sampling Techniques*. John Wiley and Sons. New York. 428 p.
- Deriso, R.B., T.J. Quinn II, and P.R. Neal. 1985. Catch-age analysis with auxiliary information. *Can. J. Fish. Aquat. Sci.* 42: 815-824.
- Dorn, M. W., and R. D. Methot. 1990. Status of the coastal Pacific whiting resource in 1989 and recommendation to management in 1990. U.S. Dep. Commer., NOAA Tech. Memo. NMFS F/NWC-182, 84 p.
- Dorn, M.W., A.B. Hollowed., E. Brown, B.A. Megrey, C. Wilson and J. Blackburn. 1999. Walleye pollock. In *Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska*. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Dorn, M.W., Hollowed, A.B., E. Brown, B. Megrey, C. Wilson, and J. Blackburn. 2001. Assessment of the walleye pollock stock in the Gulf of Alaska. In *Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska*. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Dorn, M.W., S. Barbeaux, B. M. Guttormsen, B. Megrey, A. Hollowed, M. Wilkins, and K. Spalinger. 2003. Assessment of the walleye pollock stock in the Gulf of Alaska. *In* Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Doubleday, W.G. 1976. A least-squares approach to analyzing catch at age data. *Res. Bull. Int. Comm. Northw. Atl. Fish.* 12:69-81.

Fournier, D. and C. P. Archibald. 1982. A general theory for analyzing catch at age data. *Can. J. Fish. Aquat. Sci.* 39:1195-1207.

Fritz, L. W. 1993. Trawl locations of walleye pollock and Atka mackerel fisheries in the Bering Sea, Aleutian Islands, and Gulf of Alaska from 1977-92. AFSC Processed Report 93-08. NMFS, AFSC, 7600 Sand Point Way, NE, Seattle, WA 98115. 162 p.

Grant, W.S. and F.M. Utter. 1980. Biochemical variation in walleye pollock *Theragra chalcogramma*: population structure in the southeastern Bering Sea and Gulf of Alaska. *Can. J. Fish. Aquat. Sci.* 37:1093-1100.

Greiwanck, A., and G.F. Corliss (eds.) 1991. Automatic differentiation of algorithms: theory, implementation and application. Proceedings of the SIAM Workshop on the Automatic Differentiation of Algorithms, held Jan. 6-8, Breckenridge, CO. Soc. Indust. and Applied Mathematics, Philadelphia.

Gunderson, D. R. and P. H. Dygert. 1988. Reproductive effort as a predictor of natural mortality rate. *J. Cons. int. Mer*, 44:200-209.

Guttormsen, M. A., C. D. Wilson, and S. Stienessen. 2001. Echo integration-trawl survey results for walleye pollock in the Gulf of Alaska during 2001. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M. A., C. D. Wilson, and S. Stienessen. 2002. Echo integration-trawl survey results for walleye pollock in the Gulf of Alaska during February and March 2002. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M. A., C. D. Wilson, and S. Stienessen. 2003. Results of the February and March 2003 Echo integration-trawl surveys of walleye pollock (*Theragra chalcogramma*) conducted in the Gulf of Alaska, Cruises MF2003-01 and MF2003-05. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Guttormsen, M. A. 2004. Results of the March-April 2004 echo integration-trawl surveys of walleye pollock (*Theragra chalcogramma*) conducted in the Gulf of Alaska, Cruise MF0403. *In* Stock Assessment and Fishery Evaluation Report for Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Hastie, T., and R. Tibshirani. 1990. Generalized additive models. Chapman and Hall, London. 289 pp.

Hilborn, R. and C.J. Walters. 1992. Quantitative fisheries stock assessment: choice, dynamics, and uncertainty. Chapman and Hall, New York, N.Y. 570 p.

Hollowed, A.B. and B.A. Megrey. 1990. Walleye pollock. *In* Stock Assessment and Fishery Evaluation Report for the 1991 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.

Hollowed, A.B., B.A. Megrey, P. Munro, and W. Karp. 1991. Walleye pollock. *In* Stock Assessment and Fishery Evaluation Report for the 1992 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.

Hollowed, A. B., B.A. Megrey, and W. Karp. 1992. Walleye pollock. *In* Stock Assessment and Fishery Evaluation Report for the 1993 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.

- Hollowed, A.B., C. Wilson, E. Brown, and B.A. Megrey. 1994. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for the 1995 Gulf of Alaska Groundfish Fishery. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510.
- Hollowed, A.B., E. Brown, P. Livingston, B.A. Megrey and C. Wilson. 1995. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for Gulf of Alaska As Projected for 1996. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Hollowed, A.B., E. Brown, J. Ianelli, B.A. Megrey and C. Wilson. 1998. Walleye pollock. In Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Hollowed, A.B., J.N. Ianelli, P. Livingston. 2000. Including predation mortality in stock assessments: a case study for Gulf of Alaska pollock. *ICES J. Mar. Sci.* 57:279-293.
- Hollowed, A.B., S. R. Hare and W. S. Wooster. 2001. Pacific Basin climate variability and patterns of Northeast Pacific marine fish production. *Progress In Oceanography* 49: 257-282.
- Hughes, S. E. and G. Hirschhorn. 1979. Biology of walleye pollock, *Theragra chalcogramma*, in Western Gulf of Alaska. *Fish. Bull.*, U.S. 77:263-274.
- Karp, W. A. 1990. Results of echo integration midwater-trawl surveys for walleye pollock in the Gulf of Alaska in 1990. Appendix 3 of Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Kendall, A.W. Jr. and S.J. Picquelle. 1990. Egg and larval distributions of walleye pollock *Theragra chalcogramma* in Shelikof Strait, Gulf of Alaska. *Fish. Bull.*, U.S. 88:133-154.
- Kimura, D.K. 1989. Variability, tuning, and simulation for the Doubleday-Deriso catch-at-age model. *Can. J. Fish. Aquat. Sci.* 46:941-949.
- Kimura, D.K. 1990. Approaches to age-structured separable sequential population analysis. *Can. J. Fish. Aquat. Sci.* 47:2364-2374.
- Kimura, D.K. 1991. Improved methods for separable sequential population analysis. Unpublished. Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle, Washington 98115.
- Kimura, D. K. and S. Chikuni. 1989. Variability in estimating catch-in-numbers-at-age and its impact on cohort analysis. *In* R.J. Beamish and G.A. McFarlane (eds.), Effects of ocean variability on recruitment and an evaluation of parameters used in stock assessment models. *Can. Spec. Publ. Fish. Aquat. Sci.* 108:57-66.
- McCullagh, P., and J. A. Nelder. 1983. Generalized linear models. Chapman and Hall, London. 261 p.
- McKelvey, D. 1996. Juvenile walleye pollock, *Theragra chalcogramma*, distribution and abundance in Shelikof Strait—What can we learn from acoustic survey results? p. 25-34. *In* U.S. Dep. Commer. NOAA Tech. Rep. NMFS 126.
- McKelvey, D.R. 1996. Juvenile walleye pollock, *Theragra chalcogramma*, distribution and abundance in Shelikof Strait—what can we learn from acoustic surveys. Ecology of Juvenile Walleye Pollock, *Theragra chalcogramma*. NOAA Technical Report NMFS 126, p 25-34.
- Megrey, B.A. 1989. Exploitation of walleye pollock resources in the Gulf of Alaska, 1964-1988: portrait of a fishery in transition. *Proc. International Symp. on the Biology and Management of Walleye Pollock*, Lowell Wakefield Fisheries Symp., Alaska Sea Grant Rep. 89-1, 33-58.
- Merati, N. 1993. Spawning dynamics of walleye pollock, *Theragra chalcogramma*, in Shelikof Strait, Gulf of Alaska. Unpublished MS thesis. University of Washington. 134 p.
- Martin, M.H. 1997. Data Report: 1996 Gulf of Alaska bottom trawl survey. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-82, 235 p.

- Methot, R.D. 2000. Technical description of the stock synthesis assessment program. U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-43, 46 p.
- Meuter, F.J. and B.L. Norcross. 2002. Spatial and temporal patterns in the demersal fish community on the shelf and upper slope regions of the Gulf of Alaska. *Fish. Bull.* 100:559-581.
- Mulligan, T.J., Chapman, R.W. and B.L. Brown. 1992. Mitochondrial DNA analysis of walleye pollock, *Theragra chalcogramma*, from the eastern Bering Sea and Shelikof Strait, Gulf of Alaska. *Can. J. Fish. Aquat. Sci.* 49:319-326.
- Olsen, J.B., S.E. Merkouris, and J.E. Seeb. 2002. An examination of spatial and temporal genetic variation in walleye pollock (*Theragra chalcogramma*) using allozyme, mitochondrial DNA, and microsatellite data. *Fish. Bull.* 100:752-764.
- Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *J. Cons. int. Explor. Mer.* 39(2):175-192.
- Press, W.H., S.A. Teukolsky, W.T. Vetterling, and B.P. Flannery. 1992. Numerical recipes in C. Second ed. Cambridge University Press. 994 p.
- Picquelle, S.J., and B.A. Megrey. 1993. A preliminary spawning biomass estimate of walleye pollock, *Theragra chalcogramma*, in Shelikof Strait, Gulf of Alaska, based on the annual egg production method. *Bulletin of Marine Science* 53(2):728:749.
- Ronholt, L. L., H. H. Shippen, and E. S. Brown. 1978. Demersal fish and shellfish resources of the Gulf of Alaska from Cape Spencer to Unimak Pass 1948 - 1976 (A historical review). Northwest and Alaska Fisheries Center Processed Report.
- Saunders, M.W., G.A. McFarlane, and W. Shaw. 1988. Delineation of walleye pollock (*Theragra chalcogramma*) stocks off the Pacific coast of Canada. *Proc. International Symp. on the Biology and Management of Walleye Pollock*, Lowell Wakefield Fisheries Symp., Alaska Sea Grant Rep. 89-1, 379-402.
- Schnute, J.T. and L.J. Richards. 1995. The influence of error on population estimates from catch-age models. *Can. J. Fish. Aquat. Sci.* 52:2063-2077.
- Smith, S. J. 1990. Use of statistical models for the estimation of abundance from groundfish trawl survey data. *Can. J. Fish. Aquat. Sci.* 47:894-903.
- Somerton, D. 1979. Competitive interaction of walleye pollock and Pacific Ocean perch in the northern Gulf of Alaska. *In* S. J. Lipovsky and C.A. Simenstad (eds.) *Gutshop '78, Fish food habits studies: Proceedings of the second Pacific Northwest Technical Workshop, held Maple Valley, WA (USA), 10-13 October, 1978.*, Washington Sea Grant, Seattle, WA.
- StatSci. 1993. S-Plus for DOS reference manual. Statistical Sciences Inc., Seattle, Wash.
- Sullivan, P.J., A.M. Parma, and W.G. Clark. 1997. Pacific halibut assessment: data and methods. *Int. Pac. Halibut Comm. SCI. Rept.* 97. 84 p.
- Swartzman, G., C. Huang, and S. Kaluzny. 1992. Spatial analysis of Bering Sea groundfish survey data using generalized additive models. *Can. J. Fish. Aquat. Sci.* 49:1366-1378.
- von Szalay, P. G., and E. Brown. 2001. Trawl comparisons of fishing power differences and their applicability to National Marine Fisheries Service and the Alaska Department of Fish and Game trawl survey gear. *Alaska Fishery Research Bulletin* 8:85-95.
- Wilson, C. 1994. Echo integration-trawl survey results for pollock in the Gulf of Alaska during 1994. Appendix D of Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.
- Wilson, C., M. Guttormsen, and D. McKelvey. 1995. Echo integration-trawl survey results for pollock in the Gulf of Alaska during 1995. Appendix D of Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Wilson, C.D., M.A. Guttormsen, and S.K. de Blois. 1996. Echo integration-trawl survey results for pollock in the Gulf of Alaska during 1996. In Stock Assessment and Fishery Evaluation Report for Groundfish Resources of the Gulf of Alaska. Prepared by the Gulf of Alaska Groundfish Plan Team, North Pacific Fishery Management Council, P.O. Box 103136, Anchorage, AK 99510. North Pacific Fisheries Management Council, Anchorage, AK.

Yang, M-S. and M. W. Nelson. 2000. Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990, 1993, and 1996. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-112, 174 p.

Table 1.1. Walleye pollock catch (t) in the Gulf of Alaska. The TAC for 2004 is for the area west of 140 ° W lon. (Western, Central and West Yakutat management areas) and includes the guideline harvest level for the state-managed fishery in Prince William Sound (1,700 t). Research catches are also reported.

<i>Year</i>	<i>Foreign</i>	<i>Joint Venture</i>	<i>Domestic</i>	<i>Total</i>	<i>TAC</i>	<i>Research</i>
1964	1,126			1,126	---	
1965	2,749			2,749	---	
1966	8,932			8,932	---	
1967	6,276			6,276	---	
1968	6,164			6,164	---	
1969	17,553			17,553	---	
1970	9,343			9,343	---	
1971	9,458			9,458	---	
1972	34,081			34,081	---	
1973	36,836			36,836	---	
1974	61,880			61,880	---	
1975	59,512			59,512	---	
1976	86,527			86,527	---	
1977	117,834		522	118,356	150,000	89
1978	96,392	34	509	96,935	168,800	100
1979	103,187	566	1,995	105,748	168,800	52
1980	112,997	1,136	489	114,622	168,800	229
1981	130,324	16,857	563	147,744	168,800	433
1982	92,612	73,917	2,211	168,740	168,800	110
1983	81,358	134,131	119	215,608	256,600	213
1984	99,260	207,104	1,037	307,401	416,600	311
1985	31,587	237,860	15,379	284,826	305,000	167
1986	114	62,591	25,103	87,809	116,000	1202
1987		22,823	46,928	69,751	84,000	227
1988		152	65,587	65,739	93,000	19
1989			78,392	78,392	72,200	73
1990			90,744	90,744	73,400	158
1991			100,488	100,488	103,400	16
1992			90,857	90,857	87,400	40
1993			108,908	108,908	114,400	116
1994			107,335	107,335	109,300	70
1995			72,618	72,618	65,360	44
1996			51,263	51,263	54,810	147
1997			90,130	90,130	79,980	56
1998			125,098	125,098	124,730	64
1999			95,590	95,590	94,580	35
2000			73,080	73,080	94,960	56
2001			72,076	72,076	90,690	77
2002			51,937	51,937	53,490	78
2003			50,666	50,666	49,590	38
2004					65,660	
Average (1977-2003)				112,684	128,541	156

Sources: 1964-85--Megrey (1988); 1986-90--Pacific Fishery Information Network (PacFIN), Pacific Marine Fisheries Commission. Domestic catches in 1986-90 were adjusted for discard as described in Hollowed et al. (1991). 1991-2003--NMFS Alaska Regional Office.

Table 1.2. Catch (retained and discarded) of walleye pollock (t) by management area in the Gulf of Alaska during 1992-2003 compiled by the Alaska Regional Office.

Year	Utilization	Shumagin 610	Chirikof 620	Kodiak 630	West Yakutat 640	Prince William Sound 649 (state waters)	Southeast and East Yakutat 650 & 659	Total	Percent discard
1992	Retained	16,014	14,171	47,467	160	0	0	77,812	
	Discarded	3,477	3,066	6,408	73	1	20	13,045	14.4%
	Total	19,490	17,237	53,876	233	1	20	90,857	
1993	Retained	19,791	22,080	58,188	583	0	2	100,645	
	Discarded	1,413	1,708	5,065	65	8	5	8,264	7.6%
	Total	21,204	23,788	63,253	648	8	7	108,908	
1994	Retained	16,238	19,917	58,511	6,362	0	0	101,028	
	Discarded	1,028	2,321	2,453	499	2	3	6,306	5.9%
	Total	17,266	22,239	60,963	6,862	2	3	107,335	
1995	Retained	28,473	11,032	21,989	480	2,739	46	64,759	
	Discarded	1,905	2,048	3,778	53	75	1	7,859	10.8%
	Total	30,378	13,080	25,768	533	2,813	47	72,618	
1996	Retained	23,100	10,150	11,571	510	775	0	46,107	
	Discarded	1,100	2,143	1,789	103	19	3	5,156	10.1%
	Total	24,200	12,293	13,361	613	794	3	51,263	
1997	Retained	25,253	29,736	22,064	3,938	1,807	89	82,888	
	Discarded	1,009	3,179	2,998	30	19	7	7,242	8.0%
	Total	26,262	32,916	25,062	3,968	1,826	96	90,130	
1998	Retained	28,815	48,530	38,753	6,316	1,655	8	124,077	
	Discarded	370	361	262	25	2	0	1,022	0.8%
	Total	29,185	48,892	39,015	6,341	1,657	8	125,098	
1999	Retained	22,864	37,349	29,515	1,737	2,178	1	93,643	
	Discarded	521	784	578	22	39	3	1,947	2.0%
	Total	23,385	38,133	30,093	1,759	2,216	4	95,590	
2000	Retained	21,380	11,314	35,078	1,917	1,181	0	70,870	
	Discarded	694	443	854	191	22	4	2,209	3.0%
	Total	22,074	11,757	35,933	2,108	1,203	4	73,080	
2001	Retained	30,298	17,186	19,942	2,327	1,590	0	71,344	
	Discarded	173	205	330	24	0	0	732	1.0%
	Total	30,471	17,391	20,272	2,351	1,590	0	72,076	
2002	Retained	17,046	20,106	10,615	1,808	1,216	0	50,791	
	Discarded	416	425	287	10	6	2	1,146	2.2%
	Total	17,462	20,531	10,902	1,818	1,222	2	51,937	
2003	Retained	16,347	18,972	12,225	940	1,118	0	49,603	
	Discarded	161	658	210	2	31	0	1,063	2.1%
	Total	16,508	19,630	12,435	943	1,149	0	50,666	
Average (1992-2003)		23,157	23,157	32,578	2,348	1,207	16	82,463	

Table 1.3. Catch at age (000,000s) of walleye pollock in the Gulf of Alaska.

Year	Age															Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
1976	0.00	1.91	24.21	108.69	39.08	16.37	3.52	2.25	1.91	0.31	0.00	0.00	0.00	0.00	0.00	198.25
1977	0.01	2.76	7.06	23.83	89.68	30.35	8.33	2.13	1.79	0.67	0.44	0.10	0.02	0.00	0.00	167.17
1978	0.08	12.11	48.32	18.26	26.39	51.86	12.83	4.18	1.36	1.04	0.32	0.04	0.01	0.00	0.00	176.80
1979	0.00	2.53	48.83	76.37	14.15	10.13	16.70	5.02	1.27	0.60	0.16	0.04	0.00	0.00	0.00	175.81
1980	0.25	19.01	26.50	58.31	36.63	11.31	8.61	8.00	3.89	1.11	0.50	0.21	0.08	0.03	0.00	174.42
1981	0.14	2.59	31.55	73.91	47.97	20.29	4.87	4.83	2.73	0.26	0.03	0.02	0.00	0.00	0.00	189.19
1982	0.01	10.67	55.55	100.77	71.73	54.25	10.46	1.33	0.93	0.55	0.03	0.02	0.02	0.00	0.00	306.31
1983	0.00	3.64	20.64	110.03	137.31	67.41	42.01	7.38	1.24	0.06	0.28	0.07	0.00	0.00	0.00	390.07
1984	0.34	2.37	33.00	38.80	120.80	170.72	62.55	19.31	5.42	0.10	0.07	0.03	0.03	0.00	0.00	453.54
1985	0.04	12.74	5.53	33.22	42.22	86.02	128.95	41.19	10.84	2.20	0.70	0.00	0.00	0.00	0.00	363.64
1986	0.66	8.63	20.34	10.12	19.13	7.32	8.70	9.78	2.13	0.80	0.00	0.00	0.00	0.00	0.00	87.59
1987	0.00	8.83	14.03	8.00	6.89	6.44	7.18	4.19	9.95	1.94	0.00	0.00	0.00	0.00	0.00	67.44
1988	0.17	3.05	20.80	26.95	11.94	5.10	3.45	1.62	0.34	3.21	0.00	0.00	0.00	0.00	0.00	76.62
1989	1.08	0.27	1.47	19.39	28.89	16.96	8.09	4.76	1.69	1.10	3.62	0.43	0.01	0.00	0.00	87.77
1990	0.00	2.77	2.40	2.99	9.49	40.39	13.06	4.90	1.08	0.41	0.01	0.56	0.01	0.07	0.06	78.20
1991	0.00	0.59	9.68	5.45	2.85	5.33	26.67	3.12	16.10	0.87	5.65	0.42	2.19	0.21	0.77	79.90
1992	0.05	3.25	5.57	50.61	14.13	4.02	8.77	19.55	1.02	1.49	0.20	0.73	0.00	0.00	0.00	109.41
1993	0.02	1.97	9.43	21.83	47.46	15.72	6.55	6.29	8.52	1.81	2.07	0.49	0.72	0.13	0.24	123.25
1994	0.06	1.26	4.49	9.63	35.92	31.32	12.20	4.84	4.60	6.15	1.44	1.02	0.29	0.09	0.08	113.37
1995	0.00	0.06	1.01	5.11	11.52	25.83	12.09	2.99	1.52	2.00	1.82	0.19	0.28	0.03	0.15	64.61
1996	0.00	1.27	1.37	1.12	3.50	5.11	12.87	10.60	3.14	1.53	0.80	1.43	0.35	0.23	0.16	43.48
1997	0.00	1.07	6.72	3.77	3.28	6.60	10.09	16.52	12.24	5.06	2.06	0.79	0.54	0.17	0.02	68.92
1998	0.31	0.27	26.44	36.44	15.06	6.65	7.50	11.36	14.96	10.76	3.75	0.75	0.38	0.21	0.11	134.95
1999	0.00	0.42	2.21	22.74	36.10	8.99	6.89	3.72	5.71	7.27	4.01	1.07	0.56	0.12	0.10	99.92
2000	0.08	0.98	2.84	3.47	14.65	24.63	6.24	5.05	2.30	1.24	3.00	1.52	0.30	0.14	0.04	66.48
2001	0.74	10.13	6.59	7.34	9.42	12.59	14.44	4.73	2.70	1.35	0.65	0.83	0.61	0.00	0.04	72.14
2002	0.16	12.31	20.72	6.76	4.47	8.75	5.37	6.06	1.33	0.82	0.43	0.30	0.33	0.22	0.13	68.16
2003	0.14	2.69	21.47	22.95	5.33	3.25	4.66	3.76	2.58	0.54	0.19	0.04	0.09	0.04	0.05	67.79

Table 1.4. Number of aged and measured fish in the Gulf of Alaska domestic pollock fishery used to estimate fishery age composition.

<i>Year</i>	<i>Number aged</i>			<i>Number measured</i>		
	<i>Males</i>	<i>Females</i>	<i>Total</i>	<i>Males</i>	<i>Females</i>	<i>Total</i>
1989	882	892	1,774	6,454	6,456	12,910
1990	453	689	1,142	17,814	24,662	42,476
1991	1,146	1,322	2,468	23,946	39,467	63,413
1992	1,726	1,755	3,481	31,608	47,226	78,834
1993	926	949	1,875	28,035	31,306	59,341
1994	136	129	265	24,321	25,861	50,182
1995	499	544	1,043	10,591	10,869	21,460
1996	381	378	759	8,581	8,682	17,263
1997	496	486	982	8,750	8,808	17,558
1998	924	989	1,913	78,955	83,160	162,115
1999	980	1,115	2,095	16,304	17,964	34,268
2000	1,108	972	2,080	13,167	11,794	24,961
2001	1,063	1,025	2,088	13,731	13,552	27,283
2002	1,036	1,025	2,061	9,924	9,851	19,775
2003	1,091	1,119	2,210	8,375	8,220	16,595

Table 1.5. Biomass estimates (t) of walleye pollock from NMFS echo integration trawl surveys in Shelikof Strait, NMFS bottom trawl surveys (west of 140 W. long.), egg production surveys in Shelikof Strait, and ADF&G crab/groundfish trawl surveys. The biomass of age-1 fish is not included in Shelikof Strait EIT survey estimates in 1995 and 2000 (106,900 and 54,400 t respectively). An adjustment of +1.05% was made to the AFSC bottom trawl biomass time series to account for unsurveyed biomass in Prince William Sound. In 2001, when the NMFS bottom trawl survey did not extend east of 147° W lon., an expansion factor of 2.7% derived from previous surveys was used for West Yakutat.

Year	<i>EIT Shelikof Strait survey</i>		<i>NMFS bottom</i>	<i>Shelikof Strait egg production</i>	<i>ADF&G crab/groundfish survey</i>
	<i>Biosonics</i>	<i>Simrad EK500</i>	<i>trawl west of 140° W lon.</i>		
1981	2,785,755			1,788,908	
1982					
1983	2,278,172				
1984	1,757,168		719,937		
1985	1,175,823			768,419	
1986	585,755			375,907	
1987			732,541	484,455	
1988	301,709			504,418	
1989	290,461			433,894	214,434
1990	374,731		825,592	381,475	114,451
1991	380,331			370,000	
1992	580,000	681,400		616,000	127,359
1993	295,785	408,200	754,390		132,849
1994		467,300			103,420
1995		618,300			
1996		745,400	665,745		122,477
1997		570,100			93,728
1998		489,900			81,215
1999			607,147		53,587
2000		334,900			102,871
2001		369,600	216,777		86,967
2002		229,100			96,237
2003		270,200	399,690		66,989
2004		291,215			99,358

Table 1.6. Survey sampling effort and biomass coefficients of variation (CV) for pollock in the Gulf of Alaska bottom trawl survey and the Shelikof Strait EIT survey. For the Shelikof Strait EIT survey, CVs are reported for 1981-91, while relative estimation error using a geostatistical method are reported for 1992-2003. For the Gulf of Alaska bottom trawl survey, the number of measured pollock is approximate due to subsample expansions in the database, and the total number measured includes both sexed and unsexed fish.

Year	No. of tows	No. of tows with pollock		Survey biomass CV		Number aged		Number measured		Total
		No. of tows	pollock	Males	Females	Males	Females	Males	Females	
Bottom trawl survey										
1984	929	536	0.14	1,119	1,394	2,513	8,979	13,286	24,064	
1987	783	533	0.20	672	675	1,347	8,101	15,654	24,608	
1990	708	549	0.12	503	560	1,063	13,955	18,967	35,355	
1993	775	628	0.16	879	1,013	1,892	14,496	18,692	34,921	
1996	807	668	0.15	509	560	1,069	14,653	15,961	34,526	
1999	764	567	0.38	560	613	1,173	10,808	11,314	24,080	
2001	489	302	0.30	395	519	914	NA	NA	NA	
2003	807	508	0.12	514	589	1,103	NA	NA	NA	
Shelikof Strait EIT survey										
	No. of midwater tows	No. of bottom trawl tows								
1981	36	18	0.12	1,921	1,815	3,736	NA	NA	NA	
1983	47	1	0.16	1,642	1,103	2,745	NA	NA	NA	
1984	42	0	0.18	1,739	1,622	3,361	NA	NA	NA	
1985	57	0	0.14	1,055	1,187	2,242	NA	NA	NA	
1986	38	1	0.22	642	618	1,260	NA	NA	NA	
1987	27	0	---	557	643	1,200	NA	NA	NA	
1988	26	0	0.17	537	464	1,001	NA	NA	NA	
1989	21	0	0.10	757	796	1,553	NA	NA	NA	
1990	25	16	0.17	988	1,117	2,105	NA	NA	NA	
1991	16	2	0.35	478	628	1,106	NA	NA	NA	
1992	17	8	0.04	784	765	1,549	NA	NA	NA	
1993	22	2	0.05	583	624	1,207	NA	NA	NA	
1994	42	12	0.05	554	633	1,187	NA	NA	NA	
1995	22	3	0.05	599	575	1,174	NA	NA	NA	
1996	30	8	0.04	724	775	1,499	NA	NA	NA	
1997	16	14	0.04	682	853	1,535	NA	NA	NA	
1998	22	9	0.04	863	784	1,647	NA	NA	NA	
2000	31	0	0.05	430	370	800	NA	NA	NA	
2001	15	9	0.05	314	378	692	NA	NA	NA	
2002	18	1	0.07	278	326	604	NA	NA	NA	
2003	17	2	0.05	294	322	616	NA	NA	NA	
2004	13	2	0.09	422	315	737	NA	NA	NA	

Table 1.7. Estimated number at age (000,000s) from the echo integration-trawl survey in Shelikof Strait, and from the NMFS bottom trawl survey. For the acoustic survey in 1987, when total abundance could not be estimated, the percent at age is given. Bottom trawl survey estimates are for the Western and Central Gulf of Alaska only (Management areas 610-630).

<i>Gulf of Alaska bottom trawl survey</i>																
<i>Year</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>6</i>	<i>7</i>	<i>8</i>	<i>9</i>	<i>10</i>	<i>11</i>	<i>12</i>	<i>13</i>	<i>14</i>	<i>15</i>	<i>Total</i>
1984	0.93	10.02	67.81	155.78	261.17	474.57	145.10	24.80	16.59	1.66	0.21	1.32	0.00	0.00	0.00	1159.96
1987	25.45	363.02	172.99	138.97	91.13	168.27	78.14	43.99	175.39	22.41	7.81	3.51	1.82	0.00	0.00	1292.88
1989	208.88	63.49	47.56	243.15	301.09	104.43	54.47	28.39	26.14	5.98	10.66	0.00	0.00	0.00	0.00	1094.23
1990	64.04	251.21	48.34	46.68	209.77	240.82	74.41	110.41	26.13	34.23	5.03	27.73	5.70	1.07	1.63	1147.19
1993	139.31	71.15	50.94	182.96	267.12	91.51	33.12	68.98	76.62	26.36	11.85	6.29	3.82	1.82	4.41	1036.25
1996	194.23	128.79	17.30	26.13	50.04	63.18	174.41	87.62	52.37	27.73	12.10	18.46	7.16	9.68	19.70	888.90
1999	109.73	19.17	20.94	66.76	118.94	56.80	59.04	47.71	56.40	81.97	65.18	9.67	8.28	2.50	0.76	723.85
2001	412.83	117.03	34.42	33.39	25.05	33.45	37.01	8.20	5.74	0.59	4.48	2.52	1.28	0.00	0.18	716.19
2003	75.46	18.40	128.41	140.74	73.27	44.72	36.10	25.27	14.51	8.61	3.23	1.79	1.26	0.00	0.00	571.77
<i>Shelikof Strait EIT survey</i>																
<i>Year</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>6</i>	<i>7</i>	<i>8</i>	<i>9</i>	<i>10</i>	<i>11</i>	<i>12</i>	<i>13</i>	<i>14</i>	<i>15</i>	<i>Total</i>
1981	77.65	3,481.18	1,510.77	769.16	2,785.91	1,051.92	209.93	128.52	79.43	25.19	1.73	0.00	0.00	0.00	0.00	10,121.37
1983	1.21	901.77	380.19	1,296.79	1,170.81	698.13	598.78	131.54	14.48	11.61	3.92	1.71	0.00	0.00	0.00	5,210.93
1984	61.65	58.25	324.49	141.66	635.04	988.21	449.62	224.35	41.03	2.74	0.00	1.02	0.00	0.00	0.00	2,928.07
1985	2,091.74	544.44	122.69	314.77	180.53	347.17	439.31	166.68	42.72	5.56	1.77	1.29	0.00	0.00	0.00	4,258.67
1986	575.36	2,114.83	183.62	45.63	75.36	49.34	86.15	149.36	60.22	10.62	1.29	0.00	0.00	0.00	0.00	3,351.78
1987	7.5%	25.5%	55.8%	2.9%	1.7%	1.2%	1.6%	1.2%	2.1%	0.4%	0.1%	0.0%	0.0%	0.0%	0.0%	100.0%
1988	17.44	109.93	694.32	322.11	77.57	16.99	5.70	5.60	3.98	8.96	1.78	1.84	0.20	0.00	0.00	1,266.41
1989	399.48	89.52	90.01	222.05	248.69	39.41	11.75	3.83	1.89	0.55	10.66	1.42	0.00	0.00	0.00	1,119.25
1990	49.14	1,210.17	71.69	63.37	115.92	180.06	46.33	22.44	8.20	8.21	0.93	3.08	1.51	0.79	0.24	1,782.08
1991	21.98	173.65	549.90	48.11	64.87	69.60	116.32	23.65	29.43	2.23	4.29	0.92	4.38	0.00	0.00	1,109.32
1994	155.71	30.33	42.97	29.31	146.27	79.07	40.47	25.98	42.66	46.46	14.22	6.40	1.08	2.25	0.55	663.72
1995	10,000.00	467.55	71.97	71.72	98.51	235.25	116.74	51.36	15.96	10.30	13.98	5.57	2.04	0.42	0.00	11,161.37
1996	51.50	3,193.33	110.73	23.75	51.72	68.32	193.46	114.14	38.40	12.53	10.93	5.13	2.42	0.02	0.37	3,876.75
1997	66.42	179.05	1,230.48	77.54	17.69	42.98	50.48	95.27	51.52	13.96	2.34	2.97	0.91	0.45	0.00	1,832.04
1998	390.12	85.49	123.98	467.34	133.52	13.64	30.44	34.55	70.48	24.64	13.63	6.56	0.26	0.54	0.54	1,395.74
2000	4,275.17	621.45	180.36	13.61	58.41	114.11	14.63	10.95	8.53	6.79	12.05	5.99	1.67	0.92	0.00	5,324.66
2001	272.48	3,591.22	296.13	51.47	34.83	18.99	28.53	10.81	5.10	2.20	1.00	1.55	0.57	0.41	0.20	4,315.50
2002	6.01	137.88	1,023.82	86.05	13.21	12.98	6.15	5.41	1.16	0.51	0.28	0.27	0.12	0.10	0.00	1,293.95
2003	45.19	64.26	187.48	705.94	46.12	5.43	2.92	1.01	1.31	0.57	0.19	0.00	0.12	0.00	0.00	1,060.54
2004	8.39	62.01	49.28	143.85	320.69	44.12	2.31	2.21	2.52	0.45	0.40	0.00	0.33	0.00	0.00	636.56

Table 1.8. Estimates of pollock biomass obtained from GLM model predictions of pollock CPUE and INPFC area expansions. Biomass estimates were multiplied by the von Szalay and Brown (2001) FPC of 3.84 for comparison to the NMFS triennial trawl survey biomass estimates. Coefficients of variation do not reflect the variance of the FPC estimate.

<i>Year</i>	<i>Biomass (t)</i>	<i>FPC-adjusted</i>	<i>biomass (t)</i>	<i>CV</i>
1961	50,356		193,369	0.24
1962	57,496		220,783	0.30
1970	7,979		30,640	0.42
1971	4,257		16,348	0.64
1974	1,123,447		4,314,035	0.38
1975	1,501,142		5,764,384	0.52
1978	223,277		857,383	0.31
1980	146,559		562,787	0.27
1981	257,219		987,719	0.33
1982	356,433		1,368,703	0.29

Other published estimates of pollock biomass from surveys using 400-mesh eastern trawls

<i>Year</i>	<i>Biomass (t)</i>	<i>Source</i>
1961	57,449	<i>Ronholt et al. 1978</i>
1961-62	91,075	<i>Ronholt et al. 1978</i>
1973-75	1,055,000	<i>Alton et al. 1977</i>
1973-76	739,293	<i>Ronholt et al. 1978</i>
1973-75	610,413	<i>Hughes and Hirschhorn 1979</i>

Table 1.9. Predictions of Gulf of Alaska pollock year-class strength. The FOCI prediction is the prediction of year-class strength made in the natal year of the year class, and was derived from environmental indices, larval surveys, and the time series characteristics of pollock recruitment. The McKelvey index is the estimated abundance of 9-16 cm pollock from the Shelikof Strait EIT survey.

<i>Year class</i>	<i>FOCI prediction</i>	<i>Year of EIT survey</i>	<i>McKelvey index</i>	<i>Rank abundance of McKelvey index</i>
1980		1981	0.078	10
1981				
1982		1983	0.001	21
1983		1984	0.062	12
1984		1985	2.092	3
1985		1986	0.579	4
1986				
1987		1988	0.017	18
1988		1989	0.399	5
1989		1990	0.049	15
1990		1991	0.022	17
1991		1992	0.153	9
1992	Strong	1993	0.054	14
1993	Average	1994	0.156	8
1994	Average	1995	10.004	1
1995	Average-Strong	1996	0.056	13
1996	Average	1997	0.066	11
1997	Average	1998	0.390	6
1998	Average			
1999	Average	2000	4.275	2
2000	Average	2001	0.274	7
2001	Average-Strong	2002	0.006	20
2002	Average	2003	0.045	16
2003	Average	2004	0.008	19
2004	Average		---	---

Table 1.10. Ageing error transition matrix used in the Gulf of Alaska pollock assessment model.

<i>True Age</i>	<i>St. dev.</i>	<i>Observed Age</i>										
		<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>6</i>	<i>7</i>	<i>8</i>	<i>9</i>	<i>10</i>	
1	0.18	0.9970	0.0030	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2	0.23	0.0138	0.9724	0.0138	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
3	0.27	0.0000	0.0329	0.9342	0.0329	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
4	0.32	0.0000	0.0000	0.0571	0.8858	0.0571	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
5	0.36	0.0000	0.0000	0.0000	0.0832	0.8335	0.0832	0.0000	0.0000	0.0000	0.0000	0.0000
6	0.41	0.0000	0.0000	0.0000	0.0001	0.1090	0.7817	0.1090	0.0001	0.0000	0.0000	0.0000
7	0.45	0.0000	0.0000	0.0000	0.0000	0.0004	0.1333	0.7325	0.1333	0.0004	0.0000	0.0000
8	0.50	0.0000	0.0000	0.0000	0.0000	0.0000	0.0012	0.1554	0.6868	0.1554	0.0012	0.0000
9	0.54	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0028	0.1747	0.6450	0.1775	0.0000
10	0.59	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0052	0.1913	0.8035	0.0000

Table 1.11. Maturity at age of female pollock derived from maturity stage data collected during winter EIT surveys in the Gulf of Alaska (1983-2004).

Year	2		3		4		5		6		7		8		9		10+		Total
	Mat.	Tot.																	
1983	0	145	19	115	284	356	291	303	189	194	171	174	33	35	7	7	4	4	1333
1984	0	39	25	173	97	141	349	364	507	512	237	237	132	133	21	21	1	1	1621
1985	3	204	4	79	75	177	53	102	182	196	261	263	122	123	30	30	9	9	1183
1986	0	93	1	48	6	57	62	73	46	51	71	74	151	151	57	57	14	14	618
1987	0	39	2	171	5	47	18	53	30	39	69	78	57	60	116	117	34	34	638
1988	0	49	0	136	24	115	12	68	20	33	10	15	13	13	6	7	27	28	464
1989	0	35	0	50	52	175	122	276	71	100	57	62	16	16	12	12	70	70	796
1990	0	86	0	109	19	99	182	270	468	620	202	222	103	109	58	60	268	269	1844
1991	0	47	0	159	3	27	7	85	34	60	89	111	19	22	45	46	71	71	628
1992	0	12	0	43	5	126	20	291	41	53	53	54	104	105	23	23	57	58	765
1993	0	38	1	62	6	50	59	127	48	112	37	46	61	63	58	58	67	68	624
1994	0	43	1	144	27	64	230	247	64	68	41	46	38	39	84	84	137	137	872
1995	0	147	0	61	13	85	63	88	231	239	90	92	35	38	11	12	42	43	805
1996	0	61	0	89	1	28	43	60	78	85	198	203	131	136	55	55	44	46	763
1997	0	11	0	111	7	29	19	25	123	123	135	135	234	235	125	125	49	49	843
1998	0	69	0	72	14	215	13	64	15	18	53	55	65	65	112	112	86	87	757
2000	0	29	1	81	1	8	36	57	78	100	11	19	11	13	10	10	36	39	356
2001	0	44	0	57	13	45	16	52	33	40	69	73	29	30	13	14	19	19	374
2002	0	11	2	77	15	58	51	68	84	90	76	78	83	83	13	13	21	21	499
2003	0	40	1	34	29	151	12	31	9	17	10	11	3	4	8	8	5	5	301
2004	0	30	0	24	58	104	149	219	35	47	2	3	7	7	6	6	4	4	444

Proportion mature

	2	3	4	5	6	7	8	9	10+
1983	0.000	0.165	0.798	0.960	0.974	0.983	0.943	1.000	1.000
1984	0.000	0.145	0.688	0.959	0.990	1.000	0.992	1.000	1.000
1985	0.015	0.051	0.424	0.520	0.929	0.992	0.992	1.000	1.000
1986	0.000	0.021	0.105	0.849	0.902	0.959	1.000	1.000	1.000
1987	0.000	0.012	0.106	0.340	0.769	0.885	0.950	0.991	1.000
1988	0.000	0.000	0.209	0.176	0.606	0.667	1.000	0.857	0.964
1989	0.000	0.000	0.297	0.442	0.710	0.919	1.000	1.000	1.000
1990	0.000	0.000	0.192	0.674	0.755	0.910	0.945	0.967	0.996
1991	0.000	0.000	0.111	0.082	0.567	0.802	0.864	0.978	1.000
1992	0.000	0.000	0.040	0.069	0.774	0.981	0.990	1.000	0.983
1993	0.000	0.016	0.120	0.465	0.429	0.804	0.968	1.000	0.985
1994	0.000	0.007	0.422	0.931	0.941	0.891	0.974	1.000	1.000
1995	0.000	0.000	0.153	0.716	0.967	0.978	0.921	0.917	0.977
1996	0.000	0.000	0.036	0.717	0.918	0.975	0.963	1.000	0.957
1997	0.000	0.000	0.241	0.760	1.000	1.000	0.996	1.000	1.000
1998	0.000	0.000	0.065	0.203	0.833	0.964	1.000	1.000	0.989
2000	0.000	0.012	0.125	0.632	0.780	0.579	0.846	1.000	0.923
2001	0.000	0.000	0.289	0.308	0.825	0.945	0.967	0.929	1.000
2002	0.000	0.026	0.259	0.750	0.933	0.974	1.000	1.000	1.000
2003	0.000	0.029	0.192	0.387	0.529	0.909	0.750	1.000	1.000
2004	0.000	0.000	0.558	0.680	0.745	0.667	1.000	1.000	1.000
Averages									
All years	0.001	0.023	0.259	0.553	0.804	0.895	0.955	0.983	0.989
1994-2004	0.000	0.007	0.234	0.608	0.847	0.888	0.942	0.985	0.984
1999-2004	0.000	0.014	0.284	0.551	0.762	0.815	0.913	0.986	0.985

Table 1.12. Results comparing model fits, stock status, and 2005 yield for different model configurations.

	<i>Model 1</i>	<i>Model 2a</i>	<i>Model 2b</i>	<i>Model 2c</i>	<i>Model 3</i>	<i>Model 4</i>
Model fits						
Total -log(Likelihood)	868.48	869.25	869.25	869.25	854.48	702.73
NMFS trawl q	0.80	1.00	1.00	1.00	1.00	1.00
Age composition data						
Fishery effective N	270	267	267	267	267	272
NMFS bottom trawl effective N	79	78	78	78	86	69
Shelikof Strait EIT effective N	33	32	32	32	32	25
Length composition data						
ADF&G trawl effective N	32	32	32	32	33	33
Historical trawl survey effective N	20	20	20	20	20	20
Survey abundance						
NMFS bottom trawl RMSE	0.418	0.419	0.419	0.419	0.428	0.364
Shelikof Strait EIT RMSE	0.367	0.363	0.363	0.363	0.364	0.426
ADF&G trawl RMSE	0.244	0.250	0.250	0.250	0.271	0.238
Historical trawl survey RMSE	1.523	1.523	1.523	1.523	1.522	1.525
Egg production survey RMSE	0.461	0.445	0.445	0.445	0.447	0.442
Stock status (t)						
2005 Spawning biomass	242,824	213,187	215,329	183,165	200,880	240,819
(CV)	(16%)	(11%)	(11%)	(9%)	(12%)	(12%)
2005 3+ biomass	825,733	736,202	736,202	641,166	697,380	861,539
(CV)	(16%)	(12%)	(12%)	(10%)	(13%)	(14%)
Depletion (B2005/B0)	38%	37%	38%	32%	35%	42%
B _{40%}	253,787	229,099	229,099	229,099	228,067	231,005
2005 yield (000 t)						
F_{OFL}	165.78	144.34	144.34	105.53	129.96	179.77
MaxFABC	143.43	124.80	124.80	90.89	112.23	154.53
Author's F	122.54	106.53	78.98	77.17	95.63	136.75

Comments:

$$RMSE = \sqrt{\frac{\sum \ln(obs / pred)^2}{n}}$$

Model descriptions (see text for details):

- Model 1--Estimated NMFS trawl survey catchability
- Model 2a--Last year's model configuration
- Model 2b--F50% in 2005 & 2006
- Model 2c--1999 year class set to average
- Model 3--Reduced emphasis on ADF&G survey
- Model 4--Reduced emphasis on Shelikof EIT survey

Table 1.13. Estimated selectivity at age for Gulf of Alaska pollock fisheries and surveys. The fisheries and surveys were modeled using double logistic selectivity functions, with random walk process error for the fishery logistic parameters. Fishery selectivity at age reported below is the average of the annual selectivity for the indicated time period, rescaled so that the maximum is one.

Age	Early				400-mesh			
	POP fishery (1961-71)	Foreign (1972- 84)	domestic (1985-91)	Recent domestic (1992-2004)	EIT survey	Bottom trawl survey	ADF&G bottom trawl	eastern trawl 1961-82
2	0.001	0.041	0.043	0.059	1.000	0.176	0.058	0.120
3	0.021	0.263	0.158	0.157	0.998	0.282	0.130	0.388
4	0.413	0.756	0.424	0.381	0.993	0.444	0.268	0.747
5	1.000	1.000	0.741	0.682	0.976	0.666	0.473	0.932
6	0.946	0.920	0.954	0.886	0.932	0.896	0.689	0.985
7	0.700	0.670	1.000	0.972	0.819	1.000	0.849	0.997
8	0.362	0.339	0.830	1.000	0.603	0.886	0.939	0.999
9	0.133	0.126	0.431	0.941	0.338	0.652	0.981	1.000
10	0.041	0.041	0.150	0.369	0.147	0.432	1.000	1.000

Table 1.14. Total estimated abundance at age (numbers in 000,000s) of Gulf of Alaska pollock from the age-structured assessment model.

	<i>Age</i>								
	2	3	4	5	6	7	8	9	10
1961	373	195	120	74	55	39	28	21	16
1962	413	277	145	89	55	41	29	21	28
1963	443	306	205	107	66	41	30	21	36
1964	99	328	227	152	79	49	30	22	42
1965	257	73	243	168	112	59	36	22	48
1966	136	191	54	179	123	82	43	26	52
1967	340	101	141	40	127	87	59	31	58
1968	402	252	75	103	28	91	63	43	66
1969	700	298	187	54	73	20	65	46	80
1970	330	519	220	130	35	47	13	46	93
1971	715	244	384	159	91	24	33	10	102
1972	1,343	529	181	279	112	64	18	24	83
1973	1,026	995	391	129	187	76	45	13	79
1974	3,363	760	735	279	86	126	53	32	67
1975	685	2,491	561	520	180	56	86	37	73
1976	433	507	1,823	393	361	126	40	62	81
1977	2,000	320	366	1,264	272	252	89	29	105
1978	2,741	1,479	233	252	861	187	177	64	99
1979	2,507	2,024	1,062	160	173	595	132	127	120
1980	3,523	1,853	1,465	732	109	119	420	95	182
1981	1,790	2,600	1,343	1,026	507	76	84	302	204
1982	435	1,323	1,891	937	703	349	53	60	371
1983	493	319	942	1,312	647	488	246	38	319
1984	206	362	226	636	872	433	337	177	264
1985	473	150	249	141	380	523	274	235	325
1986	1,594	345	104	152	78	203	290	175	408
1987	539	1,164	242	68	95	48	128	200	430
1988	156	396	839	167	45	61	31	85	460
1989	361	115	287	587	112	29	40	20	399
1990	1,602	267	84	205	398	72	19	25	306
1991	1,041	1,184	196	61	140	254	44	11	242
1992	418	769	869	141	42	92	163	28	167
1993	239	308	559	606	93	27	59	104	140
1994	141	176	224	391	402	60	17	37	167
1995	213	104	128	158	264	264	39	11	141
1996	830	157	76	92	109	179	179	26	107
1997	404	614	115	55	64	75	122	122	93
1998	174	298	448	82	37	41	47	76	138
1999	182	127	209	288	49	21	23	27	131
2000	272	134	91	140	176	28	12	13	100
2001	1,196	201	97	64	91	105	16	7	75
2002	746	876	144	67	41	55	63	9	56
2003	107	545	628	100	45	27	35	40	46
2004	127	78	388	437	68	30	17	23	60
Average	808	599	436	299	196	130	87	60	151

Table 1.15. Estimates of population biomass, recruitment, and harvest of Gulf of Alaska pollock from the age-structured assessment model. The harvest rate is the catch in biomass divided by the total biomass of age 3+ fish at the start of the year.

Year	2+ total		Female		Age 2		2003 Assessment results				
	biomass (1,000 t)	3+ total biomass (1,000 t)	spawn. biom.	spawn. biom.	recruits (million)	Catch (t)	Harvest rate	3+ total biomass	Female spawn. biom.	Age 2 recruits	Harvest rate
1969	701	598	143	143	700	17,553	3%	589	140	686	3%
1970	764	715	140	140	330	9,343	1%	704	138	322	1%
1971	857	752	156	156	715	9,458	1%	738	152	694	1%
1972	1,060	863	174	174	1,343	34,081	4%	844	170	1,298	4%
1973	1,295	1,144	192	192	1,026	36,836	3%	1,114	187	991	3%
1974	1,854	1,360	226	226	3,363	61,880	5%	1,320	219	3,244	5%
1975	2,275	2,174	280	280	685	59,512	3%	2,102	270	661	3%
1976	2,375	2,311	391	391	433	86,527	4%	2,231	373	418	4%
1977	2,409	2,115	490	490	2,000	118,356	6%	2,037	470	1,938	6%
1978	2,684	2,281	531	531	2,741	96,935	4%	2,196	511	2,671	4%
1979	3,150	2,781	541	541	2,507	105,748	4%	2,684	519	2,482	4%
1980	3,769	3,251	598	598	3,523	114,622	4%	3,154	569	3,505	4%
1981	4,171	3,908	488	488	1,790	147,744	4%	3,819	467	1,788	4%
1982	4,099	4,035	559	559	435	168,740	4%	3,959	538	437	4%
1983	3,480	3,407	681	681	493	215,608	6%	3,351	708	493	6%
1984	2,784	2,753	712	712	206	307,401	11%	2,710	753	207	11%
1985	2,107	2,037	644	644	473	284,826	14%	2,003	646	474	14%
1986	1,898	1,638	523	523	1,594	87,809	5%	1,608	590	1,601	5%
1987	1,783	1,696	440	440	539	69,751	4%	1,675	489	540	4%
1988	1,623	1,597	398	398	156	65,739	4%	1,583	393	157	4%
1989	1,506	1,446	385	385	361	78,392	5%	1,435	355	362	5%
1990	1,498	1,229	345	345	1,602	90,744	7%	1,222	383	1,603	7%
1991	1,518	1,344	326	326	1,041	100,488	7%	1,339	345	1,039	8%
1992	1,737	1,669	287	287	418	90,857	5%	1,666	284	416	5%
1993	1,562	1,523	323	323	239	108,908	7%	1,520	319	237	7%
1994	1,301	1,278	372	372	141	107,335	8%	1,275	371	139	8%
1995	1,096	1,076	343	343	213	72,618	7%	1,072	343	208	7%
1996	969	889	311	311	830	51,263	6%	885	310	806	6%
1997	940	901	268	268	404	90,130	10%	891	267	383	10%
1998	848	823	203	203	174	125,098	15%	806	201	153	16%
1999	690	664	184	184	182	95,590	14%	642	180	185	15%
2000	625	591	171	171	272	73,080	12%	568	165	308	13%
2001	704	579	167	167	1,196	72,076	12%	567	160	1,385	13%
2002	947	832	144	144	746	51,937	6%	891	139	682	6%
2003	980	959	143	143	107	50,666	5%	993	140	118	5%
2004	866	841	168	168	127	---	---	---	---	---	---
Average											
1969-2004	1,748	1,613	346	346	920	95,933	6%	1,606	350	932	6%
1979-2003					786						

Table 1.16. Gulf of Alaska pollock life history and fishery vectors used to estimate spawning biomass per recruit (F_{SPR}) harvest rates. Population weight at age is the average for the bottom trawl survey in 1999-2003. Proportion mature females is the average for 1983-2004 from winter EIT survey specimen data. Spawning weight at age is the average for the Shelikof Strait EIT survey in 2000-2004.

Age	Natural mortality	Fishery selectivity (Avg. 1992-2004)	Weight at age (kg)			Proportion mature females
			Spawning (March 15)	Population (June-Aug.)	Fishery (Avg. 1999-2003)	
2	0.3	0.059	0.075	0.150	0.329	0.001
3	0.3	0.157	0.203	0.421	0.499	0.023
4	0.3	0.381	0.386	0.630	0.664	0.259
5	0.3	0.682	0.583	0.800	0.875	0.553
6	0.3	0.886	0.850	0.943	1.046	0.804
7	0.3	0.972	1.144	1.078	1.191	0.895
8	0.3	1.000	1.267	1.221	1.329	0.955
9	0.3	0.941	1.455	1.312	1.440	0.983
10+	0.3	0.369	1.600	1.445	1.597	0.989

Table 1.17. Methods used to assess Gulf of Alaska pollock, 1977-2003. The basis for catch recommendation in 1977-1989 is the presumptive method by which the TAC was determined (based on the assessment and SSC minutes). The basis for catch recommendation given in 1990-2003 is the method used by the Plan Team to derive the ABC recommendation given in the SAFE summary chapter.

<i>Year</i>	<i>Assessment method</i>	<i>Basis for catch recommendation in following year</i>	<i>B40% (t)</i>
1977	Survey biomass, CPUE trends, $M=0.4$	$MSY = 0.4 * M * B_{zero}$	---
1978	Survey biomass, CPUE trends	$MSY = 0.4 * M * B_{zero}$	---
1979	Survey biomass, CPUE trends	$MSY = 0.4 * M * B_{zero}$	---
1980	Survey biomass, CPUE trends	$MSY = 0.4 * M * B_{zero}$	---
1981	Survey biomass, CPUE trends	$MSY = 0.4 * M * B_{zero}$	---
1982	CAGEAN	$MSY = 0.4 * M * B_{zero}$	---
1983	CAGEAN	Mean annual surplus production	---
1984	Projection of survey numbers at age	Stabilize biomass trend	---
1985	CAGEAN, projection of survey numbers at age, CPUE trends	Stabilize biomass trend	---
1986	CAGEAN, projection of survey numbers at age	Stabilize biomass trend	---
1987	CAGEAN, projection of survey numbers at age	Stabilize biomass trend	---
1988	CAGEAN, projection of survey numbers at age	10% of exploitable biomass	---
1989	Stock synthesis	10% of exploitable biomass	---
1990	Stock synthesis, reduce M to 0.3	10% of exploitable biomass	---
1991	Stock synthesis, assume trawl survey catchability = 1	FMSY from an assumed SR curve	---
1992	Stock synthesis	$\text{Max}[-\text{Pr}(\text{SB} < \text{Threshold}) + \text{Yld}]$	---
1993	Stock synthesis	$\text{Pr}(\text{SB} > \text{B20}) = 0.95$	---
1994	Stock synthesis	$\text{Pr}(\text{SB} > \text{B20}) = 0.95$	---
1995	Stock synthesis	$\text{Max}[-\text{Pr}(\text{SB} < \text{Threshold}) + \text{Yld}]$	---
1996	Stock synthesis	Amendment 44 Tier 3 guidelines	289,689
1997	Stock synthesis	Amendment 44 Tier 3 guidelines	267,600
1998	Stock synthesis	Amendment 44 Tier 3 guidelines	240,000
1999	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F_{ABC})	247,000
2000	AD model builder	Amendment 56 Tier 3 guidelines	250,000
2001	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F_{ABC})	245,000
2002	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F_{ABC})	240,000
2003	AD model builder	Amendment 56 Tier 3 guidelines (with a reduction from max permissible F_{ABC})	248,000

Table 1.18. Projections of Gulf of Alaska pollock spawning biomass, full recruitment fishing mortality, and catch for 2005-2017 under different harvest policies. All projections begin with estimated age composition in 2005 using base run model. Coefficients of variation are given in parentheses, and reflex only variability in recruitment in 2006-2017. The values for $B_{100\%}$, $B_{40\%}$, and $B_{35\%}$ are 573,000, 229,000, and 200,000 t, respectively.

Spawning biomass (t)	F_{OFL}		$Max F_{ABC}$		Author's recommended F		Author's recommended F with $F_{50\%}$ in 2005 & 2006		50% of max FABC		Average F		$F = 0$	
	2005	210,060	(0.00)	211,660	(0.00)	213,120	(0.00)	215,260	(0.00)	216,280	(0.00)	214,660	(0.00)	221,000
2006	183,730	(0.00)	191,460	(0.00)	198,840	(0.00)	210,230	(0.00)	215,920	(0.00)	206,670	(0.00)	244,920	(0.00)
2007	148,080	(0.01)	157,600	(0.01)	167,210	(0.01)	181,310	(0.01)	191,440	(0.01)	176,260	(0.01)	240,340	(0.01)
2008	137,410	(0.16)	146,610	(0.15)	156,270	(0.15)	165,100	(0.14)	182,090	(0.13)	161,920	(0.15)	242,450	(0.10)
2009	152,650	(0.32)	161,710	(0.31)	171,230	(0.30)	176,610	(0.29)	198,100	(0.29)	174,030	(0.32)	266,600	(0.24)
2010	178,220	(0.39)	188,430	(0.39)	198,120	(0.38)	201,090	(0.37)	229,060	(0.39)	202,140	(0.43)	306,670	(0.36)
2011	197,680	(0.38)	210,370	(0.39)	220,410	(0.38)	221,870	(0.37)	260,050	(0.42)	230,290	(0.46)	352,440	(0.41)
2012	207,530	(0.37)	222,910	(0.38)	233,130	(0.36)	233,790	(0.36)	283,540	(0.41)	251,200	(0.44)	395,150	(0.41)
2013	212,240	(0.36)	229,890	(0.37)	240,140	(0.35)	240,420	(0.35)	301,210	(0.40)	266,780	(0.43)	434,030	(0.40)
2014	215,460	(0.36)	234,700	(0.37)	244,880	(0.35)	244,990	(0.35)	314,930	(0.39)	279,090	(0.42)	467,440	(0.39)
2015	217,860	(0.36)	238,120	(0.37)	248,210	(0.35)	248,250	(0.35)	325,050	(0.39)	288,460	(0.41)	493,250	(0.38)
2016	218,940	(0.36)	239,980	(0.37)	249,880	(0.35)	249,900	(0.35)	331,930	(0.39)	294,980	(0.41)	512,740	(0.38)
2017	219,540	(0.36)	241,090	(0.37)	250,830	(0.35)	250,840	(0.35)	336,760	(0.39)	299,620	(0.41)	527,690	(0.37)

Fishing mortality	F_{OFL}		$Max F_{ABC}$		Author's recommended F		Author's recommended F with $F_{50\%}$ in 2005 & 2006		50% of max FABC		Average F		$F = 0$	
	2005	0.32	(0.00)	0.27	(0.00)	0.23	(0.00)	0.17	(0.00)	0.14	(0.00)	0.18	(0.00)	0
2006	0.28	(0.00)	0.24	(0.00)	0.21	(0.00)	0.16	(0.00)	0.14	(0.00)	0.18	(0.00)	0	---
2007	0.22	(0.01)	0.20	(0.01)	0.18	(0.01)	0.19	(0.01)	0.12	(0.01)	0.18	(0.00)	0	---
2008	0.20	(0.15)	0.18	(0.14)	0.16	(0.14)	0.17	(0.13)	0.11	(0.10)	0.18	(0.00)	0	---
2009	0.22	(0.25)	0.20	(0.23)	0.18	(0.25)	0.18	(0.23)	0.12	(0.16)	0.18	(0.00)	0	---
2010	0.25	(0.26)	0.22	(0.24)	0.20	(0.28)	0.20	(0.27)	0.13	(0.17)	0.18	(0.00)	0	---
2011	0.27	(0.24)	0.24	(0.23)	0.22	(0.26)	0.22	(0.26)	0.13	(0.16)	0.18	(0.00)	0	---
2012	0.28	(0.22)	0.25	(0.20)	0.23	(0.24)	0.23	(0.24)	0.14	(0.14)	0.18	(0.00)	0	---
2013	0.29	(0.21)	0.25	(0.19)	0.24	(0.22)	0.24	(0.22)	0.14	(0.12)	0.18	(0.00)	0	---
2014	0.29	(0.19)	0.26	(0.17)	0.24	(0.21)	0.24	(0.21)	0.14	(0.10)	0.18	(0.00)	0	---
2015	0.29	(0.19)	0.26	(0.17)	0.24	(0.20)	0.24	(0.20)	0.14	(0.09)	0.18	(0.00)	0	---
2016	0.29	(0.19)	0.26	(0.17)	0.24	(0.20)	0.24	(0.20)	0.14	(0.09)	0.18	(0.00)	0	---
2017	0.29	(0.19)	0.26	(0.16)	0.24	(0.20)	0.24	(0.20)	0.14	(0.08)	0.18	(0.00)	0	---

Catch (t)	F_{OFL}		$Max F_{ABC}$		Author's recommended F		Author's recommended F with $F_{50\%}$ in 2005 & 2006		50% of max FABC		Average F		$F = 0$	
	2005	144,340	(0.00)	124,800	(0.00)	106,530	(0.00)	78,980	(0.00)	65,540	(0.00)	86,880	(0.00)	0
2006	103,250	(0.04)	95,250	(0.04)	86,280	(0.04)	70,010	(0.04)	60,580	(0.04)	77,850	(0.04)	0	---
2007	72,250	(0.17)	68,880	(0.16)	64,440	(0.15)	75,060	(0.14)	49,670	(0.13)	70,010	(0.13)	0	---
2008	72,880	(0.57)	68,780	(0.53)	64,370	(0.54)	70,440	(0.51)	49,420	(0.41)	71,140	(0.30)	0	---
2009	96,080	(0.76)	88,270	(0.73)	82,620	(0.77)	85,770	(0.74)	58,400	(0.60)	78,600	(0.47)	0	---
2010	126,390	(0.69)	114,950	(0.67)	109,310	(0.72)	110,640	(0.71)	72,770	(0.60)	92,030	(0.50)	0	---
2011	149,040	(0.60)	136,290	(0.59)	131,380	(0.63)	131,800	(0.63)	86,650	(0.54)	105,510	(0.47)	0	---
2012	160,940	(0.55)	148,630	(0.54)	144,290	(0.58)	144,310	(0.57)	96,730	(0.49)	114,920	(0.45)	0	---
2013	165,950	(0.54)	154,510	(0.52)	150,470	(0.55)	150,380	(0.55)	102,810	(0.47)	120,650	(0.43)	0	---
2014	166,760	(0.54)	156,190	(0.52)	151,760	(0.55)	151,690	(0.55)	105,030	(0.45)	122,510	(0.43)	0	---
2015	168,390	(0.55)	157,620	(0.52)	153,580	(0.55)	153,550	(0.55)	106,890	(0.45)	124,260	(0.44)	0	---
2016	168,450	(0.54)	158,020	(0.52)	153,890	(0.55)	153,870	(0.55)	107,920	(0.45)	125,300	(0.43)	0	---
2017	168,530	(0.54)	158,250	(0.51)	154,110	(0.55)	154,100	(0.55)	108,620	(0.45)	126,010	(0.43)	0	---




Figure 1.1 Pollock catch in 2003 by 10 sq. nmi. blocks by season in the Gulf of Alaska as determined by observer-recorded haul retrieval locations. Blocks with less than 1.0 t of pollock catch are not shown. The size of the circle is proportional to the catch.




Figure 1.2. Incidental catch (including both retained and discarded species) in tows classified as pollock targets in the Gulf of Alaska (1997-2002).




Figure 1.3. 2003 catch age composition by half year and statistical area.




Figure 1.4. Gulf of Alaska pollock catch age composition (1976-2003). The diameter of the circle is proportional to the catch. Diagonal lines show strong year classes (1972, 1975, 1976, 1977, 1978, 1979, 1984, 1988, 1994, and 1999).





Figure 1.5. Biomass estimates of juvenile pollock (top) and adult pollock (bottom) from 1986-2004 Shelikof Strait EIT surveys. Bottom panel also shows the model estimate of total spawning biomass.




Figure 1.6. Biomass by length for pollock in the Shelikof Strait EIT survey (1981-2004, except 1982,1987 and 1999).




Figure 1.7. Length frequency of pollock in the ADF&G crab/groundfish trawl survey (1989-2004, except 1991 and 1995).





Figure 1.8. Trends in pollock biomass since 1986 for the Shelikof Strait EIT survey, the triennial bottom trawl survey, and the ADF&G crab/groundfish survey (top). Each survey biomass estimate is standardized to the survey average since 1986. The Shelikof Strait EIT survey is split into separate time series corresponding to the two acoustic systems used for the survey. In the bottom panel, a lowess smooth (SPLUS 1993) of the same data is compared to the estimated biomass trend from the assessment model.




Figure 1.9. Gulf of Alaska pollock catch characteristics.




Figure 1.10. Estimates of the proportion mature at age from visual maturity data collected during 2001-2004 winter EIT surveys in the Gulf of Alaska.





Figure 1.11. Age at 50% mature (top) and length at 50% mature (bottom) from annual logistic regressions for female pollock from winter EIT survey data in the Gulf of Alaska, 1983-2004.





Figure 1.12. Comparison of estimated female spawning biomass for alternative models. The top panel shows the entire modeled period, while the bottom panel shows the percent difference relative to the base model (fixed NMFS survey catchability) since 1990.




Figure 1.13. Observed and predicted fishery age composition for Gulf of Alaska pollock based on Model 3. Continuous lines are model predictions and lines with + symbol are observed proportions at age.





Figure 1.14. Residuals from Model 3 for fishery age composition (1972-2003). Circle diameters are proportional to the magnitude of the residual. Circles drawn with dotted lines indicate negative residuals. Diagonal lines show the strong year classes (1972, 1975, 1976, 1977, 1978, 1984, 1988, 1994, and 1999).

Shelikof Strait EIT survey



NMFS bottom trawl survey




Figure 1.15. Residuals from Model 3 for the Shelikof Strait EIT survey age composition (top) and NMFS bottom trawl age composition (bottom). Circle diameters are proportional to the magnitude of the residual. Circles drawn with dotted lines indicate negative residuals. Diagonal lines show the strong year classes (1972, 1975, 1976, 1977, 1978, 1979, 1984, 1988, 1994, and 1999).




Figure 1.16. Residuals from Model 3 for the ADF&G survey length composition.





Figure 1.17. Model predicted and observed survey biomass for the Shelikof Strait EIT survey. The Shelikof EIT survey is modeled with two catchability periods corresponding to the two acoustic systems used for the survey. Error bars indicate plus and minus two standard deviations. Since variance estimates are unavailable for EK500 biomass estimates, an assumed CV of 0.2 is used in the assessment model.





Figure 1.18. Model predicted and observed survey biomass for the NMFS bottom trawl survey (top), and the ADFG crab/groundfish survey (bottom). Error bars indicate plus and minus two standard deviations. Since variance estimates are unavailable for ADFG biomass estimates, an assumed CV of 0.25 is used in the assessment model.





Figure 1.19. Model predicted and observed survey biomass for the historical 400-mesh eastern trawl surveys (top), and the egg production survey (bottom). Error bars indicate plus and minus two standard deviations.




Figure 1.20. Uncertainty in the catchability coefficient for the NMFS trawl survey from a likelihood profile for Model 1.




Figure 1.21. Estimates of time-varying fishery selectivity for Gulf of Alaska pollock.





Figure 1.22. Estimated time series of Gulf of Alaska pollock spawning biomass (million t, top) and age-2 recruitment (billions of fish, bottom) from 1961 to 2005. Vertical bars represent two standard deviations. The B35% and B40% lines represent the current estimate of these benchmarks.





Figure 1.23. Retrospective plot of estimated Gulf of Alaska pollock female spawning biomass for stock assessments in the years 1994-2004 (top). For this figure, the time series of female spawning biomass for the 2004 assessment was calculated using the weight and maturity at age used in previous assessments to facilitate comparison. The bottom panel shows the estimated age composition in 2004 from the 2003 and 2004 assessments.




Figure 1.24. Gulf of Alaska pollock spawner productivity $\log(R / S)$ in 1961-2002 (top). A five-year running average is also shown. Spawner productivity in relation to female spawning biomass (bottom). The Ricker stock-recruit curve is linear in a plot of spawner productivity against spawning biomass. Horizontal lines indicate the mean spawner productivity for each decade within the range of spawning biomass indicated by the endpoints of the lines.




Figure 1.25. Gulf of Alaska pollock spawning biomass relative to the unfished level and fishing mortality relative to F_{OFL} (1961-2004). The ratio of fishing mortality to F_{OFL} is calculated using the estimated selectivity pattern in that year. Estimates of unfished spawning biomass are based on current estimates of maturity at age, weight at age, and mean recruitment. Because these estimates change as new data become available, this figure can only be used in a general way to evaluate management performance relative to biomass and fishing mortality reference levels.





Figure 1.26. Uncertainty in the estimate of recruitment abundance of the 1999 year class in 2001, 2002, 2003, and 2004 stock assessments (top) and the 2000 year class in 2002, 2003, and 2004 assessments (bottom).




Figure 1.27. Uncertainty in spawning biomass in 2005-2008 based on a thinned MCMC chain from the joint marginal likelihood for Model 3 where catch in 2005 is set to the author's recommended ABC. In 2006-2008, catch is based on the author's recommended fishing mortality schedule.





Figure 1.28. Projected spawning biomass and catches in 2005-10 under different harvest strategies.




Figure 1.29. Gulf of Alaska food web showing demersal (red) and pelagic (blue) pathways. Walleye pollock is shown in green. Pollock consumers stain green according to the importance of pollock in their diet.





Figure 1.30. Diet of walleye pollock juveniles (top) and adults (bottom) from an ECOPATH model of the Gulf of Alaska. Pollock less than 20cm are considered juveniles.





Figure 1.31. Sources of mortality for walleye pollock juveniles (top) and adults (bottom) from an ECOPATH model of the Gulf of Alaska. Pollock less than 20cm are considered juveniles.




Figure 1.32. Diet diversity of major predators of walleye pollock from an ECOPEATH model for Gulf of Alaska during 1990-94.




Figure 1.33. Trends in the biomass of major predators of walleye pollock in the Gulf of Alaska (top). Trends in annual consumption of pollock (kg consumed per kg predator biomass) by major predators. Estimates based on diet composition from stomach samples collected during NMFS bottom trawl surveys (1990-2001).




Figure 1.34. Length at age of female arrowtooth flounder in Gulf of Alaska 1975-2003.




Figure 1.36. Pair-wise Spearman rank correlation between abundance trends of walleye pollock, pollock fishery catches, Steller sea lions, arrowtooth flounder, Pacific halibut, and Pacific cod in the Gulf of Alaska. Rank correlations are based on the years in which abundance estimates are available for each pair.

Appendix A: Southeast Alaska pollock

Bottom trawl surveys indicate a substantial reduction in pollock abundance east of 140° W. lon. Stock structure in this area is poorly understood. Bailey et al. (1999) suggest that pollock metapopulation structure in southeast Alaska is characterized by numerous fiord populations. In the 1996, 1999 and 2003 bottom trawl surveys, higher pollock CPUE in southeast Alaska occurred primarily from Cape Ommaney to Dixon Entrance, where the shelf is more extensive. Pollock size composition in the 1996, 1999 and 2003 surveys was dominated by smaller fish (<40 cm). These juveniles are unlikely to influence the population dynamics of pollock in the central and western Gulf of Alaska. Ocean currents are generally northward in this area, suggesting that juvenile settlement is a result of spawning further south. Spawning aggregations of pollock have been reported from the northern part of Dixon Entrance (Saunders et al. 1988).

Historically, there has been little directed fishing for pollock in southeast Alaska (Fritz 1993). During 1991-2002, pollock catch the Southeast and East Yakutat statistical areas averaged 17 t (Table 1.2). The current ban on trawling east of 140° W. lon. prevents the development of a trawl fishery for pollock in Southeast Alaska.

Pollock biomass estimates from the bottom trawl survey are highly variable, in part due to year-to-year differences in survey coverage. Surveys in 1996, 1999, and 2003 had the most complete coverage of shallow strata in southeast Alaska, and indicate that stock size is approximately 25-75,000 t (Fig. 1.37). We recommend placing southeast Alaska pollock in Tier 5 of NPFMC harvest policy, and basing the ABC and OFL on natural mortality (0.3) and the biomass for the 2003 survey. Biomass in southeast Alaska was estimated by splitting survey strata and CPUE data in the Yakutat INPFC area at 140° W. lon. and combining the strata east of the line with comparable strata in the Southeastern INPFC area. This gives a **2005 ABC of 6,520 t** ($28,979 \text{ t} * 0.75 \text{ M}$), and a **2005 OFL of 8,694 t** ($28,979 \text{ t} * \text{M}$).




Figure 1.37. Pollock age composition in 2003 (left) and biomass trend in southeast Alaska from NMFS bottom trawl surveys in 1990-2003 (right).

Appendix B: Gulf pollock stock assessment model

Population dynamics

The age-structured model for pollock describes the relationships between population numbers by age and year. The modeled population includes individuals from age 2 to age 10, with age 10 defined as a “plus” group, i.e., all individuals age 10 and older. The model extends from 1961 to 2004 (44 yrs). The Baranov (1918) catch equations are assumed, so that

$$c_{ij} = N_{ij} \frac{F_{ij}}{Z_{ij}} [1 - \exp(-Z_{ij})]$$

$$N_{i+1,j+1} = N_{ij} \exp(-Z_{ij})$$

$$Z_{ij} = \sum_k F_{ik} + M$$

except for the plus group, where

$$N_{i+1,10} = N_{i,9} \exp(-Z_{i,9}) + N_{i,10} \exp(-Z_{i,10})$$

where N_{ij} is the population abundance at the start of year i for age j fish, F_{ij} = fishing mortality rate in year i for age j fish, and c_{ij} = catch in year i for age j fish. A constant natural mortality rate, M , irrespective of year and age, is assumed.

Fishing mortality is modeled as a product of year-specific and age-specific factors (Doubleday 1976)

$$F_{ij} = s_j f_i$$

where s_j is age-specific selectivity, and f_i is the annual fishing mortality rate. To ensure that the selectivities are well determined, we require that $\max(s_j) = 1$. Following previous assessments, a scaled double-logistic function (Dorn and Methot 1990) was used to model age-specific selectivity,

$$s'_j = \left(\frac{1}{1 + \exp[-\beta_1(j - \alpha_1)]} \right) \left(1 - \frac{1}{1 + \exp[-\beta_2(j - \alpha_2)]} \right)$$

$$s_j = s'_j / \max (s'_j)$$

where α_1 = inflection age, β_1 = slope at the inflection age for the ascending logistic part of the equation, and α_2 , β_2 = the inflection age and slope for the descending logistic part.

Measurement error

Model parameters were estimated by maximum likelihood (Fournier and Archibald 1982, Kimura 1989, 1990, 1991). Fishery observations consist of the total annual catch in tons, C_i , and the proportions at age in the catch, p_{ij} . Predicted values from the model are obtained from

$$\hat{C}_i = \sum_j w_{ij} c_{ij}$$

$$\hat{p}_{ij} = c_{ij} / \sum_j c_{ij}$$

where w_{ij} is the weight at age j in year i . Year-specific weights at age are used when available.

Log-normal measurement error in total catch and multinomial sampling error in the proportions at age give a log-likelihood of

$$\log L_k = \sum_i [\log (C_i) - \log (\hat{C}_i)]^2 / 2 \sigma_i^2 + \sum_i m_i \sum_j p_{ij} \log (\hat{p}_{ij} / p_{ij})$$

where σ_i is standard deviation of the logarithm of total catch ($\sim CV$ of total catch) and m_i is the size of the age sample. In the multinomial part of the likelihood, the expected proportions at age have been divided by the observed proportion at age, so that a perfect fit to the data for a year gives a log likelihood value of zero (Fournier and Archibald 1982). This formulation of the likelihood allows considerable flexibility to give different weights (i.e. emphasis) to each estimate of annual catch and age composition. Expressing these weights explicitly as CVs (for the total catch estimates), and sample sizes (for the proportions at age) assists in making reasonable assumptions about appropriate weights for estimates whose variances are not routinely calculated.

Survey observations consist of a total biomass estimate, B_i , and survey proportions at age π_{ij} . Predicted values from the model are obtained from

$$\hat{B}_i = q \sum_j w_{ij} s_j N_{ij} \exp [\phi_i Z_{ij}]$$

where q = survey catchability, w_{ij} is the survey weight at age j in year i (if available), s_j = selectivity at age for the survey, and ϕ_i = fraction of the year to the mid-point of the survey. Although there are multiple surveys for Gulf pollock, a subscript to index a particular survey has been suppressed in the above and subsequent equations in the interest of clarity. Survey selectivity was modeled using either a double-logistic function of the same form used for fishery selectivity, or simpler variant, such as single logistic function. The expected proportions at age in the survey in the i th year are given by

$$\hat{\pi}_{ij} = s_j N_{ij} \exp[\phi_i Z_{ij}] / \sum_j s_j N_{ij} \exp[\phi_i Z_{ij}]$$

Log-normal errors in total biomass and multinomial sampling error in the proportions at age give a log-likelihood for survey k of

$$\log L_k = \sum_i [\log(B_i) - \log(\hat{B}_i)]^2 / 2 \sigma_i^2 + \sum_i m_i \sum_j \pi_{ij} \log(\hat{\pi}_{ij} / \pi_{ij})$$

where σ_i is the standard deviation of the logarithm of total biomass (\sim CV of the total biomass) and m_i is the size of the age sample from the survey.

Process error

Process error refers to random changes in parameter values from one year to the next. Annual variation in recruitment and fishing mortality can be considered types of process error (Schnute and Richards 1995). In the pollock model, these annual recruitment and fishing mortality parameters are generally estimated as free parameters, with no additional error constraints. We use process error to describe changes in fisheries selectivity over time. To model temporal variation in a parameter γ , the year-specific value of the parameter is given by

$$\gamma_i = \bar{\gamma} + \delta_i$$

where $\bar{\gamma}$ is the mean value (on either a log scale or linear scale), and δ_i is an annual deviation subject to the constraint $\sum \delta_i = 0$. For a random walk where annual *changes* are normally distributed, the log-likelihood is

$$\log L_{Proc.Err.} = \sum \frac{(\delta_i - \delta_{i+1})^2}{2 \sigma_i^2}$$

where σ_i is the standard deviation of the annual change in the parameter. We use a process error model for all four parameters of the fishery double-logistic curve.

The total log likelihood is the sum of the likelihood components for each fishery and survey, plus a term for process error,

$$\text{Log } L = \sum_k \text{Log } L_k + \sum_p \text{Log } L_{Proc.Err.}$$

Appendix C: Seasonal distribution and apportionment of walleye pollock among management areas in the Gulf of Alaska

Since 1992, the Gulf of Alaska pollock TAC has been apportioned between management areas based on the distribution of biomass in groundfish surveys. Both single species and ecosystem considerations provide the rationale for apportioning the TAC. From an ecosystem perspective, apportioning the TAC will spatially distribute the effects of fishing on other pollock consumers (i.e., Steller sea lions), potentially reducing the overall intensity of any adverse effects. From the perspective of the pollock population, apportioning the TAC ensures that no smaller component of the stock experiences higher mortality than any other. Although no sub-stock units of pollock have yet been identified in the Gulf of Alaska, it would be precautionary to manage the fishery so that if these sub-units do exist they would not be subject to high fishing mortality. Protection of sub-stock units would be most important during spawning season, when they are spatially separated. The Steller Sea Lion Protection Measures implemented in 2001 require apportionment of pollock TAC based on the seasonal distribution of biomass. Although spatial apportionment is intended to reduce the potential impact of fishing on endangered Steller Sea Lions, it is important to recognize that apportioning the TAC based on an inaccurate or inappropriate estimate of biomass distribution could be detrimental, both to pollock population itself, and on species that depend on pollock.

Walleye pollock in the Gulf of Alaska undergo an annual migration between summer foraging habitats and winter spawning grounds. Since surveying effort has been concentrated during the summer months and prior to spawning in late winter, the dynamics and timing of this migration are not well understood. Regional biomass estimates are highly variable, indicating either large sampling variability, large interannual changes in distribution, or, more likely, both. There is a comprehensive survey of the Gulf of Alaska in summer, but historically surveying during winter has focused on the Shelikof Strait spawning grounds. Recently there has been expanded EIT surveying effort outside of Shelikof Strait in winter, but no acoustic survey has been comprehensive, covering all areas where pollock could potentially occur.

Winter distribution

In winter, an annual acoustic survey in Shelikof Strait has been conducted since 1981. A significant portion of the remaining shelf and upper slope waters in the Gulf of Alaska west of Cape Suckling has been surveyed at least once during winter by exploratory surveys and surveys with shorter time series. Therefore a “composite” approach was developed to use data from several different surveys. We used data from 1) Shelikof Strait surveys in 1992-2003, 2) surveys of the Shumagin Island area in 1995, and 2001-03 (Wilson et al. 1995, Guttormsen et al. 2001, 2002, 2003), and 3) an exploratory survey along the shelf break in 1990 (Karp 1990). Each of these surveys covered a non-overlapping portion of the Gulf of Alaska shelf and upper slope west of Cape Suckling. Surveys of the Shumagin Island area in 1994 and 1996 were not used in this analysis because most fish were in post-spawning condition, and replicated surveys of spawning pollock in Shelikof Strait indicate a rapid decline in abundance after peak spawning (Wilson 1994, Wilson et al. 1996).

The “composite” approach was to estimate the percent of the total stock surveyed during a particular survey by dividing the survey biomass by the estimated total biomass of pollock at spawning from the assessment model. The percent for each non-overlapping survey was added together to form a composite biomass distribution, which, with some luck, ought to be close to 100%. Model estimates of biomass at spawning took into account the total mortality between the start of the year and spawning, and used mean weight at age from Shelikof Strait surveys in 1992-2004.

Results indicate that an average of 65% of the pollock biomass was in Shelikof Strait in winter (Appendix table 1.1). For the Shumagin surveys in 1995, 2001, and 2002, 24% of the total stock biomass was surveyed on average. The sum of the percent biomass for all surveys was 97%, which may reflect sampling variability, interannual variation in spawning location, or differences in echo sounder/integration systems, but also suggests reasonable consistency between the aggregate biomass of pollock surveyed acoustically in winter and the assessment model estimates of abundance. After rescaling, the resulting average biomass distribution was 24.12%, 66.20%, and 9.68% in areas 610, 620, and 630. These estimates are within 1% of last year's estimates. We have not used recent survey results along the shelf break in areas 620 and 630, nor the 2003 estimates of biomass in Sanak Gully in area 610 because the relationship between these newly surveyed aggregations and those in Shelikof Strait and the Shumagin Islands is unclear.

A-season apportionment between areas 620 and 630

Last year, based on evaluation of fishing patterns which suggested that the migration to spawning areas was not complete by January 20, the plan team recommended an alternative apportionment scheme for areas 620 and 630 based on the midpoint of the summer and winter distributions in area 630. This approach was not used for area 610 because A season fishing patterns suggested that most of the fish captured in area 610 would eventually spawn in area 610. The resulting A season apportionment updated with 2004 survey data is: 610, 24.12%; 620, 56.01%; 630, 19.87%.

Middleton Island winter EIT survey results in 2003

The apportionment for area 640, which is not managed by season, has previously been based on the summer distribution of the biomass. Fishing, however, takes places primarily in winter or early spring on a spawning aggregation near Middleton Island. During 28-29 March 2003, this area was surveyed by the NOAA ship *Miller Freeman* for the first time and biomass estimate of 6900 t was obtained. Although maturity stage data suggested the timing of the survey was appropriate, discussions with fishing vessels contacted during the survey raised some questions about survey timing relative to peak biomass. Notwithstanding, a tier 5 calculation based on this spawning biomass gives an ABC of 1,550 t (6,901 t * 0.75 M), compared to 1,560 t for the author's 2005 ABC recommendation and an apportionment based on the summer biomass distribution. This suggests that the current approach of basing the area 640 apportionment on the gulfwide ABC and the summer biomass distribution is at least consistent with the biomass present near Middleton Island in the winter. We recommend continuing this approach until sufficient survey information during winter has accumulated to evaluate interannual variation in the biomass present in this area.

Summer distribution

The NMFS bottom trawl is summer survey (typically extending from mid-May to mid-August). Because of large shifts in the distribution of pollock between management areas one survey to the next, and the high variance of biomass estimates by management area, Dorn et al. (1999) recommended that the apportionment of pollock TAC be based upon the four most recent NMFS summer surveys. The four-survey average was updated with 2003 survey results in an average biomass distribution of 47.68%, 20.88%, 29.46%, and 1.98% in areas 610, 620, 630, and 640 (Fig. 1.38).

Example calculation of 2005 Seasonal and Area TAC Allowances for W/C/WYK

Warning: This example is based on hypothetical ABC of 100,000 t.

1) Deduct the Prince William Sound Guideline Harvest Level.

2) Use summer biomass distribution for the 640 allowance:

$$640 \quad 0.0198 \times \text{Total TAC} = 1,980 \text{ t}$$

3) Calculate seasonal apportionments of TAC for the A, B, C, and D seasons at 25 %, 25%, 25%, and 25 % of the remaining annual TAC west of 140° W lon.

$$\text{A season} \quad 0.25 \times (\text{Total TAC} - 1,980) = 24,505 \text{ t}$$

$$\text{B season} \quad 0.25 \times (\text{Total TAC} - 1,980) = 24,505 \text{ t}$$

$$\text{C season} \quad 0.25 \times (\text{Total TAC} - 1,980) = 24,505 \text{ t}$$

$$\text{D season} \quad 0.25 \times (\text{Total TAC} - 1,980) = 24,505 \text{ t}$$

4) For the A season, the allocation of TAC to areas 610, 620 and 630 is based on a blending of winter and summer distributions to reflect that pollock may not have completed their migration to spawning areas by Jan. 20, when the A season opens.

$$610 \quad 0.2412 * 24,505 \text{ t} = 5,911 \text{ t}$$

$$620 \quad 0.5601 * 24,505 \text{ t} = 13,725 \text{ t}$$

$$630 \quad 0.1987 * 24,505 \text{ t} = 4,869 \text{ t}$$

5) For the B season, the allocation of TAC to areas 610, 620 and 630 is based on the composite estimate of winter biomass distribution

$$610 \quad 0.2412 * 24,505 \text{ t} = 5,790 \text{ t}$$

$$620 \quad 0.6620 * 24,505 \text{ t} = 16,534 \text{ t}$$

$$630 \quad 0.0968 * 24,505 \text{ t} = 2,181 \text{ t}$$

6) For the C and D seasons, the allocation of remaining TAC to areas 610, 620 and 630 is based on the average biomass distribution in areas 610, 620 and 630 in the most recent four NMFS bottom trawl surveys.

$$610 \quad 0.4768 / (1 - 0.0198) \times 24,505 = 11,920 \text{ t}$$

$$620 \quad 0.2088 / (1 - 0.0198) \times 24,505 = 5,220 \text{ t}$$

$$630 \quad 0.2946 / (1 - 0.0198) \times 24,505 = 7,365 \text{ t}$$

$$610 \quad 0.4768 / (1 - 0.0198) \times 24,505 = 11,920 \text{ t}$$

$$620 \quad 0.2088 / (1 - 0.0198) \times 24,505 = 5,220 \text{ t}$$

$$630 \quad 0.2946 / (1 - 0.0198) \times 24,505 = 7,365 \text{ t}$$

Appendix Table 1.1. Estimates of winter pollock biomass distribution in management areas 610-630 from EIT surveys in the Gulf of Alaska.

Survey	Year	Model estimates		Survey biomass estimate ¹	Percent	Percent of biomass by management area			Percent of total biomass			
		of total 2+ biomass at spawning				Area 610	Area 620	Area 630	Area 610	Area 620	Area 630	
Shelikof Strait	1992	991,466		681,400	68.7%							
Shelikof Strait	1993	1,090,410		408,200	37.4%							
Shelikof Strait	1994	1,083,430		467,300	43.1%							
Shelikof Strait	1995	896,318		618,300	69.0%							
Shelikof Strait	1996	796,121		745,400	93.6%							
Shelikof Strait	1997	723,075		570,100	78.8%	0.0%	98.8%	1.2%				
Shelikof Strait	1998	606,500		489,900	80.8%	0.0%	97.5%	2.5%				
Shelikof Strait	2000	494,994		334,900	67.7%	0.0%	97.8%	2.2%				
Shelikof Strait	2001	497,560		369,600	74.3%	0.0%	98.3%	1.7%				
Shelikof Strait	2002	514,059		229,104	44.6%	0.0%	97.7%	2.3%				
Shelikof Strait	2003	542,508		270,200	49.8%	0.0%	97.6%	2.4%				
Shelikof Strait	2004	588,504		291,215	49.5%	0.0%	97.6%	2.4%				
Shelikof Strait	Average				63.1%	0.0%	97.9%	2.1%	0.0%	61.8%	1.3%	
Shumagin	1995	896,318		290,100	32.4%	90.0%	10.0%	0.0%				
Shumagin	2001	497,560		108,791	21.9%	84.8%	15.2%	0.0%				
Shumagin	2002	514,059		135,644	26.4%	100.0%	0.0%	0.0%				
Shumagin	2003	542,508		67,160	12.4%	99.7%	0.3%	0.0%				
Shumagin	Average				23.2%	93.6%	6.4%	0.0%	21.8%	1.5%	0.0%	
Shelf break/east side Kodiak	1990	949,901		96,610	10.2%	14.9%	6.2%	78.9%	1.5%	0.6%	8.0%	
Total					96.53%				23.28%	63.9%	9.34%	
Rescaled total					100.00%				24.12%	66.2%	9.68%	

¹The biomass of age-1 pollock was not included in Shelikof Strait survey biomass in 1995 and 2000.




Figure 1.38. Percent distribution of Gulf of Alaska pollock biomass west of 140° W lon. in NMFS bottom trawl surveys in 1984-2003. The percent in West Yakutat in 1984, 1987, and 2001 was set equal to the mean percent in 1990-99.

(This page intentionally left blank)