

5. Assessment of the Deepwater Flatfish Stock Complex in the Gulf of Alaska

Carey R. McGilliard and Wayne Palsson

November 2018

Executive Summary

Introduction

The Gulf of Alaska deepwater flatfish complex (consisting of Dover sole, Greenland turbot, and deepsea sole) is assessed every four years and was last assessed in 2015. In years without an assessment, we present an executive summary to recommend harvest levels for the next two years. Please refer to the 2015 full stock assessment report for further information regarding the assessment model (McGilliard and Palsson, 2015, available online at <http://www.afsc.noaa.gov/REFM/Docs/2015/GOAdeepflat.pdf>). A full stock assessment document with updated assessment and projection model results will be presented in 2019.

Dover sole is assessed using an age-structured model and Tier 3 determination. Thus, the single species projection model was run using parameter values from the accepted 2015 Dover sole assessment model (McGilliard and Palsson 2015), together with updated catch information for 2017-2020, to predict stock status for Dover sole in 2019 and 2020 and to make ABC recommendations for those years. Projections are conducted using numbers-at-age for Dover sole from age 3-59+ and historical recruitment of age 3 individuals to calculate OFL's and ABC's. Greenland turbot and deepsea sole fall under Tier 6. ABC's and OFL's for Tier 6 species are based on historical catch levels (average catch over the years 1978-1995) and therefore these quantities cannot be updated. ABC's and OFL's for the individual species in the deepwater flatfish complex are determined only as an intermediate step for the purpose of calculating complex-level OFL's and ABC's.

Summary of Results

As in previous years (McGilliard 2017), the species-level ABC is 179 t for Greenland turbot and the OFL is 238 t for both 2019 and 2020. The species-level ABC for deepsea sole is 4 t and the OFL is 6 t for both 2019 and 2020. The species-level ABC for Dover sole is 9,318 t in 2019 and 9,441 t in 2020 and the OFL is 11,190 t in 2019 and 11,337 t in 2020.

Based on the updated projection model results, the recommended complex-level ABC's for 2019 and 2020 are 9,501 t and 9,624 t, and the OFL's are 11,434 t and 11,581 t. The new ABC recommendation and OFL for 2019 are similar to those developed in 2018 (9,499 t and 11,431 t). The principal reference values are shown in the following table:

Species	Quantity	As estimated or <i>specified last year for:</i>		As estimated or <i>recommended this year for:</i>	
		2018	2019	2019*	2020*
Dover sole	<i>M</i> (natural mortality rate)	0.085	0.085	0.085	0.085
	Tier	3a	3a	3a	3a
	Projected total (3+) biomass (t)	144,654	145,899	145,926	147,001
	Projected Female spawning biomass (t)	49,366	49,373	49,385	49,418
	<i>B</i> _{100%}	57,871	57,871	57,871	57,871
	<i>B</i> _{40%}	23,148	23,148	23,148	23,148
	<i>B</i> _{35%}	20,255	20,255	20,255	20,255
	<i>F</i> _{OFL}	0.12	0.12	0.12	0.12
	<i>maxF</i> _{ABC}	0.1	0.1	0.1	0.1
	<i>F</i> _{ABC}	0.1	0.1	0.1	0.1
	OFL (t)	11,050	11,187	11,190	11,337
maxABC (t)	9,202	9,316	9,318	9,441	
ABC (t)	9,202	9,316	9,318	9,441	
Greenland turbot	Tier	6	6	6	6
	OFL (t)	238	238	238	238
	maxABC (t)	179	179	179	179
	ABC (t)	179	179	179	179
Deepsea sole	Tier	6	6	6	6
	OFL (t)	6	6	6	6
	maxABC (t)	4	4	4	4
	ABC (t)	4	4	4	4
Deepwater Flatfish Complex	OFL (t)	11,294	11,431	11,434	11,581
	maxABC (t)	9,385	9,499	9,501	9,624
	ABC (t)	9,385	9,499	9,501	9,624
	Status	As determined <i>last</i> year for:		As determined <i>this</i> year for:	
		2016	2017	2017	2018
	Overfishing	no	n/a	no	n/a
	Overfished	n/a	no	n/a	no
Approaching overfished	n/a	no	n/a	no	

*Projections are based on estimated catches of 209 t used in place of maximum permissible ABC for 2018 and 270 t used in place of maximum permissible ABC for 2019-2020. The 2018 projected catch was calculated as the current catch as of October 6, 2018 added to the average October 6 – December 31 catches over the 5 previous years. The 2019-2020 projected catch was calculated as the average catch over the previous 5 years.

Area Apportionment

Area apportionment for ABC of deepwater flatfish is currently based on the proportion of survey biomass of Greenland Turbot and deepsea sole found within each management area from 2001-2017 and an estimates of 2018 and 2019 survey biomass for Dover sole in each management area based on results from the random effects model. An ABC exists only at the level of the complex (deepwater flatfish) and

not for each species individually. The ABC by area for the deepwater flatfish complex is then the sum of the species-specific portions of the ABC.

The random effects model is used to fill in depth and area gaps in the Dover sole survey biomass by area and to calculate an area- and depth-specific estimate of 2019 and 2020 survey biomass. These estimates are summed over depths and the resulting relative biomass in each management area is used as the basis for apportionment of the Dover sole portion of the deepwater complex. This method of conducting area apportionment for deepwater flatfish was recommended by the GOA Plan Team in 2016 (McGilliard 2016). The method was chosen because it accounts for time and area gaps in the survey for Dover sole, which comprises nearly all of the deepwater flatfish catch and moves to deeper waters ontogenetically, and explicitly accounts for differences in the spatial distributions of Dover sole and Greenland turbot. Greenland turbot were found exclusively in the Western region by the survey over the period 2001-2017.

Species	Year	West				Total
		Western	Central	Yakutat	Southeast	
		2.5%	36.9%	35.2%	25.3%	100.0%
Dover Sole	2019	237	3,440	3,279	2,362	9,318
	2020	241	3,485	3,322	2,393	9,441
		100.0%	0.0%	0.0%	0.0%	100.0%
Greenland Turbot	2019	179	0	0	0	179
	2020	179	0	0	0	179
		0.7%	72.9%	15.3%	11.0%	100.0%
Deepsea Sole	2019	0	3	1	0	4
	2020	0	3	1	0	4
Deepwater Flatfish	2019	416	3,443	3,280	2,362	9,501
	2020	420	3,488	3,323	2,393	9,624

Figures




Figure 1. Catch:female spawning biomass ratio using spawning biomass for Dover sole only.

Tables

Table 1. Total catch of species in the deepwater flatfish complex. 2018 catch is current as of Oct. 8, 2018.

Year	Dover Sole	Deepsea Sole	Greenland Turbot	Total
1978	827	5	51	883
1979	530	5	24	559
1980	570	2	57	629
1981	457	8	8	473
1982	457	31	23	511
1983	354	11	145	510
1984	132	1	18	151
1985	43	3	0	46
1986	23	0	0	23
1987	56	0	44	100
1988	1,087	0	256	1,343
1989	1,521	0	56	1,577
1990	2,348	30	0	2,378
1991	10,067	1	127	10,196
1992	8,266	3	226	8,495
1993	6,675	6	24	6,706
1994	3,040	3	34	3,077
1995	1,865	1	345	2,211
1996	2,177	0	13	2,191
1997	3,642	1	16	3,659
1998	2,210	37	39	2,286
1999	2,261	0	22	2,283
2000	964	1	16	981
2001	801	0	2	804
2002	550	0	8	559
2003	934	0	17	951
2004	685	1	1	686
2005	413	0	5	418
2006	364	4	38	406
2007	281	0	0	281
2008	570	0	3	573
2009	466	7	3	476
2010	545	0	0	545
2011	465	0	1	466
2012	261	0	0	262
2013	240	1	1	242
2014	340	1	14	355
2015	251	1	4	256
2016	234	2	2	238
2017	253	1	5	259
2018	169	1	0	171

Table 2. Dover sole survey biomass by area and depth. Depth is reported as maximum depth in meters (e.g. "200" indicates depths of 101-200 m).

Year	Depth	Western	Central	Eastern	Total	Year	Depth	Western	Central	Eastern	Total
1984		4,460	52,469	11,592	68,521	2005		2,832	38,881	38,847	80,560
	100	34	1,870	925	2,829		100	475	4,255	1,924	6,654
	200	725	24,506	4,989	30,220		200	468	19,805	12,340	32,613
	300	355	5,598	1,975	7,928		300	275	6,691	10,732	17,697
	500	1,138	4,039	1,645	6,822		500	455	4,742	12,577	17,774
	700	1,290	5,147	1,728	8,166		700	312	1,617	1,206	3,134
	1000	919	11,309	330	12,557		1000	848	1,772	69	2,689
1987		2,623	34,577	26,194	63,394	2007		2,325	43,404	25,740	71,469
	100	5	1,260	3,137	4,401		100	78	1,748	903	2,728
	200	108	12,728	12,995	25,831		200	405	22,417	6,887	29,709
	300	32	8,587	3,419	12,039		300	110	9,543	9,945	19,598
	500	1,103	3,706	4,126	8,934		500	468	4,437	6,430	11,335
	700	1,267	6,757	2,518	10,542		700	208	3,604	1,298	5,109
	1000	108	1,539		1,647		1000	1,056	1,655	278	2,989
1990		1,649	71,109	23,839	96,597	2009		5,067	35,820	35,389	76,277
	100	161	11,233	896	12,290		100	154	2,372	4,008	6,534
	200	716	42,188	14,869	57,774		200	565	15,668	10,253	26,486
	300	50	15,644	4,290	19,985		300	88	12,619	10,979	23,685
	500	721	2,043	3,784	6,549		500	548	3,158	5,595	9,300
1993		2,379	43,515	39,664	85,557		700	3,712	1,769	4,144	9,625
	100	180	3,937	651	4,768		1000	0	236	411	646
	200	1,044	24,054	18,901	43,999	2011		833	35,548	41,150	77,531
	300	154	10,883	8,893	19,930		100	235	1,810	2,377	4,422
	500	1,001	4,640	11,219	16,861		200	146	14,528	10,065	24,739
1996		1,458	37,144	40,928	79,531		300	8	15,131	11,102	26,241
	100	134	1,674	4,753	6,561		500	134	2,578	16,704	19,416
	200	337	21,452	16,066	37,856		700	311	1,501	902	2,714
	300	290	8,691	9,121	18,101	2013		979	23,180	58,580	82,739
	500	698	5,327	10,988	17,013		100	0	1,196	23,355	24,551
1999		1,442	34,155	38,648	74,245		200	627	7,789	7,928	16,344
	100	7	3,619	2,806	6,431		300	126	9,896	11,178	21,201
	200	56	14,068	14,425	28,549		500	84	2,026	14,994	17,104
	300	43	8,085	11,448	19,576		700	142	2,273	1,125	3,540
	500	651	4,779	6,887	12,317	2015		336	20,067	32,667	53,069
	700	685	2,889	2,476	6,049		100	0	730	2,094	2,824
	1000	0	716	606	1,323		200	85	7,284	10,225	17,594
2001		895	31,529		32,424		300	34	6,044	5,254	11,332
	100	18	3,785		3,803		500	157	2,885	12,796	15,838
	200	53	16,241		16,294		700	60	1,222	2,256	3,538
	300	188	7,303		7,491		1000	0	1,901	42	1,943
	500	636	4,200		4,836	2017		260	20,495	37,552	58,307
2003		3,149	49,283	46,865	99,297		100	37	170	678	885
	100	194	2,842	7,119	10,154		200	134	7,753	20,583	28,470
	200	541	23,005	21,636	45,181		300	62	10,143	5,475	15,680
	300	270	10,070	7,491	17,832		500	27	1,663	10,398	12,089
	500	811	4,629	8,153	13,593		700	0	765	419	1,184
	700	1,333	8,738	2,466	12,537						

Literature Cited

- McGilliard, C.R. and Palsson, W. 2017. 5. Gulf of Alaska Deepwater Flatfish. In Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Gulf of Alaska. pp. 649-656. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage AK 99510.
- McGilliard, C.R. and Palsson, W. 2015. 5. Gulf of Alaska Deepwater Flatfish. In Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Gulf of Alaska. pp. 563-624. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage AK 99510.