

## Chapter 6.5 Kamchatka Flounder

By

Thomas K. Wilderbuer, Daniel G. Nichol and Robert Lauth

Alaska Fisheries Science Center  
 NMFS/NOAA 7600 Sand Point Way NE  
 Seattle WA 98115

### Executive Summary

This document is the initial analysis of stock status and harvest recommendation for Kamchatka flounder as a single species. It has heretofore been a constituent of the *Atheresthes* species complex of which arrowtooth flounder had the dominant biomass and ABC's were set based upon its productivity and stock status. Due to the recent development of a targeted fishery on Kamchatka flounder it is no longer feasible to manage this species as part of the present complex due to the potential to overharvest the resource since the complex ABC exceeds the estimated biomass of Kamchatka flounder.

Quantity/Status	Last year		This year	
	2010	2011	2011	2012
<i>M</i> (natural mortality)	--	--	<b>0.2</b>	0.2
Specified/recommended Tier	3a	3a	<b>5</b>	5
Biomass	--	--	<b>128,800 t</b>	
$F_{OFL}$ (F=M)	--	--	<b>0.2</b>	
$maxF_{ABC}$ (maximum allowable = $0.75x F_{OFL}$ )	--	--	<b>0.15</b>	
Specified/recommended $F_{ABC}$	--	--	<b>0.15</b>	
Specified/recommended OFL (t)	--	--	<b>23,600 t</b>	
Specified/recommended ABC (t)	--	--	<b>17,700 t</b>	

## Introduction

The Kamchatka flounder (*Atheresthes evermanni*) is a relatively large flatfish which is distributed from Northern Japan through the Sea of Okhotsk to the Western Bering Sea north to Anadyr Gulf (Wilimovsky et al. 1967) and east to the eastern Bering Sea shelf. In U.S. waters they are found in the Aleutian Islands where they generally decrease in abundance from west to east (Zimmerman and Goddard 1996). They are also present in Bering Sea slope waters but are absent from the Alaska Peninsula and the Gulf of Alaska.

In the eastern part of their range, Kamchatka flounder overlap with arrowtooth flounder (*Atheresthes stomias*) which are very similar in appearance and were not routinely distinguished in the commercial catches until 2007. Until about 1992, these species were also not consistently separated in trawl survey catches (Fig. 6.5-1) and were combined in the arrowtooth flounder stock assessment (Wilderbuer et al. 2009). However, managing the two species as a complex became undesirable in 2010 due to the emergence of a directed fishery for Kamchatka flounder in the BSAI management area. Since the ABC was determined by the large amount of arrowtooth flounder relative to Kamchatka flounder (complex is about 93% arrowtooth flounder) the possibility arose of an overharvest of Kamchatka flounder as the *Atheresthes* sp. ABC exceeded the Kamchatka flounder biomass. Beginning with the 2011 fishing season, arrowtooth flounder and Kamchatka flounder are managed separately.

## Catch History

Historical Kamchatka flounder catch is combined in catch records of arrowtooth flounder and Greenland turbot from the 1960s. The fisheries for Greenland turbot intensified during the 1970s and the bycatch of arrowtooth flounder and Kamchatka flounder is assumed to have also increased. Catches of these species decreased after implementation of the MFCMA and the Kamchatka flounder resource has remained lightly exploited with the combined catches with arrowtooth flounder averaging 12,831 t from 1977-2008 (Table 6.5-1). It is estimated that only a small fraction (<10%) of this catch was Kamchatka flounder. This decline resulted from catch restrictions placed on the fishery for Greenland turbot and phasing out of the foreign fishery in the U.S. EEZ. Catches in Table 6.5-1 through 2006 are for arrowtooth flounder and Kamchatka flounder combined, catches thereafter are those estimated for Kamchatka flounder only. The total catch estimated for arrowtooth and Kamchatka flounder by the Alaska Regional Office is a blend of vessel reported catch and observer at-sea sampling of the catch which was not differentiated by species through 2010. However, observers have separately identified the two species from catches aboard trawl vessels since 2007 and their sampling has indicated that the proportion of Kamchatka flounder in the combined catch has steadily increased from 10% in 2007 to 55% in 2010.

year	Percent of combined catch
2007	10
2008	31
2009	45
2010	55

The increased harvest is the result of a recently developed market for Kamchatka flounder which has now become a fishery target. The 2010 estimated catch of Kamchatka flounder (through October 16) is 19,622 t, taken primarily in area 514 and to a lesser extent in area 518. The Kamchatka/arrowtooth flounder

combined catch by week in 2010 (Fig. 6.5-2) shows that targeting for Kamchatka flounder began May 1 and continued through September.

## **Data**

The data used in this assessment includes estimates of total catch and bottom trawl survey biomass estimates from the Bering Sea shelf, slope and Aleutian Islands surveys.

### **Absolute Abundance from Trawl Surveys**

Biomass estimates (t) for Kamchatka flounder from the standard shelf survey area in the eastern Bering Sea, slope surveys and the Aleutian Islands region are shown in Table 6.5-2. Reliable estimates of Kamchatka flounder became available in 1991 and they were estimated at an average biomass of 45,500 t through 1994 on the Bering Sea shelf (Fig. 6.5-1). During the following 11 years the biomass was estimated at a lower level (26,800 t average) before increasing to high and stable levels the past 5 years (56,000 t average). On the continental shelf they are usually found in highest concentrations at depths greater than 200 meters around the Pribilof Islands and also in the large shelf area west of St. Matthew Island. Trends of abundance from the slope and Aleutian Islands surveys also indicate an increasing resource. They are common in the deeper waters of the slope area (500 to 800 meters, Zimmerman and Goddard 1996) in both the Aleutian Islands and the eastern Bering Sea slope (Figs. 6.5-3 and 6.5-4).

An estimate of total BSAI biomass for the years in which Aleutian Islands and slope surveys were not conducted was calculated by averaging the years in closest temporal (before and after) proximity.

### **Length-weight, maximum age and natural mortality**

Length-weight measurements collected in 1999 from 193 fish indicate that males and females grow by accumulating the same weight for a given size (Fig. 6.5-5). Age at length calculations from a small sample collected in 1991 indicate that males and females exhibit divergent growth after about age 5-6 with females growing larger than males (Zimmerman and Goddard 1996). Both sexes have been found in relatively equal numbers and the oldest fish have been aged at 33 years indicating that Kamchatka flounder are similar in life history to other Bering Sea flatfish. Accordingly we tentatively set the natural mortality rate at 0.2 for both sexes for this assessment.

### **Acceptable Biological Catch and exploitation rate**

Kamchatka flounder have a wide-spread distribution along the deeper waters of the Bering Sea/Aleutian Islands region and are believed to be at a high level as discerned from the increases in survey estimates from the time-series of Bering Sea shelf, slope and Aleutian Islands surveys. The 2010 combined estimate of total biomass from the three surveys is 128,800 t. Exploitation rates estimated for 2008-2010 have steadily increased from 5% in 2008, 10% in 2009 to 15% in 2010.

Given the limited amount of biological information available for Kamchatka flounder, they are qualified to be managed under Tier 5 of Amendment 56 to the BSAI groundfish management plan, and thus have harvest recommendations which are directly calculated from estimates of biomass and natural mortality. The Tier 5 formula for calculating ABC is:  $ABC = 0.75 \times M \times \text{average biomass}$ .

ABC calculated from this formula is sensitive to the fluctuations in annual biomass estimated from bottom trawl surveys (shelf survey cv is 10%, Aleutians cv = 30%). In order to lessen this effect, annual estimates of Kamchatka flounder abundance (using trawl survey estimates when they are available and filling in missing years from the average of the closest previous and future year which bracket the missing

year) from the three surveys were summed and then ABC was calculated using running averages which ranged from 3 to the 7 most recent years (all with  $M = 0.2$ ). ABC estimates from these five methods indicate that the effect of annual variability on the estimate of ABC and OFL can be dampened by including more years in the estimation calculation which was particularly evident in the years of biomass increase from the past five years (Fig. 6.5-6 and Table 6.5-3). The seven year moving average is chosen for the ABC and OFL calculations for 2011 since it has the most resilience to the trawl survey variability and gives estimates which are close to the other moving averages.

The potential yield of Kamchatka flounder in 2011, based on a combined biomass of **128,800 t** from the combined trawl survey estimates is summarized as follows:

$F_{ABC}$	$F_{OFL}$	ABC	OFL
0.15	0.20	17,700	23,600

**The estimates of  $F_{abc}$  and  $F_{ofl}$  under tier 5 are  $0.75 \times M$  and  $M$ , respectively, and the ABC and OFL levels are the product of the fishing mortality rate and the 7 year running average of estimated biomass.**

## Ecosystem Considerations

### Predators of Kamchatka flounder

Kamchatka flounder have rarely been found in the stomachs of other groundfish species in samples collected by the Alaska Fisheries Science Center. Their presence has only been documented in 17 stomach samples from the BSAI where the predators included Pacific cod, pollock, Pacific halibut, arrowtooth flounder and two sculpin species.

### Kamchatka flounder predation

The prey of Kamchatka flounder can be discerned from 152 stomachs collected in 1983 (Yang and Livingston 1986). The principle diet was composed of walleye pollock, shrimp (most Crangonidae) and euphausiids. Pollock was the most important prey item for all sizes of fish, ranging from 56 to 86% of the total stomach content weight. An examination of diet overlap with arrowtooth flounder indicated that these two congeneric species basically consume the same resources.

### Ecosystem Effects on the stock

#### 1) Prey availability/abundance trends

Arrowtooth flounder diet varies by life stage as indicated in the previous section. Regarding juvenile prey and its associated habitat, information is not available to assess the abundance trends of the benthic infauna of the Bering Sea shelf. The original description of infaunal distribution and abundance by Haflinger (1981) resulted from sampling conducted in 1975 and 1976 and has not be re-sampled since. Information on pollock abundance is available in Chapter 1 of this SAFE report. It has been hypothesized that predators on pollock, such as adult arrowtooth flounder, may be important species which control (with other factors) the variation in year-class strength of juvenile pollock (Hunt et al. 2002). The populations of arrowtooth flounder which have occupied the outer shelf and slope areas of the Bering Sea

over the past twenty years for summertime feeding do not appear food-limited. These populations have fluctuated due to the variability in recruitment success which suggests that the primary infaunal food source has been at an adequate level to sustain the arrowtooth flounder resource.

## 2) Predator population trends

As juveniles, it is well-documented from studies in other parts of the world that flatfish are prey for shrimp species in near shore areas. This has not been reported for Bering Sea arrowtooth flounder due to a lack of juvenile sampling and collections in near shore areas, but is thought to occur. As late juveniles they are found in stomachs of pollock and Pacific cod, mostly on small arrowtooth flounder ranging from 5 to 15 cm standard length..

Past, present and projected future population trends of these predator species can be found in their respective SAFE chapters in this volume. Encounters between arrowtooth flounder and their predators may be limited as their distributions do not completely overlap in space and time.

## 3) Changes in habitat quality

Changes in the physical environment which may affect Kamchatka flounder distribution patterns, recruitment success, migration timing and patterns are catalogued in the Ecosystem Considerations Appendix of this SAFE report. Habitat quality may be enhanced during years and warmer bottom water temperatures with reduced ice cover (higher metabolism with more active feeding). Environmental factors important to juvenile survival are presently not well known.

---



---

**Ecosystem effects on Kamchatka flounder**


---

Indicator	Observation	Interpretation	Evaluation
<i>Prey availability or abundance trends</i>			
Benthic infauna	Stomach contents	Stable, data limited	Unknown
<i>Predator population trends</i>			
Fish (Pollock, Pacific cod)	Stable	Possible increases to Kamchatka mortality	
<i>Changes in habitat quality</i>			
Temperature regime	Cold years Kamchatka catchability and herding may decrease	Deeper water species so less likely to affect surveyed stock	No concern (dealt with in model)
Winter-spring environmental conditions	Affects pre-recruit survival	Probably a number of factors	Causes natural variability

---

**Arrowtooth flounder effects on ecosystem**


---

Indicator	Observation	Interpretation	Evaluation
<i>Fishery contribution to bycatch</i>			
Prohibited species	Stable, heavily monitored	Minor contribution to mortality	No concern
Forage (including Pollock, shrimp and euphausiids)	Stable, heavily monitored	Bycatch levels small relative to forage biomass	No concern
HAPC biota	Low bycatch levels of (spp)	Bycatch levels small relative to HAPC biota	No concern
Marine mammals and birds	Very minor direct-take	Safe	No concern
Sensitive non-target species	Likely minor impact		No concern
Data limited, likely to be safe			
<i>Fishery concentration in space and time</i>	Recent high exploitation rate	Little detrimental effect	No concern
<i>Fishery effects on amount of large size target fish</i>	Recent high exploitation rate, but unknown effect	Natural fluctuation	No concern
<i>Fishery contribution to discards and offal production</i>	Stable trend	Improving, but data limited	Possible concern
<i>Fishery effects on age-at-maturity and fecundity</i>	Unknown	NA	Possible concern

## References

- Wilderbuer, T. K., D. G. Nichol, and K. Aydin 2009. Arrowtooth flounder. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 2010, Chapter 6. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.
- Wilimovsky, N. J., A. Peden, and J. Peppar. 1967. Systematics of six demersal fishes of the north Pacific Ocean. Fish. Res. Board Can., Tech. Rep. 34, 52 p.
- Yang, M. S. and P. A. Livingston. 1986. Food habits and diet overlap of two congeneric species, *Atheresthes stomias* and *Atheresthes evermanni*, in the eastern Bering Sea. Fish. Bull. Vol. 82 (8)615-623.
- Zimmermann, Mark, and Pamela Goddard 1996. Biology and distribution of arrowtooth (*Atheresthes stomias*) and Kamachatka (*A. evermanni*) flounders in Alaskan waters. Fish. Bull 94:358-370.

Table 6.5-1. All nation total combined catch (t) of arrowtooth and Kamchatka flounder in the eastern Bering Sea and Aleutian Islands region<sup>a</sup>, 1970-2010. Catches since 1990 are not reported by area. Beginning in 2007, when the two species were differentiated in commercial catches, catch is reported for Kamchatka flounder only in this table

Year	Eastern Bering Sea				Aleutian Island Region				Total
	Non-U.S. fisheries <sup>b</sup>	U.S. J.V. <sup>c</sup>	U.S. DAH	Total	Non-U.S. fisheries	U.S. J.V.	U.S. DAH	Total	
1970	12,598			12,598	274			274	12,872
1971	18,792			18,792	581			581	19,373
1972	13,123			13,123	1,323			1,323	14,446
1973	9,217			9,217	3,705			3,705	12,922
1974	21,473			21,473	3,195			3,195	24,668
1975	20,832			20,832	784			784	21,616
1976	17,806			17,806	1,370			1,370	19,176
1977	9,454			9,454	2,035			2,035	11,489
1978	8,358			8,358	1,782			1,782	10,140
1979	7,921			7,921	6,436			6,436	14,357
1980	13,674	87		13,761	4,603			4,603	18,364
1981	13,468	5		13,473	3,624	16		3,640	17,113
1982	9,065	38		9,103	2,356	59		2,415	11,518
1983	10,180	36		10,216	3,700	53		3,753	13,969
1984	7,780	200		7,980	1,404	68		1,472	9,452
1985	6,840	448		7,288	11	59	89	159	7,447
1986	3,462	3,298	5	6,766		78	337	415	7,181
1987	2,789	1,561	158	4,508		114	237	351	4,859
1988		2,552	15,395	17,947		22	2,021	2,043	19,990
1989		2,264	4,000	6,264			1,042	1,042	7,306
1990		660	7,315	7,975			5,083	5,083	13,058
1991									22,052
1992									10,382
1993									9,338
1994									14,366
1995									9,280
1996									14,652
1997									10,054
1998									15,241
1999									10,573
2000									12,929
2001									13,908
2002									11,540
2003									12,834
2004									17,809
2005									13,685
2006									13,309
2007									1,183
2008									6,819
2009									12,802
2010 **									19,662

<sup>a</sup>Catches from data on file Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Seattle, WA 98115.

<sup>b</sup>Japan, U.S.S.R., Republic of Korea, Taiwan, Poland, and Federal Republic of Germany.

<sup>c</sup>Joint ventures between U.S. fishing vessels and foreign processing vessels.

\*\*Catch information through 16 October, 2010 (NMFS regional office).



Table 6.5-2 Estimated biomass from the three BSAI bottom trawl surveys.

Reliable estimates of Kamchatka flounder biomass are only available after 1991.

	<b>shelf</b>	<b>slope</b>	<b>Aleutian islands</b>
<b>1982</b>	0		
<b>1983</b>	17,299		1,034
<b>1984</b>	20,695		
<b>1985</b>	31		
<b>1986</b>	0		565
<b>1987</b>	40		
<b>1988</b>	13,723		
<b>1989</b>	17,108		
<b>1990</b>	32,799		
<b>1991</b>	37,152		16,255
<b>1992</b>	50,081		
<b>1993</b>	38,376		
<b>1994</b>	56,268		49,156
<b>1995</b>	28,393		
<b>1996</b>	24,196		
<b>1997</b>	18,282		37,664
<b>1998</b>	23,474		
<b>1999</b>	18,974		
<b>2000</b>	21,551		28,535
<b>2001</b>	31,120		
<b>2002</b>	25,213	18,645	49,035
<b>2003</b>	27,531		
<b>2004</b>	29,663	14,740	39,219
<b>2005</b>	46,084		
<b>2006</b>	61,644		45,369
<b>2007</b>	65,191		
<b>2008</b>	53,967	24,822	
<b>2009</b>	47,252		
<b>2010</b>	51,927	27,875	49,069

Table 6.5-3. ABC and OFL values calculated from 5 methods using running averages from 3 to 7 years.

**running averages for ABC calculation**

	7 yr	6 yr	5 yr	4 yr	3 yr
<b>1991</b>					
<b>1992</b>					
<b>1993</b>					13,166
<b>1994</b>				15,742	15,742
<b>1995</b>			14,919	14,919	14,919
<b>1996</b>		14,351	14,351	14,351	14,745
<b>1997</b>	13,660	13,660	13,660	13,783	12,272
<b>1998</b>	13,215	13,215	13,224	11,951	11,510
<b>1999</b>	12,800	12,739	11,624	11,211	10,734
<b>2000</b>	12,351	11,356	10,972	10,554	10,441
<b>2001</b>	11,589	11,309	11,041	11,078	11,107
<b>2002</b>	11,684	11,523	11,649	11,814	12,313
<b>2003</b>	11,770	11,916	12,102	12,548	13,392
<b>2004</b>	12,006	12,175	12,547	13,180	13,243
<b>2005</b>	12,754	13,160	13,789	13,988	14,007
<b>2006</b>	13,997	14,660	14,995	15,260	15,929
<b>2007</b>	15,399	15,800	16,174	16,904	18,357
<b>2008</b>	16,243	16,628	17,303	18,493	19,250
<b>2009</b>	16,842	17,440	18,419	18,968	18,951
<b>2010</b>	17,710	18,571	19,041	19,046	18,785

**running averages for OFL**

	7 yr	6 yr	5 yr	4 yr	3 yr
<b>1991</b>					
<b>1992</b>					
<b>1993</b>					17,555
<b>1994</b>				20,989	20,989
<b>1995</b>			19,892	19,892	19,892
<b>1996</b>		19,134	19,134	19,134	19,661
<b>1997</b>	18,213	18,213	18,213	18,377	16,362
<b>1998</b>	17,620	17,620	17,633	15,935	15,347
<b>1999</b>	17,067	16,986	15,499	14,948	14,311
<b>2000</b>	16,467	15,141	14,630	14,072	13,921
<b>2001</b>	15,453	15,078	14,722	14,770	14,809
<b>2002</b>	15,578	15,365	15,532	15,752	16,418
<b>2003</b>	15,694	15,888	16,135	16,731	17,856
<b>2004</b>	16,008	16,234	16,730	17,573	17,658
<b>2005</b>	17,005	17,547	18,385	18,651	18,675
<b>2006</b>	18,663	19,547	19,993	20,346	21,238
<b>2007</b>	20,532	21,067	21,565	22,538	24,476
<b>2008</b>	21,658	22,171	23,071	24,658	25,666
<b>2009</b>	22,456	23,253	24,559	25,291	25,268
<b>2010</b>	23,613	24,761	25,387	25,395	25,047

### Comparison of species identified during the EBS survey




Figure 6.5.1—Number of hauls where arrowtooth flounder and Kamchatka flounder were identified during the annual Bering Sea shelf surveys, 1982-2010.




Figure 6.5-2 Arrowtooth and Kamchatka flounder catch (t) by week from Alaska Regional Office catch reports.

**Legend**

**speciescpue2010.csv Events**

**wgtcpue**

- 0.198298 - 7.813200
- 7.813201 - 18.309061
- 18.309062 - 34.03601
- 34.036011 - 61.24088
- 61.240890 - 136.7693




Figure 6.5-3. Distribution and relative of abundance of Kamchatka flounder from the 2010 slope survey.




Figure 6.5-4. Distribution and relative abundance of Kamchatka flounder from the 2006 Aleutian Islands survey.




Figure 6.5-4 (continued).




Figure 6.5-4 (continued).






Figure 6.5-5 Kamchatka flounder length-weight plots for male and females.




Figure 6.5-6 Estimated ABC (t), by year, from five methods each using a different number of years to calculate a moving average from shelf, slope and Aleutian Islands biomass estimates.