

**Note: January 2004: There are minor differences between the numbers of Steller sea lions reported in Table 3 and the numbers provided to the Steller sea lion recovery team for central California and northern California/Oregon. These numbers will be updated in the draft SARs for 2004.**


### STELLER SEA LION (*Eumetopias jubatus*): Eastern U. S. Stock

#### STOCK DEFINITION AND GEOGRAPHIC RANGE

Steller sea lions range along the North Pacific Rim from northern Japan to California (Loughlin et al. 1984), with centers of abundance and distribution in the Gulf of Alaska and Aleutian Islands, respectively. The species is not known to migrate, but individuals disperse widely outside of the breeding season (late May-early July), thus potentially intermixing with animals from other areas. Despite the wide ranging movements of juveniles and adult males in particular, exchange between rookeries by breeding adult females and males (other than between adjoining rookeries) appears low (NMFS 1995); however, resighting data from branded animals have not yet been analyzed.

Loughlin (1997) considered the following information when classifying stock structure based upon the phylogeographic approach of Dizon et al. (1992): 1)

Distributional data: geographic distribution continuous, yet a high degree of natal site fidelity and low (<10%) exchange rate of breeding animals between rookeries; 2) Population response data: substantial differences in population dynamics (York et al. 1996); 3) Phenotypic data: unknown; and 4) Genotypic data: substantial differences in mitochondrial DNA (Bickham et al. 1996). Based on this information, two separate stocks of Steller sea lions are now recognized within U. S. waters: an eastern U. S. stock, which includes animals east of Cape Suckling, Alaska (144°W), and a western U. S. stock, which includes animals at and west of Cape Suckling (Loughlin 1997, Fig. 3).



**Figure 2.** Approximate distribution of Steller sea lions in the eastern U.S. stock (shaded area). Major haulouts and rookeries are also depicted (points). Note: Haulouts and rookeries in British Columbia are not shown.

#### POPULATION SIZE

The previous estimate of Steller sea lion abundance in Southeast Alaska was based on comprehensive aerial surveys performed in June 1996 (Sease et al. 1999, Sease and Loughlin 1999). Data from these surveys represent actual counts of pups and non-pups at all rookeries and major haulout sites in Southeast Alaska. In 1996 a total of 14,621 Steller sea lions were counted in Southeast Alaska, including 10,907 non-pups and 3,714 pups. Aerial surveys in 1998 and 2000 included the trend sites and other major sites. There were some differences between which major sites were surveyed in 1998 and 2000, so the total counts for each survey are not entirely comparable. The counts for 1998 and 2000 were 10,939 and 12,417, respectively (Sease and Loughlin, 1999, Sease et al, 2001). Pup counts totaled 4,160 in 1997 and 4,257 in 1998 (Sease and Loughlin, 1999). The total count for Southeast Alaska in 1998 is 15,196 (10,939 non-pups plus 4,257 pups); if we assume that the pup count is roughly stable, the total count for 2000 would be 16,674 (12,417 non-pups plus 4,257 pups).

Aerial surveys and ground counts of California, Oregon, and Washington rookeries and major haulout sites were also conducted during the summer of 1996 (NMFS unpubl. data, National Marine Mammal Laboratory, 7600 Sand Point Way NE, Seattle, WA 98115; Southwest Fisheries Science Center, P. O. Box 271, La Jolla, CA 90238; ODF&W unpubl. data, Marine Science Drive, Newport, OR 97365). In 1996 a total of 6,555 Steller sea lions were counted in California (2,042), Oregon (3,990), and Washington (523), including 5,464 non-pups and 1,091 pups.

The eastern U. S. stock of Steller sea lions is a transboundary stock, including sea lions from British Columbia

rookeries (see Wade and Angliss 1997 for discussion of transboundary stocks). Aerial surveys were last conducted in British Columbia during 1994 and produced counts of 8,091 non-pups and 1,186 pups, for a total count of 9,277 (Dept. Fisheries and Oceans, unpubl. data, Pacific Biological Station, Nanaimo, BC, V9R 5K6). Complete count data are not available for British Columbia in 1996. However, because the number of Steller sea lions in British Columbia is thought to have increased since 1994 (P. Olesiuk, pers. comm., Pacific Biological Station, Canada), the 1994 counts represent a conservative estimate for the 1996 counts. Combining the total counts for the three regions results in a minimum estimated abundance of 31,028 (15,196 + 6,555 + 9,277) Steller sea lions in this stock.

Slight changes in the non-pup numbers result from changes in the non-pup count database which occurred since publication of the results from the 1998 aerial survey (Sease and Loughlin 1999). The database underwent considerable review, verification, and editing; the most significant changes related to replicate counts of individual sites. For additional information on the minor changes in the non-pup numbers, see Sease et al. (2001).

The abundance estimate for the eastern U. S. stock is based on counts of all animals (pup and non-pup) at all sites and has not corrected for animals missed because they were at sea. A reliable correction factor to account for these animals is currently not available (J. Sease, pers. comm., National Marine Fisheries Service). As a result, this represents an underestimate for the total abundance of Steller sea lions in this stock.

### Minimum Population Estimate


The minimum population estimate will be calculated by adding 1998 counts from Southeast Alaska (15,196), 1996 counts from WA/OR/CA (6,555), and Canadian counts from 1994 (9,277), which results in an  $N_{MIN}$  for the eastern U. S. stock of Steller sea lions of 31,028. Recall that this count has not been corrected for animals which were at sea, and also uses the 1994 data from British Columbia where Steller sea lion numbers are thought to have increased since 1994.

### Current Population Trend

Trend counts (an index to examine population trends) for Steller sea lions in Oregon were relatively stable in the 1980s, with uncorrected counts in the range of 2,000-3,000 sea lions (NMFS 1992). Counts in Oregon have shown a gradual increase since 1976, as the adult and juvenile state-wide count for that year was 1,486 compared to 3,648 in 2001 (Brown and Reimer 1992; Brown et al. 2002).

Steller sea lion numbers in California, especially in southern and central California, have declined from historic numbers. Counts in California between 1927 and 1947 ranged between 5,000 and 7,000 non-pups with no apparent trend, but have subsequently declined by over 50%, remaining between 1,500 and 2,000 non-pups during 1980-01. Limited information suggests that counts in northern California appear to be stable (NMFS 1995). At Año Nuevo in central California, a steady decline in ground counts started around 1970, resulting in an 85% reduction in the breeding population by 1987 (LeBoeuf et al. 1991). In vertical aerial photographic counts conducted at Año Nuevo, pups declined at a rate of 9.9% from 1990 to 1993, while non-pups declined at a rate of 31.5% over the same time period (Westlake et al. 1997). Pup counts at Año Nuevo have been steadily declining at about 5% annually since 1990 (W. Perryman, pers. comm., National Marine Fisheries Service). The most recent pup counts at Año Nuevo and the Farallons are 564 for 1999 and 349 in 2000 (M. Lowry, pers. comm). Overall, counts of non-pups at trend sites in California and Oregon have been relatively stable since the 1980s (Table 3, Fig. 4).

In Southeast Alaska, counts (no correction factors applied) of non-pups at trend sites increased by 30% from 1979-2000 from 6,376 to 9,862 (Merrick et al., 1992, Sease et al., 2001). During 1979-97, counts of pups on the three



**Figure 4.** Counts of adult and juvenile Steller sea lions at rookery and haulout trend sites throughout the range of the eastern U.S. stock, 1982-002. Data from British Columbia include all sites.

rookeries in Southeast Alaska increased by an average of 5.9% per year. Since 1989 pup counts on the three rookeries increased at a lower rate (+1.7% per year) than for the entire period (Calkins et al. 1999). A slightly lower increase in pup counts (3.3% per year from 1979-97) is reported by Sease et al. (2001). In British Columbia, counts (no correction factors applied) of non-pups throughout the Province increased at a rate of 2.8% annually during 1971-98 (Table 3, Fig. 4; P. Olesiuk, pers. comm., Pacific Biological Station, Canada). Counts of non-pups at trend sites throughout the range of the eastern U. S. Steller sea lion stock are shown in Figure 4.

**Table 3.** Counts of adult and juvenile Steller sea lions observed at rookery and haulout trend sites by year and geographical area for the eastern U. S. stock from the 1982 through 2000 (NMFS 1995, Strick et al. 1997, Sease et al. 1999, Sease and Loughlin 1999; P. Olesiuk, unpubl. data, Pacific Biological Station, Nanaimo, BC, V9R 5K6; ODF&W unpubl. data, 7118 NE Vandenberg Ave., Corvallis, OR 97330; Point Reyes Bird Observatory, unpubl. data, 4990 Shoreline Hwy., Stinson Beach, CA 94970; Sease et al., 2001). Central California data include only Año Nuevo and Farallon Islands. Trend site counts in northern California/Oregon include St. George, Rogue, and Orford Reefs. British Columbia data include counts from all sites. [Note: There are minor differences between the numbers in Table 3 and the numbers provided to the Steller sea lion recovery team for central California and northern California/Oregon (italicized) . Revisions will be completed in 2004.]

Area	1982	1990	1991	1992	1994	1996	1998	2000	2002
<i>Central CA</i>	511 <sup>1</sup>	655	537	276	512	385	208	349	
<i>Northern CA/OR</i>	3,094	2,922	3,180	3,544	2,834	2,988	3,175	n/a	
British Columbia	4,711	6,109 <sup>2</sup>	no data	7,376	8,091	no data	9,818	n/a	n/a
Southeast Alaska	6,898	7,629	8,621	7,555	9,001	8,231	8,693	9,862	9,951
Total	15,214	--	--	18,754	20,263	--	21,864	n/a	n/a

<sup>1</sup> This count includes a 1983 count from Año Nuevo. <sup>2</sup> This count was conducted in 1987.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

There are no estimates of maximum net productivity rates for Steller sea lions. Hence, until additional data become available, it is recommended that the pinniped maximum theoretical net productivity rate ( $R_{MAX}$ ) of 12% be employed for this stock (Wade and Angliss 1997).

### POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor:  $PBR = N_{MIN} \times 0.5R_{MAX} \times F_R$ . The default recovery factor ( $F_R$ ) for stocks listed as “threatened” under the Endangered Species Act (ESA) is 0.5 (Wade and Angliss 1997). However, as total population estimates for the eastern U. S. stock have remained stable or increased over the last 20 years, the recovery factor is set at 0.75; midway between 0.5 (recovery factor for a “threatened” stock) and 1.0 (recovery factor for a stock within its optimal sustainable population level). This approach is consistent with recommendations of the Alaska Scientific Review Group. Thus, for the eastern U. S. stock of Steller sea lions,  $PBR = 1,396$  animals ( $31,028 \times 0.06 \times 0.75$ ).

### ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

#### Fisheries Information

Fishery observers monitored three commercial fisheries during the period from 1990 to 2001 in which Steller sea lions from this stock were taken incidentally: the California (CA)/Oregon (OR) thresher shark and swordfish drift

gillnet, WA/OR/CA groundfish trawl, and Northern Washington (WA) marine set gillnet fisheries. The best data available on the rates of serious injury and mortality incidental to these fisheries is presented in Table 4. There have been no observed serious injuries or mortalities incidental to the CA/OR thresher shark and swordfish drift gillnet fishery in recent years. Two and one Steller sea lions were observed taken in the WA/OR/CA groundfish trawl in 1997 and 2001, respectively; these observed takes in combination with a mortality that occurred in an unmonitored haul resulted in a mean estimated annual mortality level of 0.8 (Table 4). In 1996, one Steller sea lion mortality in the northern Washington marine set gillnet fishery was observed. The mortality was not extrapolated because the coastal portion of the fishery (the portion of the fishery most likely to interact with Steller sea lions) was monitored with 100% observer coverage during 1996. This single observed mortality results in a mean annual mortality of 0.2 (CV = 1.0) Steller sea lions for the Northern Washington marine set gillnet fishery. No observer program occurred during 1994 for this fishery, and no data available after 1998. These mortalities result in a mean annual mortality rate of 1.0 (CV = 1.0). No mortalities were reported by fishery observers monitoring drift gillnet and set gillnet fisheries in Washington and Oregon this decade; though, mortalities have been reported in the past.

**Table 4.** Summary of incidental mortality of Steller sea lions (eastern U. S. stock) due to commercial and tribal fisheries from 1990 to 2001 and calculation of the mean annual mortality rate. Mean annual mortality in brackets represents a minimum estimate from self-reported fisheries information or stranding data. Data from 1997 to 2001 (or the most recent 5 years of available data) are used in the mortality calculation when more than 5 years of data are provided for a particular fishery. n/a indicates that data are not available. \* indicates a mortality seen by an observer, but during an unmonitored haul; because the haul was not monitored, no extrapolation can be done.

Fishery name	Years	Data type	Range of observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
CA/OR thresher shark and swordfish drift gillnet	96-00	obs data	4-27%	0, 0, 0, 0, 0	0, 0, 0, 0, 0	0
WA/OR/CA groundfish trawl (Pacific whiting component)	97-01	obs data	66-96%	2, 0, 0, 0, 1	2, 0, 0, 1*, 1	0.8 (CV = n/a)
Northern WA marine set gillnet (tribal fishery)	94-98	obs data	47-98%	0, 0, 1, 0, 0	0, 0, 1, 0, 0	0.2 (CV = 1.0)
Observer program total						1.0 (CV = 1.0)
				<b>Reported mortalities</b>		
Southeast Alaska salmon drift gillnet	90-01	self reports	n/a	0, 1, 2, 2, n/a, n/a, n/a, n/a, n/a, n/a, n/a,	n/a	[≥1.25]
Alaska salmon troll	92-01	strand data	n/a	0, 0, 0, 1, 0, 0, n/a, n/a, n/a	n/a	[≥0.2]

Fishery name	Years	Data type	Range of observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
British Columbia aquaculture predator control program	91-01	permit reports	n/a	14, 8, 10, 11, 6, 13, 34, 63, 91, n/a, n/a, n/a	n/a	41.4
Minimum total annual incidental mortality (estimate from observer programs plus estimates from self reports and stranding data; includes an estimate of 1.2 fishery-related strandings per year from 1996-00; see text)						3.65 (CV = 1.0)
Minimum total annual mortality (includes intentional mortalities in the BC predator control program)						45.25 (CV = 1.0)

An additional source of information on the number of Steller sea lions killed or injured incidental to commercial fishery operations is the self-reported fisheries information required of vessel operators by the MMPA. During the period between 1990 and 1998, fisher self-reports from the Southeast Alaska salmon drift gillnet fishery (Table 4) resulted in an annual mean of 1.25 mortalities from interactions with commercial fishing gear. This total is based on all available fisher self-reports for U. S. fisheries within the range of the stock, except the three fisheries for which observer data were presented above. However, because logbook records (fisher self-reports required during 1990-94) are most likely negatively biased (Credle et al. 1994), these are considered to be minimum estimates. During 1990, 11 Steller sea lion injuries incidental to the Alaska salmon troll fishery and 1 Steller sea lion injury incidental to the CA/OR/WA salmon troll fishery were reported. These injuries were not deemed serious (Angliss and DeMaster 1998) and have not been included in the Table 4. Logbook data are available for part of 1989-1994, after which incidental mortality reporting requirements were modified. Under the new system, logbooks are no longer required; instead, fishers provide self-reports. Data for the 1994-95 phase-in period is fragmentary. After 1995, the level of reporting dropped dramatically, such that the records are considered incomplete and estimates of mortality based on them represent minimums (see Appendix 7 for details).

Strandings of Steller sea lions entangled in fishing gear or with injuries caused by interactions with gear are another source of mortality data. During the 5-year period from 1995 to 1999 there were 4 fishery-related strandings in Southeast Alaska. One of these strandings has been attributed to the Alaska salmon troll fishery and has been included in Table 4. Details regarding which fishery may be responsible for other fishery-related strandings between 1994-99 is not available at this time. In 2000, there were reports of 3 Steller sea lions observed in southeast Alaska with “flashers” lodged in their mouths and one animal entangled in fishing line; all animals were alive when seen. It is not clear whether these entanglements resulted from the commercial or recreational fisheries, nor is it clear whether the interactions resulted in mortality. However, based on Angliss and DeMaster (1998), it would be appropriate to call these “serious injuries”. During the 5-year period from 1996-00, there were 6 fishery-related strandings; this results in an estimated annual mortality of 1.2 animals from this stock. This estimate is considered a minimum because not all entangled animals strand and not all stranded animals are found or reported.

Due to limited observer program coverage, no data exist on the mortality of marine mammals incidental to Canadian commercial fisheries (i.e., those similar to U.S. fisheries known to take Steller sea lions). As a result, the number of Steller sea lions taken in Canadian waters is not known.

The minimum estimated mortality rate incidental to commercial fisheries (both U.S. and Canadian) is 3.65 sea lions per year, based on observer data (1.0), self-reported fisheries information (1.25), and stranding data (0.2 + 1.2 = 1.4).

### Subsistence/Native Harvest Information

The subsistence harvest of Steller sea lions during 1997-01 is summarized in Wolfe et al. (2002). During each year, data were collected through systematic interviews with hunters and users of marine mammals in approximately 2,100 households in about 60 coastal communities within the geographic range of the Steller sea lion in Alaska.

Approximately 16 of the interviewed communities lie within the range of the eastern U. S. stock. The average number of animals harvested and struck but lost is 2 animals/year.

An unknown number of Steller sea lions from this stock are harvested by subsistence hunters in Canada. The magnitude of the Canadian subsistence harvest is believed to be small. Alaska Native subsistence hunters have initiated discussions with Canadian hunters to quantify their respective subsistence harvests, and to identify any effect these harvests may have on the cooperative management process.

### **Other Mortality**

Illegal shooting of sea lions in U.S. waters was thought to be a potentially significant source of mortality prior to the listing of sea lions as “threatened” under the ESA in 1990. Such shooting has been illegal since the species was listed as threatened. (Note: the 1994 Amendments to the MMPA made intentional lethal take of any marine mammal illegal except for subsistence hunting by Alaska Natives or where imminently necessary to protect human life). Records from NMFS enforcement indicate that there were 2 cases of illegal shootings of Steller sea lions in Southeast Alaska between 1995 and 1999: the cases involved the illegal shooting of one Steller sea lion near Sitka in 1998, and 3 Steller sea lions in Petersburg. Both cases were successfully prosecuted (NMFS, Alaska Enforcement Division).

Steller sea lions are taken in British Columbia during commercial salmon farming operations (Table 4). Preliminary figures from the British Columbia Aquaculture Predator Control Program indicated a mean annual mortality of 44 Steller sea lions from this stock over the period from 1995 to 1999 (P. Olesiuk, pers. comm., Pacific Biological Station, Canada). Note that the 1995 estimate includes one animal reported as an unidentified sea lion and the 1996 estimate is based on data from only the first three-quarters of 1996. The take of Steller sea lions has increased in recent years because of recent changes in sea lion distribution which have likely occurred in response to a shift in herring distribution (P. Olesiuk, pers. comm).

Strandings of Steller sea lions with gunshot wounds do still occur, along with strandings of animals entangled in gear that is not fishery-related. During the period from 1996 to 1999 human-related strandings of animals with gunshot wounds from this stock occurred in Oregon, Washington, and Alaska in 1996 (2 animals), 1997 (3 animals), 1998 (1 animal), and 1999 (2 animals), resulting in an estimated annual mortality of 2.0 Steller sea lions from this stock during 1996-99. This estimate is considered a minimum because not all stranded animals are found, reported, or cause of death determined (via necropsy by trained personnel). In addition, human-related stranding data are not available for British Columbia. Reports of stranded animals in Alaska with gunshot wounds have been included in the above estimates. However, it is not possible to tell whether the animal was illegally shot or if the animal was struck and lost by subsistence hunters (in which case the mortality would have been legal and accounted for in the subsistence harvest estimate). However, one of the two 1996 reports was from Alaska and has been included because there were no subsistence struck and lost reports during that year.

Stranding data may also provide information on additional sources of potential mortality. In 2000, 4 Steller sea lions were sighted entangled in some kind of rope or line that was not necessarily related to a commercial or recreational fishery, and one animal was seen entangled in a 14" tire. All of these animals were alive when sighted; the animal entangled in the tire was successfully released. It is not clear whether the occurrence of these interactions in stranding data in 2000 but not in previous years reflects an increase in these types of interactions or an increase in reporting. If the number of interactions is averaged over 5 years, the “other” interaction rate would be a minimum of one animal per year.

### **STATUS OF STOCK**

Based on currently available data, the minimum estimated fishery mortality and serious injury for this stock ( $0.7 + 1.25 + 0.2 + 41.4 + 1.2 = 45.5$ ) is less than that 10% of the calculated PBR (140) and, therefore, can be considered to be insignificant and approaching a zero mortality and serious injury rate. The estimated annual level of total human-caused mortality and serious injury from fishery interactions, subsistence harvests, and shootings ( $44 + 0 + 2 = 46$ ) does not exceed the PBR (1,396) for this stock. The eastern U. S. stock of Steller sea lion is currently listed as “threatened” under the ESA, and therefore designated as “depleted” under the MMPA. As a result, this stock is classified as a strategic stock. Although the stock size has increased in recent years, the status of this stock relative to its Optimum Sustainable Population size is unknown.

### Habitat Concerns

Unlike the observed decline in the western U. S. stock of Steller sea lion there has not been a concomitant decline in the eastern U. S. stock. Concerns regarding the possible impacts of commercial groundfish fisheries in the Gulf of Alaska and Bering Sea have been noted previously (see Habitat Concerns section in assessment report for the western U. S. stock). However, the eastern U. S. stock is stable or increasing in the northern portion of its range (Southeast Alaska and British Columbia). The stock has been declining in the southern end of its range (see Current Population Trend), where habitat concerns include reduced prey availability, contaminants, and disease (Sydeman and Allen 1997).

### CITATIONS

- Angliss, R. P., and D. P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: report of the serious injury workshop 1-2 April 1997, Silver Spring, Maryland. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-13, 48 pp.
- Bickham, J. W., J. C. Patton, and T. R. Loughlin. 1996. High variability for control-region sequences in a marine mammal: Implications for conservation and biogeography of Steller sea lions (*Eumetopias jubatus*). *J. Mammal.* 77:95-108.
- Brown, R. F., and S. D. Reimer. 1992. Steller sea lion counts in Oregon during June and July, 1975-1991. In-house Rep., Nongame Wildlife Prog., Oregon Dept. Fish and Wildl., Newport, OR, 97365. 12 pp.
- Calkins, D. G., D. C. McAllister, K. W. Pitcher, and G. W. Pendelton. 1999. Steller sea lion status and trend in Southeast Alaska: 1979-1997. *Mar. Mammal Sci.* 15(2):462-477.
- Credle, V. R., D. P. DeMaster, M. M. Merklein, M. B. Hanson, W. A. Karp, and S. M. Fitzgerald (eds.). 1994. NMFS observer programs: minutes and recommendations from a workshop held in Galveston, Texas, November 10-11, 1993. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-94-1, 96 pp.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Julian, F. 1997. Cetacean mortality in California gillnet fisheries: preliminary estimates for 1996. Unpubl. doc. submitted to Int. Whal. Comm. (SC/49/SM). 13 pp.
- Julian, F., and M. Beeson. 1998. Estimates of marine mammal, turtle, and seabird mortality for two California gillnet fisheries: 1990-1995. *Fish. Bull.* 96:271-284.
- LeBoeuf, B. J., K. Ono, and J. Reiter. 1991. History of the Steller sea lion population at Año Nuevo Island, 1961-1991. Southwest Fish. Sci. Center Admin. Rep. LJ-91-45C. U.S. Department of Commerce, La Jolla, CA, 9p + tables +figs.
- Loughlin, T. R. 1997. Using the phylogeographic method to identify Steller sea lion stocks. Pp. 329-341, *In* A. Dizon, S. J. Chivers, and W. Perrin (eds.), *Molecular genetics of marine mammals, incorporating the proceedings of a workshop on the analysis of genetic data to address problems of stock identity as related to management of marine mammals.* Soc. Mar. Mammal., Spec. Rep. No. 3.
- Loughlin, T. R., D. J. Rugh, and C. H. Fiscus. 1984. Northern sea lion distribution and abundance: 1956-1980. *J. Wildl. Manage.* 48:729-740.
- Merrick, R.L., D.G. Calkins, and D.C. McAllister. 1992. Aerial and ship-based surveys of Steller sea lions in Southeast Alaska, the Gulf of Alaska, and Aleutian Islands during June and July 1991. U.S. Dep. Commer, NOAA Tech Memo. NMFS-AFSC-1. 37p.
- National Marine Fisheries Service. 1992. Recovery Plan for the Steller Sea Lion (*Eumetopias jubatus*). Prepared by the Steller Sea Lion Recovery Team for the National Marine Fisheries Service, Silver Spring, MD. 92 pp.
- National Marine Fisheries Service. 1995. Status review of the United States Steller sea lion (*Eumetopias jubatus*) population. Prepared by the National Marine Mammal Laboratory, AFSC, NMFS, NOAA, 7600 Sand Point Way NE, Seattle, WA 98115. 61 pp.
- Sease, J. L., and T. R. Loughlin. 1999. Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1997 and 1998. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-100, 61 pp.
- Sease, J. L., J. M. Strick, R. L. Merrick, and J. P. Lewis. 1999. Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1996. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-99, 43 pp.

- Sease, J. L., W. P. Taylor, T. R. Loughlin, and K. W. Pitcher. 2001. Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1999 and 2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-122, 52p.
- Strick, J. M., L. W. Fritz, and J. P. Lewis. 1997. Aerial and ship-based surveys of Steller sea lions (*Eumetopias jubatus*) in Southeast Alaska, the Gulf of Alaska, and Aleutian Islands during June and July 1994. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-71, 55 pp.
- Sydeman, W. J., and S. G. Allen. 1997. Trends and oceanographic correlates of pinniped populations in the Gulf of the Farallones, California. U.S. Dep. Commer., Southwest Fish. Sci. Cent., Admin. Rep. LJ-97-02C, 28 pp.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Westlake, R. L., W. L. Perryman, and K. A. Ono. 1997. Comparison of vertical aerial photographic and ground censuses of Steller sea lions at Año Nuevo Island, July 1990-1993. *Mar. Mammal Sci.* 13(2): 207-218.
- Wolfe, R. J., J. A. Fall, and R. T. Stanek. 2002. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2001. Prepared for NMFS by Alaska Dept. Fish and Game, Juneau, Alaska, 90p + appendices.
- York, A. E., R. L. Merrick, and T. R. Loughlin. 1996. An analysis of the Steller sea lion metapopulation in Alaska. Chapter 12, Pp. 259-292, *In* D. R. McCullough (ed.), *Metapopulations and wildlife conservation*. Island Press, Covelo, California.