

HARBOR SEAL (*Phoca vitulina richardsi*): Bering Sea Stock

NOTE – January 2009: NMFS has new genetic information on harbor seals in Alaska which indicates that the current division of Alaskan harbor seals into the Southeast Alaska, Gulf of Alaska, and Bering Sea stocks needs to be reassessed. NMFS, in cooperation with our partners in the Alaskan Native community, is evaluating the new genetic information and hopes to make a joint recommendation regarding stock structure in 2009. In the interim, new information on harbor seal mortality levels is provided within this report. A complete revision of the harbor seal stock assessments will be postponed until new stocks are defined.

STOCK DEFINITION AND GEOGRAPHIC RANGE

Harbor seals inhabit coastal and estuarine waters off Baja California, north along the western coasts of the United States, British Columbia, and Southeast Alaska, west through the Gulf of Alaska and Aleutian Islands, and in the Bering Sea north to Cape Newenham and the Pribilof Islands. They haul out on rocks, reefs, beaches, and drifting glacial ice, and feed in marine, estuarine, and occasionally fresh waters. Harbor seals generally are non-migratory, with local movements associated with such factors as tides, weather, season, food availability, and reproduction (Scheffer and Slipp 1944; Fisher 1952; Bigg 1969, 1981). The results of recent satellite tagging studies in Southeast Alaska, Prince William Sound, and Kodiak are also consistent with the conclusion that harbor seals are non-migratory (Swain et al. 1996, Lowry et al. 2001, Small et al. 2001). However, some long-distance movements of tagged animals in Alaska have been recorded (Pitcher and McAllister 1981, Lowry et al. 2001, Small et al. 2001). Strong fidelity of individuals for haulout sites during the breeding season has been documented in several populations (Härkönen and Harding 2001), including in Alaska (Pitcher and Calkins 1979, Pitcher and McAllister 1981).




Figure 10. Approximate distribution of harbor seals in Alaska waters (shaded area).

Westlake and O’Corry-Crowe’s (2002) analysis of genetic information revealed population subdivisions on a scale of 600-820 km. These results suggest that genetic differences within Alaska, and most likely over their entire North Pacific range, increase with increasing geographic distance. New information revealed substantial genetic differences indicating that female dispersal occurs at region specific spatial scales of 150-540 km. This research identified 12 demographically independent clusters within the range of Alaskan harbor seals; however additional research is required as unsampled areas within the Alaskan harbor seal range remain (O’Corry-Crowe et al. 2003).

Currently there are three stocks of harbor seals identified in Alaska: 1) the Southeast Alaska stock - occurring from the Alaska/British Columbia border to Cape Suckling, Alaska (144°W), 2) the Gulf of Alaska stock - occurring from Cape Suckling to Unimak Pass, including animals throughout the Aleutian Islands, and 3) the Bering Sea stock - including all waters north of Unimak Pass (Fig. 8). Information concerning the three harbor seal stocks recognized along the West Coast of the continental United States can be found in the Stock Assessment Reports for the Pacific Region.

POPULATION SIZE

The National Marine Mammal Laboratory (Alaska Fisheries Science Center) routinely conducts aerial surveys of harbor seals across their entire range in Alaska. Each of five survey regions was surveyed, with one region surveyed per year. To derive an accurate estimate of population size from these surveys, a method was

developed to address the influence of external conditions on the number of seals hauled out on shore, and counted, during the surveys. Many factors influence the propensity of seals to haul out, including tides, weather, time of day, and date in the seals' annual life history cycle. A statistical model defining the relationship between these factors and the number of seals hauled out was developed for each survey region. Based on those models, the survey counts for each year were adjusted to the number of seals that would have been ashore during a hypothetical survey conducted under ideal conditions for hauling out (Boveng et al. 2003). In a separate analysis of radio-tagged seals, a similar statistical model was used to estimate the proportion of seals that were hauled out under those ideal conditions (Simpkins et al. 2003). The results from these two analyses were combined for each region to estimate the population size of harbor seals in Alaska. Discussions of estimates from a previous survey (1995) can be found in earlier stock assessment reports.

The current statewide abundance estimate for Alaskan harbor seals is 180,017 (CV = 0.03; NMFS, unpublished data), based on data collected during 1996-2000. This estimate, however, is believed to be low because it is based on incomplete coverage of terrestrial sites in Prince William Sound and of glacial sites in the Gulf of Alaska and the Southeast Alaska regions. Those problems have been addressed in the current survey (2001-2005). Prince William Sound was surveyed completely in 2001, and new methods have been developed and used for surveying glacial sites in 2001-2002. Analyses are currently underway, and a manuscript describing the regional and statewide population estimates is in preparation; the analytical methods are described in Boveng et al. (2003) and Simpkins et al. (2003) and have been presented at the 14th Biennial Conference on the Biology of Marine Mammals. The current abundance estimate for the Bering Sea stock (21,651; $18,073 \times 1.198$; CV = 0.1) is calculated from surveys in 2000 (NMFS, unpublished data).

Minimum Population Estimate

The minimum population estimate (N_{MIN}) for this stock is calculated using Equation 1 from the PBR Guidelines (Wade and Angliss 1997): $N_{\text{MIN}} = N/\exp(0.842 \times [\ln(1 + [\text{CV}(N)]^2)]^{1/2})$. Using the population estimate (N) of 21,651 from the aerial surveys and the associated CV(N) of 0.1, results in an estimate of 19,907 harbor seals. Adding the maximum count of 202 seals from the Otter Island survey results in an N_{MIN} of 20,109 for the Bering Sea harbor seal stock.

Current Population Trend

The number of harbor seals in the Bering Sea stock is thought to have declined between the 1980s and 1990s (Alaska SRG, see DeMaster 1996); however, published data to support this conclusion are unavailable. Specifically, in 1974 there were 1,175 seals reported on Otter Island. The maximum count in 1995 (202 seals) represents an 83% decline (Withrow and Loughlin 1996). However, as noted by the Alaska SRG (DeMaster 1996), the reason(s) for this decline is(are) confounded by the recolonization of Otter Island by northern fur seals since 1974, which has caused a loss of available habitat for harbor seals. Further, counts of harbor seals on the north side of the Alaska Peninsula in 1995 were less than 42% of the 1975 counts, representing a decline of 3.5% per year. The number of harbor seals in northern Bristol Bay are also lower, but have remained stable since 1990 (Withrow and Loughlin 1996). Trend counts have been conducted in Bristol Bay only between 1998 and 2001. During this period, counts indicated a non-significant trend of -1.3% (95% CI: -5.9 - 3.3; Small et al. 2003). Calculation of trends in abundance in this area is somewhat problematic due to the presence of a sympatric species, spotted seals, which may overlap the range of harbor seals but cannot be identified as a different species by aerial surveys.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Reliable rates of maximum net productivity have not been estimated for the Gulf of Alaska or Bering Sea stock of harbor seal. Population growth rates were estimated at 6% and 8% between 1991 and 1992 in Oregon and Washington, respectively (Huber et al. 1994). Harbor seals have been protected in British Columbia since 1970, and the population has responded with an annual rate of increase of approximately 12.5% since 1973 (Olesiuk et al. 1990). However, until additional data become available from which more reliable estimates of population growth can be determined, it is recommended that the pinniped maximum theoretical net productivity rate (R_{MAX}) of 12% be employed for this stock (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $\text{PBR} = N_{\text{MIN}} \times 0.5R_{\text{MAX}} \times F_R$. The recovery factor (F_R) for this stock is 0.5,

the value for pinniped stocks with unknown population status (Wade and Angliss 1997). Thus, for the Bering Sea harbor seal stock, PBR = 603 animals ($20,109 \times 0.06 \times 0.5$).

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fisheries Information

The previous stock assessment for harbor seals indicated that there were three observed commercial fisheries that operated within the range of the Bering Sea stock of harbor seals. As of 2003, changes in how fisheries are defined in the List of Fisheries have resulted in separating these fisheries into 14 fisheries based on both gear type and target species (69 FR 70094, 2 December 2004). This change does not represent a change in fishing effort, but provides managers with better information on the component of each fishery that is responsible for the incidental serious injury or mortality of marine mammal stocks in Alaska.

Observer programs in several fisheries have documented mortalities or serious injuries in the Bering Sea/Aleutian Islands flatfish trawl and the Bering Sea/Aleutian Islands Pacific cod trawl (Table 13). Over the last 5 years, there were no observed serious injuries or mortalities of harbor seals in any Bering Sea/Aleutian Islands groundfish longline fisheries, or any Bering Sea/Aleutian Islands finfish pot fisheries (Perez 2006, Perez unpubl. ms).

The estimated minimum annual mortality rate incidental to commercial fisheries for the period 2002-2006 is 2.9. However, a reliable estimate of the mortality rate incidental to commercial fisheries is currently unavailable because of the absence of observer placements in salmon gillnet fisheries known to interact with this stock. More current data on estimated fishery-related serious injury and mortality are being analyzed and will be available for inclusion in the 2010 SARs.

Table 13. Summary of incidental mortality of harbor seals (Bering Sea stock) due to commercial fisheries from 2002 through 2006 and calculation of the mean annual mortality rate.

Fishery name	Years	Data type	Range of observer coverage (%)	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Bering Sea/ Aleutian Islands Pacific cod trawl	2002	obs	38.3	0	0	1.33 (CV = 0.44)
	2003	data	42.3	1	2.0	
	2004		45.3	1	2.0	
	2005		52.8	0	0	
	2006		46.8	1	2.7	
Bering Sea/ Aleutian Islands flatfish trawl	2002	obs	58.4	0	0	1.31 (CV = 0.34)
	2003	data	64.1	0	0	
	2004		64.3	0	1.0	
	2005		68.3	2	3.0	
	2006		67.8	1	2.6	
Bering Sea/ Aleutian Islands pollock trawl	2002	obs	80.0	0	0	0.29 (CV = 0.56)
	2003	data	82.2	0	0	
	2004		81.2	0	0	
	2005		77.3	1	1.5	
	2006		73.0	0	0	
Minimum total annual mortality						2.93 (CV = 0.26)

Subsistence/Native Harvest Information

The Alaska Native subsistence harvest of harbor seals has been estimated by the Alaska Native Harbor Seal Commission (ANHSC) and the Alaska Department of Fish and Game (ADFG). The previous stock assessment reported that the estimated average harvest of the Bering Sea stock of harbor seals for 1994-1996 was 161 animals per year (including struck and lost). Recent information from the ADFG indicates the average harvest level from 2002-2006, including struck and lost, was 106.5 animals per year. Due to seasonal geographic overlap in species distribution in north Bristol Bay in combination with the difficulty in distinguishing the two species from external morphology, reports of harvest levels of harbor seals were differentiated from spotted seals based on ecological

features of the kill, primarily degree of association with seasonal ice (Wolfe et al. 2008). The estimates given in Table 14 represent the best estimate of the subsistence harvest of harbor seals, although species identifications were not confirmed; therefore, the harvest estimates for harbor seals may include some spotted seals, and some spotted seals may actually have been recorded as harbor seals (Wolfe et al. 2008).

Table 14 provides a summary of the subsistence harvest information for the Bering Sea stock. Takes from the Bering Sea stock have decreased about 63.5%, declining from 243 seals in 1992 to 88 seals in 2006 (Wolfe et al. 2008).

Table 14. Summary of the subsistence harvest data for the Bering Sea stock of harbor seals, 2003-2007. Data are from Wolfe et al. 2004; Wolfe et al. 2006; Wolfe et al. 2008; J. Fall, ADFG, pers. comm., 04 February 2009.

Year	Estimated total number taken	Number harvested	Number struck and lost
2003	82	65	17
2004	119	76	43
2005	104	64	40
2006	88	64	24
2007	88	61	27
Mean annual harvest (2002-2007)	96	66	30

Other Mortality

Illegal intentional killing of harbor seals occurs, but the magnitude of this mortality is unknown (Note: the 1994 Amendments to the MMPA made intentional lethal take of any marine mammal illegal except where imminently necessary to protect human life). The Alaska Region stranding records from 1998 to 2002 document 2-3 reports of stranded harbor seals found shot in Bristol Bay, for a maximum average of 0.6 harbor seals/year over 5 years. It is not known whether these animals were killed illegally or if they were struck but lost in the subsistence harvest. Because the reason for the shooting is not known, these animals are added to the total number of human-related mortalities.

Mortalities may occasionally occur incidental to marine mammal research activities authorized under MMPA permits issued to a variety of government, academic, and other research organizations. Between 2003-2007, there were no mortalities resulting from research on the Bering Sea stock of harbor seals (Tammy Adams, Permits, Conservation, and Education Division, Office of Protected Resources, NMFS, 1315 East-West Highway, Silver Spring, MD 20910).

STATUS OF STOCK

Harbor seals are not listed as “depleted” under the MMPA or listed as “threatened” or “endangered” under the Endangered Species Act. At present, U.S. commercial fishery-related annual mortality levels less than 60 animals per year (i.e., 10% of PBR) can be considered insignificant and approaching zero mortality and serious injury rate. A reliable estimate of the annual rate of mortality incidental to commercial fisheries is unavailable. Therefore, it is unknown whether the kill rate due to commercial fishing is insignificant. Based on the best scientific information available, the estimated level of human-caused mortality and serious injury ($2.93 + 96 + 0.6 = 99.5$) is not known to exceed the PBR (603). Therefore, the Bering Sea stock of harbor seals is not classified as a strategic stock. The status of this stock relative to its Optimum Sustainable Population size is unknown.

CITATIONS

- Bigg, M. A. 1969. The harbour seal in British Columbia. Fish. Res. Bd. Can. Bull. 172. 33 pp.
- Bigg, M. A. 1981. Harbour seal, *Phoca vitulina*, Linnaeus, 1758 and *Phoca largha*, Pallas, 1811. Pp. 1-27 In S. H. Ridgway and R. J. Harrison (eds.), Handbook of Marine Mammals, vol. 2: Seals. Academic Press, New York.
- Boveng, P. L., J. L. Bengtson, D. E. Withrow, J. C. Cesarone, M. A. Simpkins, K. J. Frost and J. J. Burns. 2003. The abundance of harbor seals in the Gulf of Alaska. Mar. Mammal Sci. 19(1):111-127.
- DeMaster, D. P. 1996. Minutes from the 11-13 September 1996 meeting of the Alaska Scientific Review Group, Anchorage, Alaska. 20 pp. + appendices. (available upon request - National Marine Mammal Laboratory, 7600 Sand Point Way, NE, Seattle, WA 98115).

- Fisher, H. D. 1952. The status of the harbour seal in British Columbia, with particular reference to the Skeena River. Fish. Res. Bd. Can. Bull. 93. 58 pp.
- Härkönen, T., and K. C. Harding. 2001. Spatial structure of harbour seal populations and the implications thereof. Canadian Journal of Zoology 79:2115-2127.
- Huber, H., S. Jeffries, R. Brown, and R. DeLong. 1994. Harbor Seal Stock Assessment in Washington and Oregon 1993. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Lowry, L. F., K. J. Frost, J. M. VerHoef, and R. A. DeLong. 2001. Movements of satellite-tagged subadult and adult harbor seals in Prince William Sound, Alaska. Mar. Mammal Sci. 17:835-861.
- O’Corry-Crowe, G. M., Martien, K. K., and B. L. Taylor. 2003. The analysis of population genetic structure in Alaskan harbor seals, *Phoca vitulina*, as a framework for the identification of management stocks. Administrative Report LJ-03-08, Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA, 8604 La Jolla Shores Dr., La Jolla, CA 92037.
- Olesiuk, P. F., M. A. Bigg, and G. M. Ellis. 1990. Recent trends in the abundance of harbour seals, *Phoca vitulina*, in British Columbia. Can. J. Fish. Aquat. Sci. 47:992-1003.
- Perez, M. A. 2006. Analysis of marine mammal bycatch data from the trawl, longline, and pot groundfish fisheries of Alaska, 1998-2004, defined by geographic area, gear type, and target groundfish catch species. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-167.
- Perez, M. A. Unpubl. ms. Bycatch of marine mammals by the groundfish fisheries in the U.S. EEZ of Alaska, 2005. 67 pp. Available NMML-AFSC.
- Pitcher, K. W., and D. G. Calkins. 1979. Biology of the harbor seal (*Phoca vitulina richardsi*) in the Gulf of Alaska. U.S. Dep. Commer., NOAA, OCSEAP Final Rep. 19(1983):231-310.
- Pitcher, K. W., and D. C. McAllister. 1981. Movements and haul out behavior of radio-tagged harbor seals, *Phoca vitulina*. Can. Field Nat. 95:292-297.
- Scheffer, V. B., and J. W. Slipp. 1944. The harbor seal in Washington state. Amer. Midl. Nat. 32:373-416.
- Simpkins, M. A., D. E. Withrow, J. C. Cesarone and P. L. Boveng. 2003. Stability in the proportion of harbor seals hauled out under locally ideal conditions. Mar. Mammal Sci. 19(4):791-805.
- Small, R. J., and D. P. DeMaster. 1995. Alaska marine mammal stock assessments 1995. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-57, 93 pp.
- Small, R. J., G. W. Pendleton, and K. W. Pitcher. 2003. Trends in abundance of Alaska harbor seals, 1983-2001. Mar. Mammal Sci. 19(2):344-362.
- Small, R. J., L. F. Lowry, K. J. Frost, J. M. VerHoef, R. A. DeLong, and M. J. Rehberg. 2001. Movements of satellite-tagged harbor seal pups in Prince William Sound and the Gulf of Alaska. Mar. Mammal Sci. 17:835-861.
- Swain, U., J. Lewis, G. Pendleton, and K. Pitcher. 1996. Movements, haulout, and diving behavior of harbor seals in southeast Alaska and Kodiak Island. Pp. 59-144, *In* Annual Report: Harbor seal investigations in Alaska. NOAA Grant NA57FX0367. Alaska Dep. Fish and Game, Division of Wildlife Conservation. Douglas, AK.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Westlake, R. L., and G. O’Corry-Crowe. 2002. Macrogeographic structure and patterns of genetic diversity in harbor seals (*Phoca vitulina*) from Alaska to Japan. J. Mamm. 83(4):1111-1126.
- Withrow, D. E., and T. R. Loughlin. 1996. Abundance and distribution of harbor seals (*Phoca vitulina richardsi*) along the north side of the Alaska Peninsula and Bristol Bay during 1995. Annual report to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910.
- Wolfe, R. J., J. A. Fall, and M. Riedel. 2008. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2006. Alaska Dep. Fish and Game, Division of Subsistence Technical Paper No. 339. Juneau, AK.
- Wolfe, R. J., J. A. Fall, and R. T. Stanek. 2004. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2003. Alaska Dep. Fish and Game, Division of Subsistence Technical Paper No. 291. Juneau, AK.

Wolfe, R. J., J. A. Fall, and R. T. Stanek. 2006. The subsistence harvest of harbor seals and sea lions by Alaska Natives in 2005. Alaska Dep. Fish and Game, Division of Subsistence Technical Paper No. 319. Juneau, AK.